arXiv:gr-qc/0510127v1 31 Oct 2005

Imprints of Spacetime Topology in the Hawking-Unruh Effect

by Paul Pierre Langlois, MMath
Division of Applied Mathematics

School of Mathematical Sciences
University of Nottingham

Thesis submitted to the University of Nottingham
for the Degree of Doctor of Philosophy, August 2005

Abstract

The Unruh and Hawking effects are investigated on certain families of topologically nontrivial spacetimes using a variety of techniques. The thesis is split into three main parts. The first presents the Bogolubov transformation between Rindler and Minkowski quantizations on two flat spacetimes with topology $\mathbb{R}^3 \times S^1$ (denoted M_0 and M_-) for massive Dirac spinors. The two inequivalent spin structures on each spacetime are considered. Results show modifications to the Minkowski space Unruh effect. This provides a flat space model for the Hawking effect on the Kruskal and $\mathbb{RP}^3$ geon black hole spacetimes which is the subject of the rest of this part.

The second part presents the expectation values of the stress tensor for massive scalar and spinor fields on M_0 and M_-, and for massive scalar fields on Minkowski space with a single infinite plane boundary, in the Minkowski-like vacua.

The final part investigates particle detector models. The aim is two-fold. First we investigate a recent paper by Schlicht which highlights the importance of choosing a suitable regularization when calculating the response of monopole detectors in a causal way. We extend Schlicht's regularization to an arbitrary spacetime dimension, to quotient spaces of Minkowski space, to non-linear couplings of the scalar field, to a massless Dirac field, and to conformally flat spacetimes. Secondly we present some detector responses. Our main new results are the time dependent responses of inertial and uniformly accelerated detectors on M_- and M with boundary with motion perpendicular to the boundary. Responses are also considered for static observers in the exterior of the $\mathbb{RP}^3$ geon and comoving observers in $\mathbb{RP}^3$ de Sitter space, via those in the associated global embedding Minkowski spaces (GEMS). The response on $\mathbb{RP}^3$ de Sitter space, found both directly and in its GEMS, provides support for the validity of the GEMS procedure applied to detector responses.

Acknowledgements

Firstly I would like to thank my main thesis supervisor Dr Jorma Louko for suggesting to me these interesting topics, for many interesting and stimulating discussions and for reading this manuscript. Thanks also to Professor Edward Armour as a secondary supervisor and for many useful discussions.

I would like to thank the University of Nottingham for a University of Nottingham postgraduate studentship and the States of Guernsey for additional funding during my writing up period.

Thanks of course go to my parents for all their support in the past and for never discouraging me from doing anything I choose to do.

Thanks to all my friends in Nottingham and all over the world who have helped make my life away from my little Island both stimulating and very enjoyable, in particular thanks to all those associated with BSTN, you know who you are, and to Elaine and Eadie in Taiwan.

Finally I would like the thank Belinda for all the love and care she has shown towards me and for making my life complete.

Examiners

Bernard Kay
John Barrett

Contents

Acknowledgements **ii**

1 Introduction **1**

2 Unruh effect for massive spinors **9**
- 2.1 Introduction . . . 9
- 2.2 The spacetimes M_0 and M_- . . . 10
- 2.3 Spinor field quantization on M_0 and M_- . . . 13
- 2.4 Bogolubov transformation on M_0 . . . 16
- 2.5 Bogolubov transformation on M_- . . . 28

3 Hawking effect for massive spinors **34**
- 3.1 Introduction . . . 34
- 3.2 Kruskal spacetime and the $\mathbb{RP}^3$ geon . . . 35
- 3.3 Boulware vacuum . . . 38
- 3.4 Bogolubov transformation on the geon . . . 47

4 Stress tensor **53**
- 4.1 Introduction . . . 53
- 4.2 Massive scalar field . . . 54
  - 4.2.1 M_0 . . . 57
  - 4.2.2 M_- . . . 57
  - 4.2.3 Minkowski space with infinite plane boundary . . . 58
- 4.3 Massless spinor field . . . 59
  - 4.3.1 M_0 . . . 63
  - 4.3.2 M_- . . . 64
- 4.4 Massive spinor field . . . 66
  - 4.4.1 M_0 . . . 67
  - 4.4.2 M_- . . . 68

5 Particle detector models **70**
- 5.1 Introduction . . . 70
- 5.2 Background . . . 71
- 5.3 Causal detector with linear coupling . . . 81
  - 5.3.1 Massless scalar field in d-dimensional Minkowski space . . . 81
  - 5.3.2 Massive scalar field . . . 95

5.3.3 The regularization as an ultraviolet cut-off . . . 97
5.3.4 An alternative detector form . . . 98
5.4 Quotients of Minkowski space . . . 100
5.4.1 The automorphic field detector . . . 100
5.4.2 Scalar detector on M_0 . . . 102
5.4.3 Minkowski space with an infinite plane boundary . . . 104
5.4.4 Conical singularity and generalisations . . . 114
5.4.5 Scalar detector on M_- . . . 116
5.5 Causal detector with non-linear coupling . . . 119
5.5.1 Minkowski space . . . 119
5.5.2 Automorphic fields . . . 121
5.6 Causal detector for the Dirac field . . . 124
5.6.1 Minkowski space . . . 124
5.6.2 Dirac detector for automorphic fields . . . 131
5.6.3 Dirac detector on M_0 . . . 133
5.6.4 Dirac detector on M_- . . . 136
5.7 Static detectors on the $\mathbb{RP}^3$ geon . . . 138
5.8 Detectors in conformally flat spacetimes . . . 143
5.9 De Sitter and $\mathbb{RP}^3$ de Sitter spaces . . . 147
5.9.1 Detectors in de Sitter space . . . 148
5.9.2 $\mathbb{RP}^3$ de Sitter space . . . 153

6 Conclusions 158

A Analytic continuation of Rindler modes across the horizons 164

B Evaluating integrals using hyperspherical coordinates 166

C Expressions in 2 dimensions 172
C.1 The correlation function . . . 172
C.2 Inertial detector . . . 175
C.3 Uniformly accelerated detector . . . 177

D Integrals on uniformly accelerated worldline 178

E Details for automorphic field detector 181

F Evaluating (5.95) for inertial detector approaching boundary 183

G Asymptotic behaviour of (5.101) and (5.115) 185

Bibliography 190

List of Figures

2.1 Conformal diagrams of M_0 and M_-. 11

2.2 Rindler space. 17

3.1 Conformal diagram of Kruskal spacetime. 36

3.2 Conformal diagram of the $\mathbb{RP}^3$ geon. 37

5.1 Transition rate of inertial detector on M_0 for an untwisted field and $\omega < 0$. . 104

5.2 Transition rate of inertial detector on M with boundary, motion parallel to boundary, $\omega < 0$.106

5.3 Transition rate of inertial detector on M with boundary, motion perpendicular to boundary, $\omega > 0$.108

5.4 Transition rate of inertial detector on M with boundary, motion perpendicular to boundary, $\omega < 0$.109

5.5 Transition rate of uniformly accelerated detector switched on at $\tau_0 = -\infty$ on M with boundary, motion

5.6 Image term of the transition rate of a uniformly accelerated detector switched on at a finite time on M

5.7 Image term of the transition rate of a uniformly accelerated detector switched on at a finite time on M

5.8 Image term of the transition rate of a uniformly accelerated detector switched on at a finite time on M

5.9 Transition rate associated with the response of a static detector on the $\mathbb{RP}^3$ geon.143

5.10 Transition rate associated with the response of a comoving detector on $\mathbb{RP}^3$ de Sitter space.157

D.1 Contour used in the integration of (D.2). 179

F.1 Contour used in the integration of (F.1). 184

List of Tables

4.1 $\langle T_{\mu\nu} \rangle$ for a left-handed two-component spinor in the Minkowski-like vacua on M_0 and M_- in the ortho

Chapter 1

Introduction

In this thesis we consider quantum field theory on certain families of topologically non-trivial flat and curved spacetimes. We present a collection of new results in three related areas of research. These include the Unruh effect on topologically non-trivial flat spacetimes and Hawking radiation on black hole spacetimes via Bogolubov transformations, the evaluation of expectation values of the stress-energy tensor, and the theory of particle detector models. A good first reference to all these subjects is the well known book by Birrell and Davies [1] [1].

Quantum field theory in curved spacetime has been an active area of research now for many decades, and its results and methods have been of interest to researchers in many related fields. The theory was constructed as a step in the direction of a full, and not yet realised, theory of quantum gravity coupled to matter. The theory considers quantized fields propagating on a fixed curved background manifold, neglecting the dynamical degrees of freedom of the gravitational field. The gravitational field therefore is left unquantized and we are making the assumption that the back reaction of the quantum field on the spacetime is negligible. It is clear that quantum field theory in curved spacetime must be a low energy approximation to a deeper theory whose effects become important as we approach the Planck scale. Despite this limited, and as yet not fully understood, domain in which the theory is considered to be an accurate representation of reality, quantum field theory in curved spacetime has had remarkable successes. Further it is expected that many of its features and predictions must be included and explained within a full theory of quantum gravity.

---

[1] Other noteworthy books on the topic of quantum fields in curved spacetimes include those of Fulling [2] and Wald [3].

One of the major successes of the theory is the description of particle production by time dependent gravitational fields. Particle creation effects may occur due to the non uniqueness in choosing a time coordinate within classical general relativity. In quantum field theory in Minkowski space we define particles using those solutions to the field equations which are positive frequency with respect to the timelike Killing vector ∂_t, in a standard Minkowski coordinate system (t, x, y, z). Due to Lorentz invariance a positive frequency solution in one inertial frame is positive frequency in all inertial frames. Thus the number of particles and in particular the vacuum state (defined as the state free from positive frequency solutions) is the same for all inertial observers. In a general curved spacetime however we no longer have global Lorentz invariance and there may be no Killing vectors at all with respect to which we can define positive frequency modes. Here two observers may not agree on which parts of a solution are positive frequency, even when both observers follow geodesic worldlines, and as a result one observer sees particles in the other's vacuum state. Particles have been created. As we shall see this may be true even in Minkowski space if at least one of the observers is not inertial. The first thorough accounts of particle creation in a gravitational background were given by Parker [4] in the late 60's in relation to cosmology. An expanding universe may create particles in the process of expansion. Since then a large amount of literature has been written on various aspects of these effects.

In 1975 Hawking published his much celebrated result [5] that a black hole formed by a star collapse radiates with a thermal spectrum at temperature

$$T = \frac{\hbar\kappa}{2\pi ck} \ , \tag{1.1}$$

as seen by an observer at infinity, where κ is the hole's surface gravity. This finding influenced a large number of papers in the following years as a systematic study of black hole radiation was undertaken. Further this demonstration of black hole temperature put the speculations of Bekenstein [6] and others, on an analogy between the laws of classical thermodynamics and certain relations in black hole mechanics, on a firmer basis. In particular it led to the acceptance of one quarter of the area [6] of the horizon as the physical entropy of the hole [2]. It was realised shortly after the collapsing star analysis that the same temperature and entropy could be obtained by considering the Kruskal-Szekeres [10, 11] extension of Schwarzschild and on it a quantum state that describes a black hole in thermal equilibrium with its environment [1, 3, 12–16]. The defining characteristics of this Hartle-Hawking state are that it is regular everywhere on the Kruskal manifold and invariant under the continuous isometries [3, 16].

In 1976 Unruh [14], following work by Fulling [17], showed that these thermal effects are not restricted to curved spacetime. He found that a uniformly accelerated observer in Minkowski space perceives the usual Minkowski vacuum as a thermal bath at temperature

$$T = \frac{\hbar a}{2\pi c k} \ , \tag{1.2}$$

where a is the observer's proper acceleration. This effect is widely known as the Unruh effect. Since Unruh's original work the Unruh effect has attracted a lot of interest building a firm foundation for it, including the possibility that the effect could be experimentally observed. Perhaps the most promising proposal so far, though far from being realised in practice, is that of using high intensity lasers to accelerate electrons at very high magnitudes by Chen and Tajima [18].

In this thesis we consider aspects of Hawking radiation and the Unruh effect on certain topologically non-trivial spacetimes. In particular we consider Hawking radiation for Dirac spinors on the spacetime known in the terminology of [19] as the $\mathbb{RP}^3$ geon (for earlier work on the classical properties of this spacetime, see [20, 21]). The $\mathbb{RP}^3$ geon is a $\mathbb{Z}_2$ quotient of Kruskal, it is space and time orientable, it contains a black and white hole, but it only has one static region, isometric to standard exterior Schwarzschild. The geon is an eternal black hole and so as with the Kruskal manifold we do not expect to see one in the sky, as an astrophysical object formed in a star collapse, but its interest lies in the fact that it is an unconventional black hole and in particular has non-trivial topology. It provides an interesting arena in which we can probe our understanding of black hole physics. In particular here we concentrate on the effect of non-trivial topology on the thermal properties. From the physical point of view, a puzzling feature of the Hartle-Hawking state on Kruskal spacetime is its reliance on the whole manifold. The manifold has two static regions, causally disconnected from each other and separated by a bifurcate Killing horizon, but the thermal properties manifest themselves when the state is probed in only one of the static regions. To explore the significance of the second exterior region, we may consider a spacetime such as the $\mathbb{RP}^3$ geon which is locally isometric to Kruskal but which only contains one exterior region. In [22] Louko and Marolf investigated scalar field quantization on the $\mathbb{RP}^3$ geon. It was shown that the Hartle-Hawking like quantum state on the geon does not appear thermal to all observers in the exterior region, but it does appear thermal in the standard Hawking temperature when probed by suitably restricted operators. In particular, the state appears thermal in the standard temperature to every operator far from the hole and with support at asymptotically late (or early) Schwarzschild times.

---

[2] The leading order corrections to this entropy, due to quantum gravity effects have been investigated extensively in the recent literature. The leading corrections are logarithmic in the area (see e.g. [7–9] and references therein.)

In chapter 3 we extend the scalar field analysis of [22] to a massive Dirac field. The main new issue with fermions is that while exterior Schwarzschild and Kruskal both have spatial topology $\mathbb{R} \times S^2$ and hence a unique spin structure, the $\mathbb{RP}^3$ geon has spatial topology $\mathbb{RP}^3 \setminus \{\text{point}\}$ and thus admits two inequivalent spin structures. The geon thus has *two* Hartle-Hawking like states for fermions, one for each spin structure. Our first aim is to examine whether these states appear thermal when probed in the exterior region: We shall find that they do, in a limited sense similar to what was found for the scalar field in [22]. Our second aim is to examine whether these two states can be distinguished by observations limited to the exterior region. We shall find that they can be in principle distinguished by suitable interference experiments: The states contain Boulware-type excitations in correlated pairs, and the spin structure affects the relative phase between the members of each pair.

This means that the restriction of the Hartle-Hawking type state to the geon exterior not only tells that the classical geometry behind the horizons differs from Kruskal but also is sensitive to a quantization ambiguity whose existence cannot be deduced from the exterior geometry. In this sense, probing the quantum state in the exterior region reveals in principle both classical and quantum information from behind the horizons. How this information might be uncovered in practice, for example by particle detectors will be investigated in chapter 5.

As a preparation for this analysis on the $\mathbb{RP}^3$ geon, we begin in chapter 2 by analysing the Unruh effect for the massive Dirac field on a family of Rindler spaces, denoted by M_0 and M_-, whose topology respectively mimics that of Kruskal and the geon [22]. The results are interesting in their own right as they illustrate a particular example of the effect of topology on the thermality of the Unruh effect. They further serve as a flat space model of the Hawking radiation on the geon. While the results are qualitatively similar to those on the geon, the effects of the spin structure appear in a much more transparent form, and these Rindler spaces thus offer a testing ground for localised particle detector models that aim to resolve the phase factors determined by the spin structure. As a by-product, we also obtain the Bogolubov transformation for massive Dirac fermions on (ordinary) Rindler space in $(3+1)$ dimensions, which to the knowledge of the author has not appeared in the literature. Chapters 2 and 3 are based on the paper [23].

One of the most important observables in quantum field theory in curved spacetime is the stress-energy tensor. Classically the stress-energy tensor provides all the information on the energy and momentum content of the spacetime, and provides the source of the gravitational field on the right hand side of Einstein's equations. In the quantum theory the expectation value of the stress-energy tensor in a given state provides information on the energy and momentum content of that state, and it may further be used in a semiclassical analysis as the source of a semiclassical Einstein equation which provides information on the back reaction of the quantum field on the spacetime geometry. In chapter 4 we calculate the expectation values of the stress-energy tensor for massive scalar and spinor fields on the flat spacetimes M_0 and M_-, introduced in chapter 2, in the Minkowski-like vacuum states. We also present the expectation values for massless two-component spinors, showing that the values are independent of the handedness. This chapter is mainly based on the paper [24] and, with the massless spinor results, also on [23].

Finally in chapter 5 we investigate particle detector models. We begin in section 5.2 with a discussion of the history and background of particle detector models in quantum field theory in curved spacetime. We then provide a number of new results. The focus of the chapter is a recent model proposed by Schlicht [25, 26]. The model is that of a monopole detector as first introduced in the context of quantum field theory in curved spacetime by DeWitt [27]. In [25, 26] the author considers monopole detectors following general trajectories in Minkowski space which are switched off at a finite value of the proper time, in contrast to the asymptotic detections usually considered. Although not the first to consider such finite time detections Schlicht seems to be the first to highlight the importance of the regularization of the Wightman function in such calculations. In particular if the Wightman function is regularized using the usual $i\epsilon$ procedure [1] then we are led to an unphysical result when considering a uniformly accelerated detector switched on in the infinite past $\tau_0 = -\infty$ and off at $\tau < \infty$. Schlicht [25, 26] offers an alternative regularization of the correlation function by considering a model of a spatially extended detector. The massless scalar field operator which the detector is coupled to is smeared in the detector's proper reference frame against a suitable window function which represents the shape of the detector. The response depends on the detector's size; however, in the limit as the size goes to 0 the usual result for a uniformly accelerated detector is recovered (that is, a thermal response at temperature (1.2) [14, 27]). The aim of the chapter is twofold. The first aim is to extend Schlicht's model in a number of directions. We first make some comments on Schlicht's regularization and some possible alternatives. We then extend the model to Minkowski space in arbitrary dimension, to the massive scalar field, to fields on quotient spaces of Minkowski space, to the massless non-linearly coupled scalar field and massless Dirac field and to conformally flat spacetimes. The second aim is to investigate some specific detector responses. In particular we consider inertial and uniformly accelerated detectors on the flat, topologically non-trivial spacetimes M_0 and M_- introduced in chapter 2 and on Minkowski space with an infinite plane boundary. The time dependent responses on M_- and on Minkowski space with boundary as the detector approaches the boundary are presented for the first time. Further we address the issue, in the case of the Dirac field, as to whether or not such a detector can distinguish the two spin structures on M_0 and M_-. Finally we consider static detectors in the exterior of the $\mathbb{RP}^3$ geon and comoving observers in $\mathbb{RP}^3$ de Sitter space via their global embedding Minkowski spaces (GEMS) (see e.g [28]). Although until now the GEMS procedure has only been applied to kinematical arguments we expect that at least in some cases the response of detectors in the original curved spaces and the corresponding ones in their GEMS should be related in some way. The case of a comoving detector on $\mathbb{RP}^3$ de Sitter space is a particularly interesting one as we are able to present the calculation both in the curved space and in the embedding space. It is found that the responses are qualitatively very similar. This case should therefore be very useful in assessing the validity of applying the GEMS procedure to cases involving quotient spaces and time dependent detector responses. Chapter 5 is an extended version of [29].

I end this introduction with a note on the terminology and conventions used throughout this thesis. I use the term "Unruh effect" when discussing the thermal (or partial thermal) properties of vacuum states in all flat (possibly topologically non-trivial) spacetimes from the point of view of uniformly accelerated observers. The term "Hawking effect" is used when discussing thermal properties of states in black hole spacetimes. This terminology is in contrast to that used in some of the literature (for example [3]), where "Hawking effect" is reserved for particle production in a collapsing star spacetime and "Unruh effect" is used for the existence of a thermal equilibrium state in a spacetime with a bifurcate Killing horizon.

We work throughout in natural units $\hbar = c = G = 1$ and with metric signature $(+, -, \ldots, -)$. In d-dimensional Minkowski space, the spatial $(d-1)$-vectors are denoted by bold face characters $\mathbf{x} \in \mathbb{R}^{d-1}$ with $\cdot$ the usual scalar product in $\mathbb{R}^{d-1}$, while d-vectors (used occasionally) are given by an italic script x with $x \cdot y = g_{\mu\nu} x^\mu y^\nu$, where $g_{\mu\nu}$ is the

Minkowski metric.

Chapter 2

Unruh effect for massive spinors on flat spacetimes of topology $\mathbb{R}^3 \times S^1$

2.1 Introduction

In this chapter we consider the Unruh effect for a massive Dirac field, using a Bogolubov transformation analysis, on two flat spacetimes with non-trivial topology. Denoted here as M_0 and M_-, these two spacetimes are built as quotients of Minkowski space under the groups generated by the discrete isometries $J_0 : (t, x, y, z) \mapsto (t, x, y, z + 2a)$ and $J_- : (t, x, y, z) \mapsto (t, -x, -y, z + a)$ respectively, where a is a prescribed positive constant. As a by-product we obtain the Bogolubov transformation for massive Dirac fermions on (ordinary) Rindler space in $(3 + 1)$ dimensions, which to the knowledge of the author has not appeared in the literature.[1] The underlying interest in M_0 and M_- arises from their role in modelling, in the context of accelerated observers in flat spacetimes, the Hawking (-Unruh) effect on the

---

[1] Thermality for massive fermions on $(3+1)$-dimensional Rindler space is demonstrated by other methods in [30, 31]. The massive $(1 + 1)$-dimensional case is considered in [32]. The massive $(3 + 1)$ case is addressed in [33, 34] but the Rindler modes constructed therein are not suitably orthonormal in the Dirac inner product.

Kruskal manifold and the $\mathbb{RP}^3$ geon respectively [22]. The Hawking effect on these black hole spacetimes for massive spinors is the topic of chapter 3.

The chapter is constructed as follows. We begin in section 2.2 by describing the spacetimes M_0 and M_- and their properties. In section 2.3 we then quantize the spinor field in these spacetimes in terms of globally defined Minkowski-like modes and construct the Minkowski-like vacuum states $|0_0\rangle$ and $|0_-\rangle$. Uniformly accelerated observers in M_0 are introduced in section 2.4, and the Rindler particle content of the Minkowski like vacuum $|0_0\rangle$ is found from the explicit Bogolubov transformation. A similar analysis for uniformly accelerated observers in M_- is given in section 2.5.

2.2 The spacetimes M_0 and M_-

Let M denote $(3+1)$-dimensional Minkowski space and let (t, x, y, z) be a standard set of Minkowski coordinates. The metric reads

$$ds^2 = dt^2 - dx^2 - dy^2 - dz^2 \quad . \tag{2.1}$$

The flat spacetimes M_0 and M_- are built as quotients of M under the isometry groups generated respectively by the isometries

$$M_0 := M/J_0 \ , \qquad J_0 : (t, x, y, z) \mapsto (t, x, y, z + 2a) \ , \tag{2.2}$$

$$M_- := M/J_- \ , \quad J_- : (t, x, y, z) \mapsto (t, -x, -y, z + a) \ , \tag{2.3}$$

where $a > 0$ is a prescribed constant. J_0 and J_- preserve space and time orientation and act freely and properly discontinuously. M_0 and M_- are thus space and time orientable flat Lorentzian manifolds.

As M is the universal covering space of both M_0 and M_-, we may understand M_0 and

M_- to be coordinatized by (t, x, y, z) with the identifications

$$M_0 : \quad (t, x, y, z) \sim (t, x, y, z + 2a) \ , \tag{2.4}$$

$$M_- : \quad (t, x, y, z) \sim (t, -x, -y, z + a) \ . \tag{2.5}$$

Note further that M_0 is a double cover of M_-. Field theory on M_0 and M_- can be considered via field theory on M by the method of images (see e.g [35, 36]).

Like M, M_0 and M_- are static spacetimes with respect to the global timelike Killing vector ∂_t. They are globally hyperbolic and have spatial topology $\mathbb{R}^2 \times S^1$. M_0 admits seven Killing vectors, which generate translations in all 4 coordinates, boosts in x and y and rotations in the (x, y)-plane. M_- only admits three Killing vectors, generating translations in t and z and rotations in the (x, y)-plane. Conformal diagrams for M_0 and M_- in which the local y, z coordinates have been suppressed are given in figure 2.1.



Figure 2.1: Conformal diagrams of the spacetimes M_0 and M_-. On M_0 each point on the diagram represents a cylinder of circumference $2a$. On M_- the region $x > 0$ is identical to that on M_0, however at $x = 0$ each point represents a suppressed Mobius strip. The infinities of the figures are not faithful to the infinity structures of the spacetimes due to the suppressed dimensions.

Due to the S^1 factor in the spatial topology, M_0 and M_- each admit two inequivalent spin structures (see e.g [37]). That is there are two inequivalent but equally consistent ways of attaching spinors to each. Consider first M_0. We refer to the vierbein

$$\begin{aligned} &V_0 = \partial_t \quad V_2 = \partial_y \\ &V_1 = \partial_x \quad V_3 = \partial_z \ , \end{aligned} \tag{2.6}$$

as the standard vierbein on M_0. In the standard vierbein the two spin structures amount to imposing respectively periodic and antiperiodic boundary conditions as we go around the compact z-direction. Labelling the spin structures by the index $\eta \in \{1, -1\}$, this means

$$\psi(t, x, y, z + 2na) = \eta^n \psi(t, x, y, z) \ , \tag{2.7}$$

where $n \in \mathbb{Z}$ and $\eta = 1$ for periodic spinors (untwisted spinors in the terminology of [1, 37]) and $\eta = -1$ for antiperiodic (twisted) spinors.

Alternatively, another useful vierbein on M_0 is one which rotates counterclockwise by 2π in the (x, y)-plane as z increases by $2a$, that is,

$$\begin{aligned} V_0 &= \partial_t \\ V_1 &= \cos(\pi z/a)\partial_x + \sin(\pi z/a)\partial_y \\ V_2 &= -\sin(\pi z/a)\partial_x + \cos(\pi z/a)\partial_y \\ V_3 &= \partial_z \ . \end{aligned} \tag{2.8}$$

Spinors that are periodic with respect to this rotating vierbein are antiperiodic with respect to the standard vierbein and vice versa. Both these vierbeins are invariant under the action of J_0. One could further introduce a vierbein that rotates clockwise by 2π in the (x, y)-plane as z increases by $2a$ (replace π with $-\pi$ in (2.8)), but periodic (respectively antiperiodic) boundary conditions in this vierbein are equivalent to periodic (antiperiodic) boundary conditions in vierbein (2.8). This shows that neither spin stucture on M_0 involves a preferred spatial orientation.

Now consider M_-. The standard Minkowski vierbein (2.6) is not invariant under J_-

and therefore does not provide a globally defined vierbein on M_-. However both the counterclockwise rotating vierbein (2.8) and its clockwise rotating analogue are invariant under J_- and hence are well defined on M_-. We may therefore specify the two spin structures on M_- by working in the vierbein (2.8) and imposing respectively periodic and antiperiodic boundary conditions under J_-. Labelling the spin structures by the index $\rho \in \{1, -1\}$ this means

$$\psi(t, (-1)^n x, (-1)^n y, z + na) = \rho^n \psi(t, x, y, z) \ , \tag{2.9}$$

where $n \in \mathbb{Z}$ and $\rho = 1$ for the periodic spinors and $\rho = -1$ for the antiperiodic spinors with respect to vierbein (2.8). We could equivalently work in the clockwise rotating vierbein and interchange the periodic and antiperiodic boundary conditions. This shows that the choice of a spin structure on M_- determines a preferred spatial orientation. For concreteness, we shall throughout specify the spin structure with respect to vierbein (2.8).

2.3 Spinor field quantization on M_0 and M_-

In this subsection we consider the quantum theory of a free Dirac field ψ with mass $m \geq 0$ on M_0 and M_-. We introduce the Minkowski-like vacua $|0_0\rangle$ and $|0_-\rangle$ for which positive and negative frequencies are defined with respect to the global timelike Killing vector ∂_t. We work in the rotating tetrad (2.8) throughout.

In a general curved spacetime the spinor Lagrangian in the vierbein formalism is [1]

$$\mathcal{L} = \det V \left(\frac{1}{2} i \left[\bar{\psi} \gamma^\mu \nabla_\mu \psi - (\nabla_\mu \bar{\psi}) \gamma^\mu \psi \right] - m \bar{\psi} \psi \right) \ , \tag{2.10}$$

where V_α^μ is a set of vierbeins, $V_\alpha = V_\alpha^\mu \partial_\mu$, and $\gamma^\mu = V_\alpha^\mu \gamma^\alpha$ are the curved space counterparts of the Dirac matrices which satisfy

$$\{\gamma^\mu, \gamma^\nu\} = 2 g^{\mu\nu} \ . \tag{2.11}$$

$\nabla_\alpha = V_\alpha^\mu(\partial_\mu + \Gamma_\mu)$ is a spinor covariant derivative with

$$\Gamma_\mu = \frac{1}{8} V_\alpha^\nu V_{\beta\nu;\mu} [\gamma^\alpha, \gamma^\beta] \ . \tag{2.12}$$

Variation of the action with respect to $\bar{\psi} \equiv \psi^\dagger \gamma^0$ yields the covariant Dirac equation

$$i\gamma^\mu \nabla_\mu \psi - m\psi = 0 \ . \tag{2.13}$$

It will be useful to work in the local Minkowski coordinates (t, x, y, z) and in the rotating vierbein (2.8) which is well defined on M_0 and M_-. The Dirac equation (2.13) then becomes

$$\begin{aligned} i\{\gamma^0\partial_t + \gamma^1\left(\cos\left(\tfrac{\pi z}{a}\right)\partial_x + \sin\left(\tfrac{\pi z}{a}\right)\partial_y\right) + \gamma^2\left(-\sin\left(\tfrac{\pi z}{a}\right)\partial_x + \cos\left(\tfrac{\pi z}{a}\right)\partial_y\right) \\ +\gamma^3\left(\partial_z - \tfrac{\pi}{4a}\left(\gamma^1\gamma^2 - \gamma^2\gamma^1\right)\right) + im\}\psi = 0 \ . \end{aligned} \tag{2.14}$$

The inner product on M is

$$\langle \psi_1, \psi_2 \rangle = \int dx\, dy\, dz\ \psi_1^\dagger \psi_2 \ , \tag{2.15}$$

on a constant t hypersurface. We denote the inner products on M_0 and M_- by $\langle \psi_1, \psi_2 \rangle_0$ and $\langle \psi_1, \psi_2 \rangle_-$ respectively.

Consider first M_0. To construct solutions to (2.14) that are positive and negative frequency with respect to ∂_t, we begin with the Minkowski space positive and negative frequency solutions in the standard vierbein (2.6) (see for example [38, 39]) and transform to the rotating vierbein (2.8) by the spinor transformation associated with a rotation by π in the (x, y)-plane as z increases by a. Working here and throughout this section in the standard representation of the γ matrices [39], $\gamma^0 = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$ and $\gamma^i = \left(\begin{smallmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{smallmatrix}\right)$, where σ_i are the Pauli matrices, $\sigma_1 = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, $\sigma_2 = \left(\begin{smallmatrix} 0 & -i \\ i & 0 \end{smallmatrix}\right)$ and $\sigma_3 = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$, this transformation reads

$$\psi \mapsto e^{-\frac{\gamma^1\gamma^2\pi z}{2a}}\psi = \operatorname{diag}\left(e^{\frac{i\pi z}{2a}}, e^{\frac{-i\pi z}{2a}}, e^{\frac{i\pi z}{2a}}, e^{\frac{-i\pi z}{2a}}\right)\psi \ . \tag{2.16}$$

Solutions on M_0 can be found from the Minkowski space solutions via the method of images. The periodic and antiperiodic boundary conditions will restrict the momentum in the z-direction. We find that a complete set of normalized positive frequency solutions is $\{U_{j,k_x,k_y,k_z}\}$, where

$$U_{j,k_x,k_y,k_z} = \frac{1}{4\pi}\sqrt{\frac{(\omega+m)}{a\omega}}e^{-i\omega t+ik_x x+ik_y y+ik_z z}\,u_{j,k_x,k_y,k_z} \ , \tag{2.17}$$

with

$$u_1 = \begin{pmatrix} e^{\frac{i\pi z}{2a}} \\ 0 \\ e^{\frac{i\pi z}{2a}}\frac{k_z}{\omega+m} \\ e^{-\frac{i\pi z}{2a}}\frac{k_+}{\omega+m} \end{pmatrix} \ , \quad u_2 = \begin{pmatrix} 0 \\ e^{-\frac{i\pi z}{2a}} \\ e^{\frac{i\pi z}{2a}}\frac{k_-}{\omega+m} \\ e^{-\frac{i\pi z}{2a}}\frac{-k_z}{\omega+m} \end{pmatrix} \ , \tag{2.18}$$

$k_\pm = k_x \pm ik_y$ and $\omega = \sqrt{m^2+k_x^2+k_y^2+k_z^2}$. For spinors that are periodic in the standard vierbein (2.6), $k_z = n\pi/a$ with $n \in \mathbb{Z}$, and in the other spin structure $k_z = (n+\frac{1}{2})\pi/a$ with $n \in \mathbb{Z}$. k_x and k_y take all real values. The orthonormality relation is

$$\langle U_{i,k_x,k_y,k_z}, U_{j,k'_x,k'_y,k'_z}\rangle_0 = \delta_{ij}\delta_{nn'}\delta(k_x-k'_x)\delta(k_y-k'_y) \ . \tag{2.19}$$

Note that the corresponding delta-normalized modes on Minkowski space are $\sqrt{\frac{a}{\pi}}U_{j,k_x,k_y,k_z}$ with $k_z \in \mathbb{R}$.

Consider now M_-. As M_0 is a double cover of M_-, a complete set of modes can be built from (2.17) as linear combinations which are invariant under J_-, that is by superposing the modes (2.17) and their images under J_- with phase factors that lead to the appropriate (anti-)periodicity properties. We choose the set $\{V_{j,k_x,k_y,k_z}\}$ given by

$$\begin{aligned} V_{1,k_x,k_y,k_z} &= U_{1,k_x,k_y,k_z} + \rho i e^{ik_z a}U_{1,-k_x,-k_y,k_z} \ , \\ V_{2,k_x,k_y,k_z} &= U_{2,k_x,k_y,k_z} - \rho i e^{ik_z a}U_{2,-k_x,-k_y,k_z} \ , \end{aligned} \tag{2.20}$$

where $k_z = (n+\frac{1}{2})\pi/a$, $n \in \mathbb{Z}$ and $\rho = 1$ ($\rho = -1$) gives spinors that are periodic (antiperiodic) in the rotating vierbein (2.8). As with the scalar field [22] there is a redundancy in these V-modes in that V_{j,k_x,k_y,k_z} and $V_{j,-k_x,-k_y,k_z}$ are proportional, and we understand this redundancy to be eliminated by taking for example $k_y > 0$. The orthonormality condition reads

$$\langle V_{i,k_x,k_y,k_z}, V_{j,k'_x,k'_y,k'_z}\rangle_- = \delta_{ij}\delta_{nn'}\delta(k_x - k'_x)\delta(k_y - k'_y) \ . \tag{2.21}$$

Note that k_z takes in both spin structures the same set of values, which coincides with the set of modes in the twisted spin structure on M_0.

Given these complete sets of modes we may proceed to canonically quantize the field in the usual way. That is, we expand the field in these sets and impose the usual anticommutation relations on the coefficients, promoted to operators (operator valued distributions). Let $|0\rangle$ be the usual Minkowski vacuum on M, defined by the set $\{U_{j,k_x,k_y,k_z}\}$. We denote by $|0_0\rangle$ the vacuum on M_0 defined by the set $\{U_{j,k_x,k_y,k_z}\}$ with the suitably restricted values of k_z and by $|0_-\rangle$ the vacuum on M_- defined by the set $\{V_{j,k_x,k_y,k_z}\}$. $|0_0\rangle$ and $|0_-\rangle$ both depend on the respective spin structures, but in what follows we will not need an explicit index to indicate this dependence.

2.4 The Unruh construction and Bogolubov transformation on Minkowski space and M_0

In this section we consider a uniformly accelerated observer on the spacetimes M and M_0. We find the Rindler particle content of the Minkowski vacuum $|0\rangle$ on M and the Minkowski-like vacuum $|0_0\rangle$ on M_0.

Let R_0 be the right-hand-side Rindler wedge of M_0, $x > |t|$. We introduce in R_0 the usual Rindler coordinates (τ, ξ, y, z) by

$$\begin{aligned} t &= \xi \sinh\tau \ , \\ x &= \xi \cosh\tau \ , \end{aligned} \tag{2.22}$$

with $\xi > 0$ and $-\infty < \tau < \infty$, understood with the identification $(\tau, \xi, y, z) \sim (\tau, \xi, y, z+2a)$. The metric reads

$$ds^2 = \xi^2 d\tau^2 - d\xi^2 - dy^2 - dz^2 \quad . \tag{2.23}$$

R_0 is a globally hyperbolic spacetime with the complete timelike Killing vector $\partial_\tau = t\partial_x + x\partial_t$, which generates boosts in the (t, x)-plane. The worldines at constant ξ, y and z are those of observers accelerated uniformly in the x-direction with acceleration ξ^{-1} and proper time $\xi\tau$.



Figure 2.2: Rindler space. The surfaces $x = t$ and $x = -t$ form horizons for the Rindler observers. For example an observer in R following the hyperbolic path shown can neither send any information to region L nor receive any information from there. The two regions are causally disconnected.

We need in R_0 a set of orthonormal field modes that are positive frequency with respect to ∂_τ. Working in the vierbein aligned along the Rindler coordinate axes,

$$V_a^\mu = \mathrm{diag}(\xi^{-1}, 1, 1, 1) \ , \tag{2.24}$$

the Dirac equation (2.13) becomes

$$(i\partial_\tau + i\xi\gamma^0\gamma^1\partial_\xi + i\xi\gamma^0\gamma^2\partial_y + i\xi\gamma^0\gamma^3\partial_z + i\gamma^0\gamma^1/2 - m\xi\gamma^0)\psi = 0 \ , \tag{2.25}$$

where the γ matrices are the usual flat space $\gamma's$. A complete set of mode solutions, positive frequency with respect to Rindler Killing time ∂_τ, on R_0 can then be found by directly separating (2.25) by an ansatz of simultaneous eigenfunctions of $-i\partial_y$, $-i\partial_z$ and the Rindler Hamiltonian. In view of comparison with M_- in section 2.5, we wish the solutions to have simple transformation properties under J_-. Modes that achieve this are

$$\psi^R_{j,\Omega,k_y,k_z}(\tau,\xi,y,z) = N_j\left(X^R_j K_{i\Omega-\frac{1}{2}}(\kappa\xi) + Y^R_j K_{i\Omega+\frac{1}{2}}(\kappa\xi)\right)e^{-i\Omega\tau+ik_y y+ik_z z} \ , \tag{2.26}$$

where

$$X^R_1 = \begin{pmatrix} \frac{k_z}{|k_z|}(k_y - im) \\ -i(|k_z|-\kappa) \\ -i(|k_z|-\kappa) \\ \frac{k_z}{|k_z|}(k_y - im) \end{pmatrix} , \qquad Y^R_1 = \begin{pmatrix} \frac{k_z}{|k_z|}(|k_z|-\kappa) \\ i(k_y - im) \\ -i(k_y - im) \\ -\frac{k_z}{|k_z|}(|k_z|-\kappa) \end{pmatrix} ,$$

$$X^R_2 = \begin{pmatrix} \frac{k_z}{|k_z|}(|k_z|-\kappa) \\ i(k_y + im) \\ i(k_y + im) \\ \frac{k_z}{|k_z|}(|k_z|-\kappa) \end{pmatrix} , \qquad Y^R_2 = \begin{pmatrix} \frac{k_z}{|k_z|}(k_y + im) \\ -i(|k_z|-\kappa) \\ i(|k_z|-\kappa) \\ -\frac{k_z}{|k_z|}(k_y + im) \end{pmatrix} , \tag{2.27}$$

and

$$\begin{aligned} N_1 =& \frac{e^{-\frac{i\pi}{4}}\sqrt{\cosh(\pi\Omega)(\kappa^2-k_z^2)}}{4\pi(k_y - im)\sqrt{a\pi(\kappa-|k_z|)}} \ , \\ N_2 =& \frac{e^{-\frac{i\pi}{4}}\sqrt{\cosh(\pi\Omega)(\kappa^2-k_z^2)}}{4\pi(k_y + im)\sqrt{a\pi(\kappa-|k_z|)}} \ , \end{aligned} \tag{2.28}$$

$j = 1, 2$, $\kappa = (m^2 + {k_y}^2 + {k_z}^2)^{1/2}$, $\Omega > 0$ and $k_y \in \mathbb{R}$. In the spin structure where the spinors are periodic (respectively antiperiodic) in the nonrotating vierbein (2.24), the values of k_z are $n\pi/a$ (respectively $(n+\frac{1}{2})\pi/a$) with $n \in \mathbb{Z}$. $K_{iM+\frac{1}{2}}$ is a modified Bessel function [40]. For $k_z = 0$, we understand the formulas in (2.26)–(2.28) and in what follows to stand for their limiting values as $k_z \to 0_+$,

$$\psi^R_{j,\Omega,k_y,0} := \lim_{k_z\to 0^+} \psi^R_{j,\Omega,k_y,k_z} .$$

The modes are orthonormal as

$$\langle \psi^R_{i,\Omega,k_y,k_z}, \psi^R_{j,\Omega',k'_y,k'_z} \rangle_{R_0} = \delta_{ij}\delta_{nn'}\delta(\Omega-\Omega')\delta(k_y-k'_y) \ , \tag{2.29}$$

where the inner product is (see e.g [30])

$$\langle \psi_1, \psi_2 \rangle_{R_0} = \int d\xi \, dy \, dz \ \psi_1^\dagger \psi_2 \ , \tag{2.30}$$

taken on an $\tau =$ constant hypersurface.

While the above modes would be sufficient for quantizing in R_0 in its own right, they are not suitable for analytic continuation arguments across the horizons, as the vierbein (2.24) becomes singular in the limit $x \to |t|$. We therefore express the modes in the rotating vierbein (2.8), which is globally defined on M_0. This choice of vierbein will further make the comparison to M_- transparent in section 2.5. The Lorentz transformation between (2.24) and (2.8) is a boost by rapidity $-\tau$ in the (τ, ξ)-plane followed by a rotation by π as $z \mapsto z + a$ in the (x, y)-plane. The corresponding transformation on the spinors is

$$\psi \mapsto e^{-\frac{\gamma^1\gamma^2\pi z}{2a}} e^{\frac{\gamma^0\gamma^1\tau}{2}} \psi \ , \tag{2.31}$$

which in our conventions reads

$$\begin{pmatrix} e^{\frac{i\pi z}{2a}}\cosh\frac{\tau}{2} & 0 & 0 & e^{\frac{i\pi z}{2a}}\sinh\frac{\tau}{2} \\ 0 & e^{\frac{-i\pi z}{2a}}\cosh\frac{\tau}{2} & e^{\frac{-i\pi z}{2a}}\sinh\frac{\tau}{2} & 0 \\ 0 & e^{\frac{i\pi z}{2a}}\sinh\frac{\tau}{2} & e^{\frac{i\pi z}{2a}}\cosh\frac{\tau}{2} & 0 \\ e^{\frac{-i\pi z}{2a}}\sinh\frac{\tau}{2} & 0 & 0 & e^{\frac{-i\pi z}{2a}}\cosh\frac{\tau}{2} \end{pmatrix}\psi \ . \tag{2.32}$$

In the vierbein (2.8), our solutions thus become

$$\begin{aligned} \psi^R_{j,\Omega,k_y,k_z}(t,x,y,z) = N_j \Big(& X'^R_j K_{i\Omega-\frac{1}{2}}(\kappa\xi) e^{-(i\Omega-\frac{1}{2})\tau} \\ & + Y'^R_j K_{i\Omega+\frac{1}{2}}(\kappa\xi) e^{-(i\Omega+\frac{1}{2})\tau} \Big) e^{ik_y y + ik_z z} \ , \end{aligned} \tag{2.33}$$

where

$$\begin{aligned} X'^R_1 &= \begin{pmatrix} e^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y - im) \\ -ie^{-\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ -ie^{\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ e^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y - im) \end{pmatrix} , \qquad Y'^R_1 = \begin{pmatrix} e^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \\ ie^{-\frac{i\pi z}{2a}}(k_y - im) \\ -ie^{\frac{i\pi z}{2a}}(k_y - im) \\ -e^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \end{pmatrix} , \\ X'^R_2 &= \begin{pmatrix} e^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \\ ie^{-\frac{i\pi z}{2a}}(k_y + im) \\ ie^{\frac{i\pi z}{2a}}(k_y + im) \\ e^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \end{pmatrix} , \qquad Y'^R_2 = \begin{pmatrix} e^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y + im) \\ -ie^{-\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ ie^{\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ -e^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y + im) \end{pmatrix} . \end{aligned} \tag{2.34}$$

In the left hand Rindler wedge $x < -|t|$ of M_0, denoted L_0, we may proceed similarly. On L_0 we define Rindler coordinates (τ, ξ, y, z) by

$$\begin{aligned} t &= -\xi \sinh\tau \ , \\ x &= -\xi \cosh\tau \ , \end{aligned} \tag{2.35}$$

again with $\xi > 0$ and $-\infty < \tau < \infty$ and with the identifications $(\tau, \xi, y, z) \sim (\tau, \xi, y, z + 2a)$

understood. Note that ∂_τ is now past-pointing. In the vierbein (2.8), a complete orthonormal set of positive frequency modes with respect to ∂_τ is $\{\psi^L_{j,\Omega,k_y,k_z}\}$ with

$$\begin{aligned}\psi^L_{j,\Omega,k_y,k_z}(t,x,y,z) = N_j \Big(& X'^L_j K_{i\Omega-\frac{1}{2}}(\kappa\xi) e^{-(i\Omega-\frac{1}{2})\tau} \\ & + Y'^L_j K_{i\Omega+\frac{1}{2}}(\kappa\xi) e^{-(i\Omega+\frac{1}{2})\tau} \Big) e^{ik_y y + ik_z z} \ ,\end{aligned} \tag{2.36}$$

where

$$X'^L_1 = \begin{pmatrix} e^{\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(k_y - im) \\ -ie^{-\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ -ie^{\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ e^{-\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(k_y - im) \end{pmatrix} , \qquad Y'^L_1 = \begin{pmatrix} -e^{\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(|k_z| - \kappa) \\ -ie^{-\frac{i\pi z}{2a}}(k_y - im) \\ ie^{\frac{i\pi z}{2a}}(k_y - im) \\ e^{-\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(|k_z| - \kappa) \end{pmatrix} ,$$

$$X'^L_2 = \begin{pmatrix} -e^{\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(|k_z| - \kappa) \\ -ie^{-\frac{i\pi z}{2a}}(k_y + im) \\ -ie^{\frac{i\pi z}{2a}}(k_y + im) \\ -e^{-\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(|k_z| - \kappa) \end{pmatrix} , \qquad Y'^L_2 = \begin{pmatrix} e^{\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(k_y + im) \\ -ie^{-\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ ie^{\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ -e^{-\frac{i\pi z}{2a}} \frac{k_z}{|k_z|}(k_y + im) \end{pmatrix} , \tag{2.37}$$

and the ranges of Ω, k_y and k_z are as in R_0. When $k_z = 0$, we define

$$\psi^L_{j,\Omega,k_y,0} := \lim_{k_z \to 0^+} \psi^L_{j,\Omega,k_y,k_z} \ .$$

These solutions satisfy the orthonormality relation

$$\langle \psi^L_{i,\Omega,k_y,k_z}, \psi^L_{j,\Omega',k'_y,k'_z} \rangle_{L_0} = \delta_{ij}\delta_{nn'}\delta(\Omega - \Omega')\delta(k_y - k'_y) \ , \tag{2.38}$$

where the inner product reads as in (2.30) with R_0 replaced by L_0.

We may now quantize the field in R_0 and L_0 in the usual manner. A complete set of positive frequency modes with respect to the future-pointing timelike Killing vector is $\{\psi^R_{j,\Omega,k_y,k_z}(t,x,y,z)\}$ in R_0 and $\{\psi^L_{j,-\Omega,k_y,k_z}(t,x,y,z)\}$ in L_0, both with $\Omega > 0$: The minus sign in ψ^L arises because ∂_τ is past-pointing in L_0. The expansion of the field in these modes and their charge conjugates reads

$$\Psi = \sum_j \sum_n \int_0^\infty d\Omega \int_{-\infty}^\infty dk_y \left(a^R_{j,\Omega,k_y,k_z} \psi^R_{j,\Omega,k_y,k_z} + a^L_{j,\Omega,k_y,k_z} \psi^L_{j,-\Omega,k_y,k_z} \right.$$
$$\left. + b^{R\dagger}_{j,\Omega,k_y,k_z} \psi^{R,c}_{j,\Omega,k_y,k_z} + b^{L\dagger}_{j,\Omega,k_y,k_z} \psi^{L,c}_{j,-\Omega,k_y,k_z} \right) \ . \tag{2.39}$$

In the standard representation the charge conjugation reads

$$\psi^c = i\gamma^2 \psi^* = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \psi^* \ , \tag{2.40}$$

where the superscript ^{c} stands for charge conjugation and * for normal complex conjugation. It follows that

$$\begin{aligned} \psi^{R,c}_{1,\Omega,k_y,k_z}(t,x,y,z) &= \ -i\psi^R_{1,-\Omega,-k_y,-k_z}(t,x,y,z) \ , \\ \psi^{R,c}_{2,\Omega,k_y,k_z}(t,x,y,z) &= i\psi^R_{2,-\Omega,-k_y,-k_z}(t,x,y,z) \ , \end{aligned} \tag{2.41}$$

with analogous expressions for ψ^L_1 and ψ^L_2. For modes at $k_z = 0$ these expressions are understood in the limit $\lim_{k_z \to 0^+}$. We now impose the usual anticommutation relations for the annihilation and creation operators

$$\begin{aligned} \left\{ a^R_{i,\Omega,k_y,k_z}, a^{R\dagger}_{j,\Omega',k'_y,k'_z} \right\} &= \delta_{ij}\delta_{nn'}\delta(\Omega - \Omega')\delta(k_y - k'_y) \ , \\ \left\{ b^R_{i,\Omega,k_y,k_z}, b^{R\dagger}_{j,\Omega',k'_y,k'_z} \right\} &= \delta_{ij}\delta_{nn'}\delta(\Omega - \Omega')\delta(k_y - k'_y) \ , \end{aligned} \tag{2.42}$$

with similar relations holding for a^L_{i,Ω,k_y,k_z} and $a^{L\dagger}_{j,\Omega',k'_y,k'_z}$ and all mixed anticommutators vanishing. This construction defines the Rindler-like vacuum states $|0_{R_0}\rangle$, and $|0_{L_0}\rangle$ on M_0

as the states annihilated by all the appropriate annihilation operators.

$$\begin{aligned} a^R_{i,\Omega,k_y,k_z}|0_{R_0}\rangle &= a^L_{i,\Omega,k_y,k_z}|0_{L_0}\rangle = 0 \ , \\ b^R_{i,\Omega,k_y,k_z}|0_{R_0}\rangle &= b^L_{i,\Omega,k_y,k_z}|0_{L_0}\rangle = 0 \ , \qquad \forall i, \Omega, k_y, k_z \ . \end{aligned} \tag{2.43}$$

Now, we wish to find the Rindler mode content of the Minkowski-like vacuum $|0_0\rangle$. While we could proceed directly by calculating the Bogolubov transformation between the modes (2.17) and (2.33) it is easier to follow Unruh's analytic continuation method [14]. That is, we build from the Rindler modes (the ψ modes) a complete set of linear combinations, denoted here as W-modes, which are purely positive frequency with respect to the global timelike Killing vector ∂_t. The vacua of these W-modes is $|0_0\rangle$ and the Bogolubov transformation may be read off easily from the expansion of the field in them.

As the positive frequency Minkowski modes can be defined by the property that they are analytic and bounded in the entire lower half complex t plane, we may construct the W-modes by analytically continuing the set $\{\psi^R_{j,\Omega,k_y,k_z}\}$ across the horizons into the other wedges in the lower half complex t-plane. The continuation into the F wedge transforms our solutions as [2]

$$\psi^R_{j,\Omega,k_y,n}(t,x,y,z) \mapsto \psi^F_{j,\Omega,k_y,n}(t,x,y,z) \ , \tag{2.44}$$

where

$$\begin{aligned} \psi^F_{1,\Omega,k_y,k_z}(t,x,y,z) &= e^{(i\Omega-\frac{1}{2})\frac{i\pi}{2}} N_1 \Big(X'^F_1 K_{i\Omega-\frac{1}{2}}(i\kappa\xi) e^{-(i\Omega-\frac{1}{2})\tau} \\ &\qquad + Y'^F_1 K_{i\Omega+\frac{1}{2}}(i\kappa\xi) e^{-(i\Omega+\frac{1}{2})\tau} \Big) e^{ik_y y + ik_z z} \ , \end{aligned} \tag{2.45}$$

$$\begin{aligned} \psi^F_{2,\Omega,k_y,k_z}(t,x,y,z) &= e^{(i\Omega+\frac{1}{2})\frac{i\pi}{2}} N_2 \Big(X'^F_2 K_{i\Omega-\frac{1}{2}}(i\kappa\xi) e^{-(i\Omega-\frac{1}{2})\tau} \\ &\qquad + Y'^F_2 K_{i\Omega+\frac{1}{2}}(i\kappa\xi) e^{-(i\Omega+\frac{1}{2})\tau} \Big) e^{ik_y y + ik_z z} \ , \end{aligned} \tag{2.46}$$

---

[2] More details of this analytic continuation are given in appenix A.

the modified Bessel function $K_{i\Omega\pm\frac{1}{2}}(i\kappa\xi)$ can be given in terms of Hankel functions as [40],

$$K_{i\omega\pm\frac{1}{2}}(i\kappa\xi) = -\frac{i\pi}{2}e^{-\frac{i\pi}{2}(i\Omega\pm\frac{1}{2})}H^{(2)}_{i\Omega\pm\frac{1}{2}}(\kappa\xi)\ . \tag{2.47}$$

The spinorial factors are given by

$$X_1^{\prime F} = \begin{pmatrix} e^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y - im) \\ -ie^{-\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ -ie^{\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ e^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y - im) \end{pmatrix}, \qquad Y_1^{\prime F} = \begin{pmatrix} ie^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \\ -e^{-\frac{i\pi z}{2a}}(k_y - im) \\ e^{\frac{i\pi z}{2a}}(k_y - im) \\ -ie^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \end{pmatrix},$$

$$X_2^{\prime F} = \begin{pmatrix} -ie^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \\ e^{-\frac{i\pi z}{2a}}(k_y + im) \\ e^{\frac{i\pi z}{2a}}(k_y + im) \\ -ie^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(|k_z| - \kappa) \end{pmatrix}, \qquad Y_2^{\prime F} = \begin{pmatrix} e^{\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y + im) \\ -ie^{-\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ ie^{\frac{i\pi z}{2a}}(|k_z| - \kappa) \\ -e^{-\frac{i\pi z}{2a}}\frac{k_z}{|k_z|}(k_y + im) \end{pmatrix}, \tag{2.48}$$

and the coordinates (τ, ξ) are in F defined by

$$\begin{aligned} t &= \xi\cosh\tau\ , \\ x &= \xi\sinh\tau\ , \end{aligned} \tag{2.49}$$

with $\xi > 0$, $-\infty < \tau < \infty$. These modes are by construction positive frequency with respect to ∂_t for all $\Omega \in \mathbb{R}$. Continuing further into L_0, we find

$$\begin{aligned} \psi^F_{1,\Omega,k_y,k_z} &\mapsto e^{(i\Omega-\frac{1}{2})i\pi}\psi^L_{1,\Omega,k_y,k_z}\ , \\ \psi^F_{2,\Omega,k_y,k_z} &\mapsto e^{(i\Omega+\frac{1}{2})i\pi}\psi^L_{2,\Omega,k_y,k_z}\ , \end{aligned} \tag{2.50}$$

where (τ, ξ) are in L given by (2.35). Making use of the analogues of (2.41) in L_0, we thus see that a complete set of normalised positive frequency W-modes $\{W_{j,\Omega,k_y,k_z}\}$ on M_0 is given by

$$W_{j,\Omega,k_y,k_z}(t,x,y,z) = \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}}\psi^{R}_{j,\Omega,k_y,k_z} + e^{-\frac{\pi\Omega}{2}}\psi^{L,c}_{j,-\Omega,-k_y,-k_z}\right) \ , \tag{2.51}$$

with $\Omega \in \mathbb{R}$.

It will be useful to split these modes into two sets, as

$$\begin{aligned} W^{(1)}_{j,\Omega,k_y,k_z}(t,x,y,z) &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}}\psi^{R}_{j,\Omega,k_y,k_z} + e^{-\frac{\pi\Omega}{2}}\psi^{L,c}_{j,-\Omega,-k_y,-k_z}\right) \ , \\ W^{(2)}_{j,\Omega,k_y,k_z}(t,x,y,z) &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(-e^{-\frac{\pi\Omega}{2}}\psi^{R,c}_{j,\Omega,-k_y,-k_z} + e^{\frac{\pi\Omega}{2}}\psi^{L}_{j,-\Omega,k_y,k_z}\right) \ , \end{aligned} \tag{2.52}$$

where now $\Omega > 0$, and we have made further use of (2.41) and their analogues in L_0. For $k_z = 0$ we interpret (2.52) in the sense $k_z \to 0^+$. The normalization is

$$\langle W_{i,\Omega,k_y,k_z}, W_{j,\Omega',k'_y,k'_z}\rangle_0 = \delta_{ij}\delta_{nn'}\delta(\Omega-\Omega')\delta(k_y-k'_y) \ . \tag{2.53}$$

We may now expand the field in terms of the W-modes (2.52) as

$$\begin{aligned} \Psi = \sum_j \sum_n \int_0^\infty d\Omega \int_{-\infty}^{\infty} dk_y \Big(& c^{(1)}_{j,\Omega,k_y,k_z} W^{(1)}_{j,\Omega,k_y,k_z} + c^{(2)}_{j,\Omega,k_y,k_z} W^{(2)}_{j,\Omega,k_y,k_z} \\ & + d^{(1)\dagger}_{j,\Omega,k_y,k_z} W^{(1),c}_{j,\Omega,k_y,k_z} + d^{(2)\dagger}_{j,\Omega,k_y,k_z} W^{(2),c}_{j,\Omega,k_y,k_z}\Big) \end{aligned} \tag{2.54}$$

and impose the usual anticommutation relations on the annihilation and creation operators in (2.54). The vacuum state for the W-modes is by construction $|0_0\rangle$, with

$$c^{(a)}_{i,\Omega,k_y,k_z}|0_0\rangle = d^{(a)}_{i,\Omega,k_y,k_z}|0_0\rangle = 0 \ , \qquad \forall a, i, \Omega, k_y, k_z \ . \tag{2.55}$$

Equating the expressions (2.54) and (2.39) and taking inner products with the ψ modes, we find that the Bogolubov transformation between the Rindler modes and W-modes reads

$$\begin{aligned} a^R_{j,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}}c^{(1)}_{j,\Omega,k_y,k_z} - e^{-\frac{\pi\Omega}{2}}d^{(2)\dagger}_{j,\Omega,-k_y,-k_z}\right) , \\ a^L_{j,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}}c^{(2)}_{j,\Omega,k_y,k_z} + e^{-\frac{\pi\Omega}{2}}d^{(1)\dagger}_{j,\Omega,-k_y,-k_z}\right) , \\ b^{R,\dagger}_{j,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}}d^{(1)\dagger}_{j,\Omega,k_y,k_z} - e^{-\frac{\pi\Omega}{2}}c^{(2)}_{j,\Omega,-k_y,-k_z}\right) , \\ b^{L,\dagger}_{j,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}}d^{(2)\dagger}_{j,\Omega,k_y,k_z} + e^{-\frac{\pi\Omega}{2}}c^{(1)}_{j,\Omega,-k_y,-k_z}\right) . \end{aligned} \tag{2.56}$$

To find the expectation value of the number operator for Rindler modes in the vacuum $|0_0\rangle$, we use (2.56) to compute

$$\langle 0_0|a^{R\dagger}_{i,\Omega,k_y,k_z}a^R_{j,\Omega',k'_y,k'_z}|0_0\rangle = \frac{1}{(e^{2\pi\Omega}+1)}\delta_{ij}\delta_{nn'}\delta(k_y-k'_y)\delta(\Omega-\Omega') . \tag{2.57}$$

From (2.57) we see immediately that the Minkowski-like vacuum on M_0 contains Rindler particles. Further, the expected number of Rindler particles in the Minkowski-like vacuum is given by the Fermi distribution function. Recall that an observer with constant ξ is uniformly accelerating with proper time $\xi\tau$ and acceleration ξ^{-1}. The proper energy of a Rindler particle seen by such an observer is thus not Ω but Ω/ξ. The usual interpretation of (2.57) is then that a Rindler observer at constant ξ sees the Minkowski vacuum as a thermal bath at temperature $T = \frac{a}{2\pi}$ where $a = 1/\xi$ is the observer's acceleration. We must note of course that setting primed and unprimed indices in (2.57) equal gives a divergent result, due to the continuum normalization of our modes. A finite result may be obtained by considering wave packets, as done for the scalar field in [22].

We can gain more insight into the situation by writing the Minkowski-like vacuum $|0_0\rangle$ in terms of the Rindler vacuum $|0_{R_0}\rangle$. Making use of the inverse Bogolubov transformation to (2.56), we find

$$|0_0\rangle = \prod_{\sigma,j,\Omega,k_y,k_z}\frac{1}{(e^{-2\pi\Omega}+1)^{\frac{1}{2}}}\sum_{q=0,1}(-1)^q e^{-\pi\Omega q}|q\rangle^R_{\sigma,j,\Omega,k_y,k_z}|q\rangle^L_{-\sigma,j,\Omega,-k_y,-k_z} , \tag{2.58}$$

where we have introduced $\sigma = +1, (-1)$ to label particle (antiparticle) modes respectively and the notation on the right hand side is adapted to the tensor product structure of the Hilbert space considered:

$$\begin{aligned} |q\rangle^R_{+,j,\Omega,k_y,k_z} &= (a^{R\dagger}_{j,\Omega,k_y,k_z})^q |0_{R_0}\rangle \quad |q\rangle^L_{+,j,\Omega,k_y,k_z} = (a^{L\dagger}_{j,\Omega,k_y,k_z})^q |0_{L_0}\rangle \ , \\ |q\rangle^R_{-,j,\Omega,k_y,k_z} &= (b^{R\dagger}_{j,\Omega,k_y,k_z})^q |0_{R_0}\rangle \quad |q\rangle^L_{-,j,\Omega,k_y,k_z} = (b^{L\dagger}_{j,\Omega,k_y,k_z})^q |0_{L_0}\rangle \ . \end{aligned} \tag{2.59}$$

The result (2.58) is the massive fermion version of the familiar bosonic result [1], indicating an entangled state in which the right and left Rindler excitations appear in correlated pairs. Suppose we consider an operator $\hat{A}^{(1)}$ whose support is in the right hand Rindler wedge only. This operator will not couple to any of the degrees of freedom of the field in the left hand wedge, and we must trace out these degrees of freedom. The operator has expectation value $\langle 0_0|\hat{A}^{(1)}|0_0\rangle = \mathrm{Tr}(\hat{A}^{(1)}\rho^{(1)})$, where $\rho^{(1)}$ is a fermionic thermal density matrix in R_0,

$$\rho^{(1)} = \prod_{\sigma,j,\Omega,k_y,k_z} \sum_{q=0,1} \frac{e^{-2q\pi\Omega}}{\sum_{m=0,1} e^{-2m\pi\Omega}} |q\rangle^R_{\sigma,j,\Omega,k_y,k_z} \, {}^{R}_{\sigma,j,\Omega,k_y,k_z}\langle q| \ . \tag{2.60}$$

We can conclude that such an operator sees $|0_0\rangle$ as a thermal bath at the usual Unruh temperature. Similar conclusions can be made for any operator whose support is entirely in the left hand wedge.

It is important to note the formal nature of (2.58): The normalisation factor is actually divergent, owing to the infinite product over modes and indicating the non-existence of a unitary operator linking the two vacua. This is not a problem however as the thermality arguments can in fact be made mode for mode (see e.g [31]). Further we note that the usual discussion may follow about the relation of (2.58) to Thermofield dynamics (as first discussed by Israel in [13]).

The result (2.58) incorporates the two spin structures on M_0, R_0 and L_0 in the allowed values of k_z, and these values are the same in all our mode sets. The twisted (respectively untwisted) $|0_0\rangle$ thus induces twisted (untwisted) thermal states in both R_0 and L_0.

To end this section, we note that the Bogolubov transformation on Minkowski space can be read off from our formulas on M_0 with minor systematic changes. There is now only one spin structure and k_z takes all real values. The expressions for the various mode functions include the additional factor $\sqrt{a/\pi}$, sums over n become integrals over k_z, and in the normalization and anticommutation relations the discrete delta $\delta_{nn'}$ is replaced by the delta-function $\delta(k_z - k'_z)$. The formulas involve still a because the spinors are expressed in the rotating vierbein (2.8). Translation into the standard vierbein (2.6) can be accomplished by the appropriate spinor transformation.

2.5 The Unruh construction and Bogolubov transformation on M_-

In this section we repeat the analysis of the previous section for M_-.

Let R_- denote the Rindler wedge on M_-, given in our local coordinates by $|x| > |t|$. As R_- is isometric to R_0, we may introduce in R_- the Rindler-coordinates (τ, ξ, y, z) by (2.22), again with the identifications $(\tau, \xi, y, z) \sim (\tau, \xi, y, z + 2a)$, and quantize as in R_0, defining the positive and negative frequencies with respect to the Killing vector ∂_τ. The set $\{\psi^R_{j,\Omega,k_y,k_z}\}$ (2.33) is therefore a complete set of positive frequency Rindler modes on R_-. We could work directly with this set, but for convenience of phase factors in the W-modes (2.72) below, we use instead the mode set $\{\Psi^R_{j,\Omega,k_y,k_z}\}$, defined as in (2.33) except with the normalization factors (2.28) replaced by

$$\begin{aligned} N_1 &= \frac{e^{-\frac{ik_z a}{2}}\sqrt{\cosh(\pi\Omega)(\kappa^2 - k_z^2)}}{4\pi(k_y - im)\sqrt{a\pi(\kappa - |k_z|)}} \ , \\ N_2 &= \frac{e^{\frac{ik_z a}{2}}\sqrt{\cosh(\pi\Omega)(\kappa^2 - k_z^2)}}{4\pi(k_y + im)\sqrt{a\pi(\kappa - |k_z|)}} \ . \end{aligned} \tag{2.61}$$

A key difference from R_0 arises now, from the requirement that the spinors on R_- must be extendible into spinors in one of the two spin structures on M_-. By the discussion in section 2.4, this implies that k_z is restricted to the values $k_z = (n + \frac{1}{2})\pi/a$ with $n \in \mathbb{Z}$. Both of the spin structures on M_- thus induce on R_- the same spin structure, in which the spinors are antiperiodic in the nonrotating vierbein (2.6) [3]. $\{\Psi^R_{j,\Omega,k_y,k_z}\}$ is a complete set of positive frequency solutions in R_-, written in the globally defined rotating vierbein (2.8). The inner product on R_- is as in (2.30) and the orthonormality relation is

$$\langle \Psi^R_{i,\Omega,k_y,k_z}, \Psi^R_{j,\Omega',k'_y,k'_z} \rangle_{R_-} = \delta_{ij}\delta_{nn'}\delta(\Omega-\Omega')\delta(k_y-k'_y) \ . \tag{2.62}$$

We quantize the field in R_- by expanding Ψ as

$$\begin{aligned}\Psi = \sum_n \int_0^\infty d\Omega \int_{-\infty}^\infty dk_y \Big(& a_{1,\Omega,k_y,k_z} \Psi^R_{1,\Omega,k_y,k_z} + a_{2,\Omega,k_y,k_z} \Psi^R_{2,\Omega,k_y,k_z} \\ & + b^\dagger_{1,\Omega,k_y,k_z} \Psi^{R,c}_{1,\Omega,k_y,k_z} + b^\dagger_{2,\Omega,k_y,k_z} \Psi^{R,c}_{2,\Omega,k_y,k_z} \Big)\end{aligned} \tag{2.63}$$

and imposing the usual anticommutation relations for the annihilation and creation operators. The Rindler-vacuum $|0_{R_-}\rangle$ on R_- is the state annihilated by all the annihilation operators in (2.63),

$$a_{i,\Omega,k_y,k_z}|0_{R_-}\rangle = b_{i,\Omega,k_y,k_z}|0_{R_-}\rangle = 0 \ , \qquad \forall i, \Omega, k_y, k_z \ . \tag{2.64}$$

To find the Rindler-mode content of $|0_-\rangle$, we again use the analytic continuation method. Working in the local coordinates (t,x,y,z), we continue the modes $\{\Psi^R_{j,\Omega,k_y,k_z}\}$ across the horizons in the lower half-plane in complexified t and form linear combinations that are globally well-defined on M_-. We begin by continuing the $\{\Psi^R_{j,\Omega,k_y,k_z}\}$ into the F region in the lower half complex t-plane as on M_0. We obtain the modes $\{\Psi^F_{j,\Omega,k_y,k_z}\}$ identical to the $\{\psi^F_{j,\Omega,k_y,k_z}\}$ of (2.45) and (2.46) but with N_j as in (2.61). These modes are a complete set of W-modes in the F region of M_0 (with the suitable restrictions on k_z). The analogous modes on M_- may be built from these via the method of images. As we are working in the globally defined rotating vierbein which is invariant under J_-, the W-modes on F_- are simply given by (see e.g. [36])

---

[3] This is easily seen by noting that a vierbein which rotates by π (repectively $-\pi$) as $z \to z+a$ thus rotates by 2π (-2π) as $z \to z+2a$. As we noted earlier these two vierbeins on M_0 are equivalent when used to specify the spin structure.

$$W^F_{j,\Omega,k_y,k_z}(t,x,y,z) \propto \Psi^F_{j,\Omega,k_y,k_z}(t,x,y,z) + \rho\Psi^F_{j,\Omega,k_y,k_z}(t,-x,-y,z+a) \ , \tag{2.65}$$

where $\Omega \in \mathbb{R}$ and we have used the parameter $\rho = +1, (-1)$ to label W-modes with periodic (antiperiodic) boundary conditions respectively. Now it is easy to show

$$\begin{aligned}
\Psi^F_{1,\Omega,k_y,k_z}(t,-x,-y,z+a) &= e^{-\pi\Omega}\Psi^F_{2,-\Omega,-k_y,k_z}(t,x,y,z) \ , &(2.66)\\
\Psi^F_{2,\Omega,k_y,k_z}(t,-x,-y,z+a) &= e^{-\pi\Omega}\Psi^F_{1,-\Omega,-k_y,k_z}(t,x,y,z) \ , &(2.67)
\end{aligned}$$

so that

$$\begin{aligned}
W^F_{1,\Omega,k_y,k_z}(t,x,y,z) &\propto \Psi^F_{1,\Omega,k_y,k_z}(t,x,y,z) + \rho e^{-\pi\Omega}\Psi^F_{2,-\Omega,-k_y,k_z}(t,x,y,z) \ , &(2.68)\\
W^F_{2,\Omega,k_y,k_z}(t,x,y,z) &\propto \Psi^F_{2,\Omega,k_y,k_z}(t,x,y,z) + \rho e^{-\pi\Omega}\Psi^F_{1,-\Omega,-k_y,k_z}(t,x,y,z) \ , &(2.69)
\end{aligned}$$

with $\Omega \in \mathbb{R}$. There is a redundancy in the W-modes (2.68) and (2.69) in that

$$W^F_{1,\Omega,k_y,k_z} \propto W^F_{2,-\Omega,-k_y,k_z} \ , \tag{2.70}$$

and we eliminate this redundancy by taking $\Omega > 0$ and $-\infty < k_y < \infty$. Continuing back to R_- in the lower half complex t plane, $\Psi^F_{j,\Omega,k_y,k_z} \mapsto \Psi^R_{j,\Omega,k_y,k_z}$, and making use of the relations

$$\begin{aligned}
\Psi^{R,c}_{1,\Omega,k_y,k_z}(t,x,y,z) &= -\Psi^R_{1,-\Omega,-k_y,-k_z}(t,x,y,z) \ ,\\
\Psi^{R,c}_{2,\Omega,k_y,k_z}(t,x,y,z) &= \Psi^R_{2,-\Omega,-k_y,-k_z}(t,x,y,z) \ , &(2.71)
\end{aligned}$$

we find a complete set $\{W^R_{j,\Omega,k_y,k_z}\}$ of W-modes on R_- to be

$$\begin{aligned} W^R_{1,\Omega,k_y,k_z}(t,x,y,z) &= \frac{1}{\sqrt{2\cosh\pi\Omega}}\left(e^{\frac{\pi\Omega}{2}}\Psi^R_{1,\Omega,k_y,k_z} + \rho e^{-\frac{\pi\Omega}{2}}\Psi^{R,c}_{2,\Omega,k_y,-k_z}\right) \ , \\ W^R_{2,\Omega,k_y,k_z}(t,x,y,z) &= \frac{1}{\sqrt{2\cosh\pi\Omega}}\left(e^{\frac{\pi\Omega}{2}}\Psi^R_{2,\Omega,k_y,k_z} - \rho e^{-\frac{\pi\Omega}{2}}\Psi^{R,c}_{1,\Omega,k_y,-k_z}\right) \ , \end{aligned} \tag{2.72}$$

with $\Omega > 0$. The orthonormality condition is

$$\langle W^R_{i,\Omega,k_y,k_z}, W^R_{j,\Omega',k'_y,k'_z}\rangle_{R_-} = \delta_{ij}\delta_{nn'}\delta(\Omega-\Omega')\delta(k_y-k'_y) \ . \tag{2.73}$$

The expansion of the field in W-modes reads

$$\begin{aligned} \Psi = \sum_n \int_0^\infty d\Omega \int_{-\infty}^\infty dk_y \Big(& c_{1,\Omega,k_y,k_z} W^R_{1,\Omega,k_y,k_z} + c_{2,\Omega,k_y,k_z} W^R_{2,\Omega,k_y,k_z} \\ & + d^\dagger_{1,\Omega,k_y,k_z} W^{R,c}_{1,\Omega,k_y,k_z} + d^\dagger_{2,\Omega,k_y,k_z} W^{R,c}_{2,\Omega,k_y,k_z} \Big) \ , \end{aligned} \tag{2.74}$$

where $k_z = \frac{(2n+1)\pi}{2a}$. The annihilation and creation operators in (2.74) satisfy the usual anticommutation relations, and $|0_-\rangle$ is the state annihilated by the annihilation operators. It follows that the Bogolubov transformation between the Rindler-modes and the W-modes reads

$$\begin{aligned} a_{1,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}} c_{1,\Omega,k_y,k_z} - \rho e^{-\frac{\pi\Omega}{2}} d^\dagger_{2,\Omega,k_y,-k_z}\right) \ , \\ a_{2,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}} c_{2,\Omega,k_y,k_z} + \rho e^{-\frac{\pi\Omega}{2}} d^\dagger_{1,\Omega,k_y,-k_z}\right) \ , \\ b^\dagger_{1,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}} d^\dagger_{1,\Omega,k_y,k_z} - \rho e^{-\frac{\pi\Omega}{2}} c_{2,\Omega,k_y,-k_z}\right) \ , \\ b^\dagger_{2,\Omega,k_y,k_z} &= \frac{1}{\sqrt{2\cosh(\pi\Omega)}}\left(e^{\frac{\pi\Omega}{2}} d^\dagger_{2,\Omega,k_y,k_z} + \rho e^{-\frac{\pi\Omega}{2}} c_{1,\Omega,k_y,-k_z}\right) \ . \end{aligned} \tag{2.75}$$

Now consider the expectation values of the Rindler mode number operators in the vacuum

$|0_-\rangle$. Using (2.75), we find

$$\langle 0_-|a^\dagger_{i,\Omega,k_y,k_z} a_{j,\Omega',k_y',k_z'}|0_-\rangle = \frac{1}{(e^{2\pi\Omega}+1)}\delta_{ij}\delta_{nn'}\delta(k_y-k_y')\delta(\Omega-\Omega') \ , \tag{2.76}$$

which is identical to the M_0 result (2.57) and hence compatible with a bath of Rindler particles at the usual Unruh temperature. Note further that the expectation values are the same in both spin structures. More information may however be gained by expressing $|0_-\rangle$ in terms of $|0_{R_-}\rangle$. The expression analogous to (2.58) is

$$|0_-\rangle = \prod_{\sigma,\Omega,k_y,k_z} \frac{1}{(e^{-2\pi\Omega}+1)^{\frac{1}{2}}} \sum_{q=0,1} (-\rho)^q e^{-\pi\Omega q} |q\rangle_{\sigma,1,\Omega,k_y,k_z} |q\rangle_{-\sigma,2,\Omega,k_y,-k_z} \ , \tag{2.77}$$

where $\sigma = +1(-1)$ and

$$\begin{aligned} |q\rangle_{+,j,\Omega,k_y,k_z} &= (a^\dagger_{j,\Omega,k_y,k_z})^q |0_{R_-}\rangle \ , \\ |q\rangle_{-,j,\Omega,k_y,k_z} &= (b^\dagger_{j,\Omega,k_y,k_z})^q |0_{R_-}\rangle \ . \end{aligned} \tag{2.78}$$

From (2.77) it is seen that $|0_-\rangle$ is in fact an entangled state of Rindler-excitations, and the correlations are between a particle and an antiparticle with opposite eigenvalues of k_z both located in R_-. This is in some sense what we would expect as the involution giving M_- from M_0 identifies the two Rindler wedges. As all the excitations are in the unique Rindler wedge R_-, the expectation values of generic operators in R_- are not thermal. However, for any operator that couples to one member of each correlated pair in R_-, the expectation values will be indistinguishable from those in the corresponding state $|0_0\rangle$ in R_0, indicating thermality in the standard Unruh temperature. Consider for example an operator $\hat{A}$, concentrated on an accelerated worldine in R_-, such that $\hat{A}$ only couples to one of the Rindler modes, say ψ_{1,Ω,k_y,k_z}. The expectation value of $\hat{A}$ is then

$$\begin{aligned} \langle 0_-|\hat{A}|0_-\rangle &= \prod_{\sigma,\Omega,k_y,k_z} \frac{1}{(1+e^{-2\pi\Omega})} \sum_{q=0,1} e^{-2q\pi\Omega}{}_{\sigma,1,\Omega,k_y,k_z}\langle q|\hat{A}|q\rangle_{\sigma,1,\Omega,k_y,k_z} \ , \\ &= \mathrm{Tr}\,(\hat{A}\rho^{(1)}) \ , \end{aligned} \tag{2.79}$$

where $\rho^{(1)}$ has the form of a thermal density matrix. We can conclude that $\hat{A}$ sees $|0_-\rangle$ as a thermal bath at the usual temperature. This is also the case for example for any operator that only couples to excitations with a definite sign of k_z. In particular, we have seen that the number operator expectation values are indistinguishable from (2.57). Further it can be argued from the isometries as in [22] that the experiences of a Rindler-observer become asymptotically thermal at early or late times.

A key result of our analysis is that while both spin structures on M_- induce a state in the same spin structure in R_-, the explicit appearance of ρ in (2.77) shows that the two states differ. A Rindler-observer in R_- can therefore in principle detect the spin structure on M_- from the non-thermal correlations. How these correlations could be detected in practice, for example by particle detectors with a local coupling to the field, is an interesting question that we will touch upon in chapter 5. As the restriction of $|0_-\rangle$ to R_- is not invariant under the Killing vector ∂_τ, and as the isometry arguments show that the correlations disappear in the limit of large $|\tau|$, investigating this question requires a particle detector formalism that can accommodate time dependent situations [25].

Chapter 3

Hawking (-Unruh) effect for massive spinors on the Kruskal manifold and the $\mathbb{RP}^3$ geon

3.1 Introduction

In this chapter we analyze the Hawking effect on the $\mathbb{RP}^3$ geon black hole spacetime for a massive Dirac field. Section 3.2 begins by reviewing some relevant properties of the Kruskal manifold and the $\mathbb{RP}^3$ geon. Here we also discuss the spin structures that these spacetimes admit. In section 3.3 we consider spinor field theory on the geon and we construct the Boulware vacuum $|0_B\rangle$ in one exterior region. This construction is analogous to that of the Rindler vacuum $|0_R\rangle$ in section 2.3. In section 3.4 we construct the Hartle-Hawking-like vacuum $|0_G\rangle$ on the geon and compute the Bogolubov transformation between $|0_B\rangle$ and $|0_G\rangle$. We also give an expression for $|0_G\rangle$ in terms of $|0_B\rangle$, analogous to equation (2.77).

3.2 Kruskal spacetime and the $\mathbb{RP}^3$ geon

In the notation of [22], the Kruskal metric in the Kruskal coordinates (T, X, θ, ϕ) reads

$$ds^2 = \frac{32M^3}{r} e^{-r/(2M)}(dT^2 - dX^2) - r^2(d\theta^2 + \sin^2\theta d\phi^2) \ , \tag{3.1}$$

where $M > 0$, $T^2 - X^2 < 1$ and r is determined as a function of T and X by $T^2 - X^2 = 1 - r/(2M)$. The manifold consists of the right and left exteriors, denoted respectively by R and L, and the black hole (future) and white hole (past) interiors, denoted respectively by F and P, separated from each other by the bifurcate Killing horizon at $|T| = |X|$ (see the conformal diagram, figure 3.1). The four regions may be individually covered by the Schwarzschild coordinates (t, r, θ, ϕ), in which the metric reads

$$ds^2 = \left(1 - \frac{2M}{r}\right) dt^2 - \frac{dr^2}{\left(1 - \frac{2M}{r}\right)} - r^2(d\theta^2 + \sin^2\theta d\phi^2) \ , \tag{3.2}$$

where $2M < r < \infty$ and t (r) is timelike (spacelike) in the exteriors while $0 < r < 2M$ and t (r) is spacelike (timelike) in the interiors. The transformation between Kruskal and Schwarzschild coordinates reads

$$\begin{aligned} T =& \pm \left(\frac{r}{2M} - 1\right)^{\frac{1}{2}} e^{r/(4M)} \sinh\left(\frac{t}{4M}\right) \ , \\ X =& \pm \left(\frac{r}{2M} - 1\right)^{\frac{1}{2}} e^{r/(4M)} \cosh\left(\frac{t}{4M}\right) \ , \end{aligned} \tag{3.3}$$

in R $(+)$ and L $(-)$, and

$$\begin{aligned} T =& \pm \left(-\frac{r}{2M} + 1\right)^{\frac{1}{2}} e^{r/(4M)} \cosh\left(\frac{t}{4M}\right) \ , \\ X =& \pm \left(-\frac{r}{2M} + 1\right)^{\frac{1}{2}} e^{r/(4M)} \sinh\left(\frac{t}{4M}\right) \ , \end{aligned} \tag{3.4}$$

in F $(+)$ and P $(-)$. The exteriors are static, with the timelike Killing vector ∂_t. The Kruskal manifold is a globally hyperbolic, inextendible black hole spacetime. The analogy between the accelerated observers in Minkowski space with constant ξ, y, z and the static observers in the Schwarzschild region R with constant r, θ, ϕ is clear. The vector ∂_t in Kruskal is analogous to ∂_τ which generated boosts in the flat spaces, while ∂_T is analogous to ∂_t in Minkowski space (however ∂_T is not a Killing vector on the Kruskal manifold). The Kruskal manifold has two static regions, each isometric to the Schwarzschild spacetime, causally disconnected from each other and separated by a bifurcate Killing horizon. No event in R can causally influence any event in L, and vice versa, as the two regions are connected only by spacelike curves.



Figure 3.1: Conformal diagram for the Kruskal Spacetime. Each point on the diagram represents a suppressed 2-sphere.

The $\mathbb{RP}^3$ geon is the quotient of the Kruskal manifold under the $\mathbb{Z}_2$ isometry group generated by the map

$$J : (T, X, \theta, \phi) \mapsto (T, -X, \pi - \theta, \phi + \pi) \ . \tag{3.5}$$

The construction is analogous to that of M_- from M_0 in section 2.2. Taking the quotient of the Kruskal manifold under the map (3.5) identifies the two exterior regions with a reflection in X accompanied by an antipodal map on the 2-sphere. (3.5) preserves space and time orientation and acts freely and properly discontinuously. The $\mathbb{RP}^3$ geon is thus a globally hyperbolic inextendible black hole spacetime which is locally isometric to Kruskal but which contains only the one exterior region R. The regions $X > 0$ of the Kruskal manifold and the $\mathbb{RP}^3$ geon are isometric. Figure 3.2 is a conformal diagram of the geon.



Figure 3.2: Conformal diagram of the $\mathbb{RP}^3$ geon spacetime. The regions $X > 0$ of Kruskal and the $\mathbb{RP}^3$ geon are isometric. Points $X > 0$ represent a suppressed 2-sphere while points at $X = 0$ represent a copy of $\mathbb{RP}^2$.

As the Kruskal manifold has spatial topology $\mathbb{R} \times S^2$, it is simply connected and has a unique spin structure. The quotient construction implies [41] that the geon has fundamental group $\mathbb{Z}_2$ and admits two spin structures. As in section 2.2, we describe these spin structures in terms of periodic and antiperiodic boundary conditions for spinors in a specified vierbein. On Kruskal, a standard reference vierbein is

$$\begin{aligned} V_0 &= \sqrt{\frac{r}{32M^3}} e^{\frac{r}{4M}} \partial_T \qquad & V_2 &= \tfrac{1}{r}\partial_\theta \\ V_1 &= \sqrt{\frac{r}{32M^3}} e^{\frac{r}{4M}} \partial_X & V_3 &= \tfrac{1}{r\sin\theta}\partial_\phi \ , \end{aligned} \tag{3.6}$$

which is aligned along Kruskal coordinate axes. This vierbein is not invariant under J. A second useful vierbein which is invariant under J is

$$\begin{aligned}
V_0 &= \sqrt{\frac{r}{32M^3}}e^{\frac{r}{4M}}\partial_T \\
V_1 &= \cos\phi\sqrt{\frac{r}{32M^3}}e^{\frac{r}{4M}}\partial_X + \frac{\sin\phi}{r}\partial_\theta \\
V_2 &= -\sin\phi\sqrt{\frac{r}{32M^3}}e^{\frac{r}{4M}}\partial_X + \frac{\cos\phi}{r}\partial_\theta \\
V_3 &= \frac{1}{r\sin\theta}\partial_\phi \ , \qquad (3.7)
\end{aligned}$$

which rotates by π in the (X, θ) tangent plane as ϕ increases by π. The vierbein (3.7) is well defined on the geon, and when the spinors are written with respect to it, the two spin structures correspond to respectively periodic and antiperiodic boundary conditions as $\phi \mapsto \phi + \pi$. Alternatively we could consider an invariant vierbein which rotates by $-\pi$ as $\phi \mapsto \phi + \pi$ again imposing suitable boundary conditions. Periodic spinors with respect to this vierbein would be antiperiodic with respect to (3.7) and vice versa. Both (3.6) and (3.7) are singular at $\theta = 0$ and $\theta = \pi$, but these coordinate singularities on the sphere can be handled by usual methods and will not affect our discussion.

In practice, we will work with the standard Kruskal vierbein (3.6). Although this vierbein is not invariant under J it is the simplest one to work with on the Kruskal manifold. The boundary conditions appropriate for the two geon spin structures will be found by the method-of-images technique of the Appendix of [36].

3.3 Spinor field theory on Schwarzschild spacetime and the Boulware vacuum

In this section we review the construction of the Boulware vacuum $|0_B\rangle$ for a massive Dirac field in one exterior region of either Kruskal or the $\mathbb{RP}^3$ geon [42]. The Boulware vacuum is that empty of particles when positive frequency is defined with respect to Schwarzschild Killing time ∂_t. While this vacuum as such is well known, we will need to decompose the field in a novel basis in order to make contact with the geon in subsection 3.4.

We consider the quantization of the spinor field ψ in one exterior region R in Schwarzschild coordinates. Here there are two natural choices of vierbein to work with. The first is aligned along Schwarzschild coordinate axes, the second aligned along a set of axes oriented relative to fixed orthogonal directions [43]. We choose to work with a vierbein aligned along the Schwarzschild coordinate axes, given by

$$V_a^\mu = \mathrm{diag}\left(\frac{1}{\left(1-\frac{2M}{r}\right)^{\frac{1}{2}}}, \left(1-\frac{2M}{r}\right)^{\frac{1}{2}}, \frac{1}{r}, \frac{1}{r\sin\theta} \right) . \tag{3.8}$$

This choice has the disadvantage of a more complex angular component analysis but the advantage of being analogous to the vierbein (2.24) used in our Rindler calculations.

The Dirac equation (2.13) then reads

$$\begin{aligned}\Bigg[m + \frac{\gamma^2}{ir\sin^{\frac{1}{2}}\theta}\partial_\theta \sin^{\frac{1}{2}}\theta + \frac{\gamma^3}{ir\sin\theta}\partial_\phi + \frac{\gamma^0}{i\left(1-\frac{2M}{r}\right)^{\frac{1}{2}}}\partial_t \\ + \frac{\gamma^1}{ir}\left(1-\frac{2M}{r}\right)^{\frac{1}{4}}\partial_r\left(1-\frac{2M}{r}\right)^{\frac{1}{4}} r \Bigg]\psi = 0 ,\end{aligned} \tag{3.9}$$

where the γ matrices are flat space γ's.

Let us at this point choose a particular representation of the γ matrices. We take

$$\gamma^0 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} , \quad \gamma^1 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} ,$$

$$\gamma^2 = \begin{pmatrix} -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix} , \quad \gamma^3 = \begin{pmatrix} 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \end{pmatrix} . \tag{3.10}$$

This particular choice is made in order to simplify the angular analysis and the radial equations. It has the advantage that charge conjugation takes the simple form

$$\psi^c = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \psi^* \ , \tag{3.11}$$

where * stands for complex conjugation as in chapter 2.

We seek a complete set of orthonormal solutions to (3.9), positive frequency with respect to Schwarzschild Killing time ∂_t. We use the separation ansatz

$$\psi_{\omega,k',m'}(t,r,\theta,\phi) = N \frac{e^{-i\omega t}}{r\left(1-\frac{2M}{r}\right)^{\frac{1}{4}}} \begin{pmatrix} F(r) Y^{k'}_{m'}(\theta,\phi) \\ G(r) Y^{k'}_{m'}(\theta,\phi) \end{pmatrix}_{\omega k'} \ , \tag{3.12}$$

where $\omega > 0$ for positive frequency modes. The angular eigenfunctions $Y^{k'}_{m'}(\theta,\phi)$ are an orthonormal set of eigenfunctions of J_3, J^2 and K where

$$\begin{aligned} J_3 &= -i\partial_\phi \ , \\ J_\pm &= e^{\pm i\phi}\left(\pm\partial_\theta - \cot\theta \frac{1}{i}\partial_\phi + \frac{1}{2}\frac{i\gamma^2\gamma^3}{\sin\theta}\right) \ , \\ K &= i\beta\gamma^1\left(\gamma^2 \frac{1}{i\sin^{\frac{1}{2}}\theta}\partial_\theta \sin^{\frac{1}{2}}\theta + \gamma^3 \frac{1}{i\sin\theta}\partial_\phi\right) \ , \end{aligned} \tag{3.13}$$

with the properties

$$\begin{aligned} J_3 Y^{k'}_{m'}(\theta,\phi) &= m' Y^{k'}_{m'}(\theta,\phi) \ , \\ J_\pm Y^{k'}_{m'}(\theta,\phi) &= [j(j+1) - m'(m'\pm 1)]^{\frac{1}{2}} Y^{k'}_{m'\pm 1}(\theta,\phi) \ , \\ K Y^{k'}_{m'}(\theta,\phi) &= k' Y^{k'}_{m'}(\theta,\phi) \ , \end{aligned} \tag{3.14}$$

where $j = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \ldots$, $m' = -j, -j+1, \ldots, j$ and $k' = \pm(j+\frac{1}{2})$. These operators are obtained from the usual angular operators J_3, $J_\pm$, K used in Minkowski space quantum field theory (i.e relative to a Cartesian set of axes) using the relevant spinor transformation which corresponds to the transformation between a Cartesian vierbein oriented relative to fixed orthogonal directions and the vierbein (3.8). This transformation is discussed in Boulware [42] (see also [43] for a discussion of these two simple choices of vierbein in the Schwarzschild geometry). J_3, for example, is the total (orbital plus spin) angular momentum about the x^3-axis and so m' takes half odd integer values. This is clear when written in the Cartesian vierbein oriented relative to fixed orthogonal directions but is not clear from the form in (3.14) as (3.14) is written with respect to vierbein (3.8). Following Boulware (see Appendix A of [42]) we may now construct explicit expressions for the $Y^{k'}_{m'}(\theta,\phi)$ eigenfunctions. We find

$$Y^{k'}_{m'}(\theta,\phi) = \frac{e^{im'\phi}}{(2\pi)^{\frac{1}{2}}}\left[\frac{(j+m')!}{(j-m')!}\right]^{\frac{1}{2}} \frac{(\tan\frac{\theta}{2})^{\sigma_2/2}}{\sin^{m'}\theta}\left(\frac{\partial}{\partial\cos\theta}\right)^{j-m'}\frac{\sin^{2j}\theta(\tan\frac{\theta}{2})^{-\sigma_2}}{2^j(j-\frac{1}{2})!}\psi_{k'}\ , \quad (3.15)$$

where $\psi_{k'} = \frac{1}{\sqrt{2}|k'|}\begin{pmatrix} |k'| \\ k' \end{pmatrix}$. Further we find the following relations which will be useful later

$$\begin{aligned} Y^{k'}_{m'}(\theta,\phi)^* &= -i(-1)^{m'-1/2}\frac{k'}{|k'|}Y^{-k'}_{-m'}(\theta,\phi)\ , && (3.16)\\ Y^{k'}_{m'}(\theta,\phi) &= -i\frac{k'}{|k'|}\sigma_2 Y^{-k'}_{m'}(\theta,\phi)\ , && (3.17)\\ Y^{k'}_{m'}(\pi-\theta,\phi+\pi) &= i(-1)^{j-1/2}\frac{k'}{|k'|}\sigma_1 Y^{k'}_{m'}(\theta,\phi)\ , && (3.18) \end{aligned}$$

where σ_j are Pauli matrices. These spinor harmonics are normalized such that

$$\int_{\text{angles}} \sin\theta\, d\theta\, d\phi\; Y^{k''}_{m''}(\theta,\phi)^\dagger Y^{k'}_{m'}(\theta,\phi) = \delta^{k''k'}\delta_{m''m'}\ . \quad (3.19)$$

For a complete set of orthonormal mode solutions we must now find a basis of solutions for the radial functions $F(r)$ and $G(r)$ which are suitably orthonormal. The radial functions satisfy

$$\left(1-\frac{2M}{r}\right)\partial_r F - i\omega F = \left(1-\frac{2M}{r}\right)^{\frac{1}{2}}\left(\frac{k'}{r}-im\right)G \ , \tag{3.20a}$$

$$\left(1-\frac{2M}{r}\right)\partial_r G + i\omega G = \left(1-\frac{2M}{r}\right)^{\frac{1}{2}}\left(\frac{k'}{r}+im\right)F \ . \tag{3.20b}$$

Following Chandrasekhar [44], we reduce the radial equations (3.20) to a pair of Schrödinger-like equations. Writing

$$\begin{pmatrix} F(r) \\ G(r) \end{pmatrix} = \begin{pmatrix} \frac{1}{2}(Z_+ + Z_-)e^{(-\frac{i}{2}\tan^{-1}(\frac{mr}{k'}))} \\ \frac{1}{2}(Z_+ - Z_-)e^{(\frac{i}{2}\tan^{-1}(\frac{mr}{k'}))} \end{pmatrix} \ , \tag{3.21}$$

we find that $Z_\pm$ satisfy

$$\left(\frac{d}{d\hat{r}_*}\mp W\right)Z_\pm = i\omega Z_\mp \ , \tag{3.22}$$

where $\hat{r}^* = r^* + \frac{1}{2\omega}\tan^{-1}(\frac{mr}{k'})$, $r^* = r + 2M\ln(|r-2M|/2M)$ and

$$W = \frac{(r^2-2Mr)^{\frac{1}{2}}(k'^2+m^2r^2)^{\frac{3}{2}}}{r^2(k'^2+m^2r^2)+\frac{k'm}{2\omega}(r^2-2Mr)} \ . \tag{3.23}$$

$Z_\pm$ hence satisfy the one-dimensional time-independent Schrödinger equations

$$\left(\frac{d^2}{d\hat{r}_*^2}+\omega^2\right)Z_\pm = V_\pm Z_\pm \ , \tag{3.24}$$

where

$$V_\pm = W^2 \pm \frac{dW}{d\hat{r}_*} \ . \tag{3.25}$$

From the usual one-dimensional scattering theory the eigenvalue spectrum for ω is seen to be continuous and consists of the entire positive real line [45].

Suppose first $\omega^2 > m^2$. The eigenvalue spectrum is then degenerate of order 2. One way we can break this degeneracy and obtain a complete set of orthonormal solutions would be to choose the solutions that have the scattering theory asymptotic form,

$$\overleftarrow{Z}_{\pm}= \begin{cases} \overleftarrow{B}_{\pm}\, e^{-i\omega\hat{r_*}} & \hat{r_*}\to -\infty \\ e^{-i(p\hat{r_*}+\frac{Mm^2}{p}\ln(\frac{\hat{r_*}}{2M}))}+ \overleftarrow{A}_{\pm}\, e^{i(p\hat{r_*}+\frac{Mm^2}{p}\ln(\frac{\hat{r_*}}{2M}))} & \hat{r_*}\to \infty \quad , \end{cases} \tag{3.26a}$$

$$\overrightarrow{Z}_{\pm}= \begin{cases} e^{i\omega\hat{r_*}}+ \overrightarrow{A}_{\pm}\, e^{-i\omega\hat{r_*}} & \hat{r_*}\to -\infty \\ \overrightarrow{B}_{\pm}\, e^{i(p\hat{r_*}+\frac{Mm^2}{p}\ln(\frac{\hat{r_*}}{2M}))} & \hat{r_*}\to \infty \quad , \end{cases} \tag{3.26b}$$

where $p = \sqrt{(\omega^2 - m^2)}$. $\overleftarrow{Z}_{\pm}$ is purely ingoing at the horizon and $\overrightarrow{Z}_{\pm}$ is purely outgoing at infinity. Considering the Wronskian between these solutions in each asymptotic region yields relations between the transmission and reflection coefficients

$$\begin{aligned} (1-\mid \overleftarrow{A}_{\pm}\mid^2) &= \frac{\omega}{p}\mid \overleftarrow{B}_{\pm}\mid^2 \ , \\ (1-\mid \overrightarrow{A}_{\pm}\mid^2) &= \frac{p}{\omega}\mid \overrightarrow{B}_{\pm}\mid^2 \ , \\ \omega\, \overleftarrow{B}_{\pm} &= p\, \overrightarrow{B}_{\pm} \ , \\ -\overrightarrow{B}_{\pm}^{\;*}\ \overleftarrow{A}_{\pm} &= \omega \overrightarrow{A}_{\pm}^{\;*}\ \overleftarrow{B}_{\pm} \ , \\ \overleftarrow{A}_{\pm}^{\;*} &= -\frac{\overrightarrow{B}_{\pm}^{\;*}\ \overrightarrow{A}_{\pm}}{\overrightarrow{B}_{\pm}} \ , \\ \mid \overrightarrow{A}_{\pm}\mid^2 &= \mid \overleftarrow{A}_{\pm}\mid^2 \ . \end{aligned} \tag{3.27}$$

From (3.22) it further follows that $\overleftarrow{B}_{+}= -\overleftarrow{B}_{-}$ and $\overrightarrow{A}_{+}= -\overrightarrow{A}_{-}$. We may therefore take for our radial modes the following basis,

$$\overset{\longleftarrow}{\begin{pmatrix} F(r) \\ G(r) \end{pmatrix}} = \sqrt{\frac{\omega}{2\pi p}} \begin{cases} \begin{pmatrix} 0 \\ \overset{\leftarrow}{B}_+ \end{pmatrix} e^{-i\omega r_*} & r_* \to -\infty \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{-iq_{\hat{r}_*}} + \begin{pmatrix} 0 \\ \overset{\leftarrow}{A}_+ \end{pmatrix} e^{iq_{\hat{r}_*}} & \hat{r_*} \to \infty \end{cases} ,$$

$$\overset{\longrightarrow}{\begin{pmatrix} F(r) \\ G(r) \end{pmatrix}} = \sqrt{\frac{1}{2\pi}} \begin{cases} \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{i\omega r_*} + \begin{pmatrix} 0 \\ \vec{A}_+ \end{pmatrix} e^{-i\omega r_*} & r_* \to -\infty \\ \begin{pmatrix} 0 \\ \vec{B}_+ \end{pmatrix} e^{iq_{\hat{r}_*}} & \hat{r_*} \to \infty \end{cases} , \tag{3.28}$$

where $q_{\hat{r}_*} = (\frac{1}{2}\tan^{-1}(\frac{mr}{k'})) + (p\hat{r_*} + \frac{Mm^2}{p}\ln(\frac{\hat{r_*}}{2M}))$. Here $\overset{\longleftarrow}{\begin{pmatrix} F(r) \\ G(r) \end{pmatrix}}$ is a purely ingoing radial mode at the horizon, while $\overset{\longrightarrow}{\begin{pmatrix} F(r) \\ G(r) \end{pmatrix}}$ is purely outgoing at infinity. However, to be able to handle the geon in subsection 3.4, we will need modes that transform simply under charge conjugation (3.11) and under J (3.5) when continued analytically into the F region. Using the properties of the reflection and transmission coefficients (3.27) and the properties of the spinor spherical harmonics (3.16)-(3.18), we find after considerable effort that a convenient set of positive frequency Boulware modes is $\{\Psi^{\pm}_{\omega,k',m'}\}$, given by

$$\Psi^{+}_{\omega,m',k'} = \frac{e^{-\frac{i\pi}{2}(j+m'+(\frac{k'}{|k'|}-1)/2)}e^{-i\omega t}}{r(1-\frac{2M}{r})^{\frac{1}{4}}} \begin{pmatrix} u(r)Y^{k'}_{m'}(\theta,\phi) \\ v(r)Y^{k'}_{m'}(\theta,\phi) \end{pmatrix}^{+}_{\omega k'} , \tag{3.29a}$$

$$\Psi^{-}_{\omega,m',k'} = \frac{e^{-\frac{i\pi}{2}(j-m'+(\frac{k'}{|k'|}-1)/2)}e^{-i\omega t}}{r(1-\frac{2M}{r})^{\frac{1}{4}}} \begin{pmatrix} u(r)Y^{k'}_{m'}(\theta,\phi) \\ v(r)Y^{k'}_{m'}(\theta,\phi) \end{pmatrix}^{-}_{\omega k'} , \tag{3.29b}$$

where the radial functions with superscript $^+$ are specified by the horizon asymptotic behaviour

$$\begin{pmatrix} u(r) \\ v(r) \end{pmatrix}^{+}_{\omega k'} = \frac{1}{\sqrt{4\pi}} \left\{ \sqrt{1+\sqrt{1-|\vec{A}\,|^2}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{i\omega r_*} \right.$$

$$\left. + \frac{\vec{A}_+}{\sqrt{1+\sqrt{1-|\vec{A}\,|^2}}} \begin{pmatrix} 0 \\ 1 \end{pmatrix} e^{-i\omega r_*} \right\} , \qquad \hat{r}^* \to -\infty \ , \tag{3.30a}$$

and

$$\begin{pmatrix} u(r) \\ v(r) \end{pmatrix}^{-}_{\omega k'} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} u(r) \\ v(r) \end{pmatrix}^{+*}_{\omega k'} . \tag{3.30b}$$

The key property for charge conjugation is (3.30b). The modes $\{\Psi^{\pm}_{\omega,k',m'}\}$ are complete for $\omega^2 > m^2$ and delta-orthonormal in the Dirac inner product

$$\langle \psi_1, \psi_2 \rangle = \int_{\text{angles}} \sin\theta \, d\theta \, d\phi \int_{2M}^{\infty} dr \, \frac{r^2}{\left(1-\frac{2M}{r}\right)^{\frac{1}{2}}} \psi_1^\dagger \psi_2 \ , \tag{3.31}$$

taken on a constant t hypersurface.

Suppose then $0 < \omega^2 < m^2$ in (3.24). There is now only one linearly independent delta-normalizable solution for each ω. This solution vanishes at infinity and has at the horizon the behaviour

$$Z_{\pm} = a_{\pm} \cos(\omega \hat{r}^* + \delta_{\pm}) \ , \quad \hat{r}^* \to -\infty \ , \tag{3.32}$$

where $a_{\pm}$, and $\delta_{\pm}$ are real constants. Physically these solutions correspond to particles that do not reach infinity. Proceeding as above, we find that a convenient set of positive frequency Boulware modes, complete for $0 < \omega^2 < m^2$ and delta-orthonormal in the Dirac inner product (3.31), is

$$\psi_{\omega,k',m'}(t,r,\theta,\phi) = e^{-\frac{i\pi}{2}(j+|m'|+(1-\frac{k'}{|k'|})/2)} \frac{e^{-i\omega t}}{r(1-\frac{2M}{r})^{\frac{1}{4}}} \begin{pmatrix} F(r) Y^{k'}_{m'}(\theta,\phi) \\ G(r) Y^{k'}_{m'}(\theta,\phi) \end{pmatrix}_{\omega k'} , \tag{3.33}$$

where $0 < \omega < m$ and the radial functions are specified by the horizon asymptotic behaviour

$$\left(\begin{array}{c} F(r) \\ G(r) \end{array}\right)_{\omega k'} = \frac{1}{\sqrt{2\pi}} \left\{ \left(\begin{array}{c} e^{i\delta_+} \\ 0 \end{array}\right) e^{i\omega r_*} + \left(\begin{array}{c} 0 \\ e^{-i\delta_+} \end{array}\right) e^{-i\omega r_*} \right\} , \quad \hat{r}^* \mapsto -\infty . \tag{3.34}$$

Up to this point we have used the Schwarzschild vierbein (3.8). To make contact with the geon in subsection 3.4, we need to express the modes in a vierbein that is regular at the horizons. We therefore now transform our modes to the Kruskal vierbein (3.6) by the spinor transformation $\psi \mapsto e^{\frac{t}{8M}\gamma^0\gamma^1}\psi$. We suppress the explicit transformed expressions and continue to use the same symbols for the mode functions.

We are now ready to quantize. The field is expanded in our orthonormal modes and their charge conjugates as

$$\begin{aligned} \Psi = & \sum_{k',m'} \int_0^m d\omega \left(a_{\omega,k',m'} \psi_{\omega,k',m'} + b^\dagger_{\omega,k',m'} \psi^c_{\omega,k',m'} \right) \\ & + \sum_{k',m'} \int_m^\infty d\omega \Big(a_{+,\omega,k',m'} \Psi^+_{\omega,k',m'} + a_{-,\omega,k',m'} \Psi^-_{\omega,k',m'} \\ & \qquad + b^\dagger_{+,\omega,k',m'} \Psi^{+,c}_{\omega,k',m'} + b^\dagger_{-,\omega,k',m'} \Psi^{-,c}_{\omega,k',m'} \Big) , \end{aligned} \tag{3.35}$$

where the annihilation and creation operators satisfy the usual anticommutation relations. The vacuum annihilated by the annihilation operators is the Boulware vacuum $|0_B\rangle$

$$a_{\omega,k',m'}|0_B\rangle = b_{\omega,k',m'}|0_B\rangle = a_{\pm,\omega,k',m'}|0_B\rangle = b_{\pm,\omega,k',m'}|0_B\rangle = 0 , \quad \forall \omega, k', m' . \tag{3.36}$$

The Boulware vacuum here is analogous to the Rindler vacuum $|0_{R_-}\rangle$ in section 2.5. $|0_B\rangle$ is by construction the state void of particles defined with respect to the Schwarzschild Killing time. It is the natural vacuum for static observers in the exterior region R.

3.4 The Hartle-Hawking-like vacuum and Bogolubov transformation on the geon

In this section we decompose the geon Hartle-Hawking-like vacuum into Boulware excitations. We use the analytic continuation method, following closely subsection 2.5.

The Hartle-Hawking vacuum on the Kruskal manifold is defined by mode functions that are purely positive frequency with respect to the horizon generators and hence analytic in the lower half-plane in the complexified Kruskal time T. It follows that on Kruskal we can construct W-modes whose vacuum is the Hartle-Hawking vacuum by analytically continuing the Boulware-modes across the horizons in the lower half-plane in T. The quotient from Kruskal to the geon defines in each spin structure on the geon the Hartle-Hawking-like vacuum $|0_G\rangle$, by restriction to the Kruskal W-modes that are invariant under the map J (3.5). Our task is to find these modes.

We begin by taking our positive frequency ($\omega > 0$) modes (3.29a), (3.29b) and (3.33) written in the Kruskal vierbein (3.6) and continuing them across the future horizon into the interior black hole region F in the lower half complex T-plane. Let us consider a general positive frequency mode solution to equation (3.9) near the horizon $r = 2M$. In the Kruskal vierbein this can take the form

$$\psi^R(t,r,\theta,\phi) = N\frac{e^{-i\omega t}}{r(1-\frac{2M}{r})^{\frac{1}{4}}}\left\{\begin{pmatrix} A \\ 0 \end{pmatrix} e^{i\omega r_* + \frac{t}{8M}} + \begin{pmatrix} 0 \\ B \end{pmatrix} e^{-i\omega r_* - \frac{t}{8M}}\right\} Y^{k'}_{m'}(\theta,\phi) \ , \tag{3.37}$$

where A and B are arbitrary constants, N is a normalization factor and $\omega > 0$. Writing this in Kruskal coordinates, it becomes

$$\psi^R(T,X,\theta,\phi) = N\left[\begin{pmatrix} A \\ 0 \end{pmatrix}_{\omega k'} (X-T)^{4M\omega i}\left(\frac{X+T}{X-T}\right)^{\frac{1}{4}} \right.$$
$$\left. + \begin{pmatrix} 0 \\ B \end{pmatrix}_{\omega k'} (X+T)^{-4M\omega i}\left(\frac{X-T}{X+T}\right)^{\frac{1}{4}}\right]\frac{1}{r}\frac{(r/2M)^{1/2}e^{r/8M}}{(X^2-T^2)^{\frac{1}{4}}} Y^{k'}_{m'}(\theta,\phi) \ . \tag{3.38}$$

Continuation to F in the lower half complex T-plane is then implemented by the substitutions $(X-T)^{1/4} \mapsto (T-X)^{1/4}e^{i\pi/4}$ and $\ln(X-T) \mapsto \ln(T-X)+i\pi$, under which $\psi^R \mapsto \psi^F$ where

$$\psi^F(T,X,\theta,\phi) = N\left[\begin{pmatrix} A \\ 0 \end{pmatrix}_{\omega k'} (T-X)^{4M\omega i} e^{-4M\omega\pi - \frac{i\pi}{2}} \left(\frac{X+T}{T-X}\right)^{\frac{1}{4}}\right.$$
$$\left. + \begin{pmatrix} 0 \\ B \end{pmatrix}_{\omega k'} (X+T)^{-4M\omega i} \left(\frac{T-X}{X+T}\right)^{\frac{1}{4}}\right] \frac{1}{r}\frac{(r/2M)^{1/2}e^{r/8M}}{(T^2-X^2)^{\frac{1}{4}}} Y^{k'}_{m'}(\theta,\phi)\ . \tag{3.39}$$

We need to build from (3.39) modes that are well-defined in the F region of the geon by the method of images. If we were working in the rotating Kruskal vierbein which is invariant under the involution J the method of images could be applied directly and we would have

$$W^F(T,X,\theta,\phi) = \psi^F(T,X,\theta,\phi) + \epsilon\psi^F(T,-X,\pi-\theta,\phi+\pi)\ , \tag{3.40}$$

where $\epsilon = 1, (-1)$ labels fields with periodic (antiperiodic) boundary conditions respectively. However we are instead working in the standard Kruskal vierbein which is not invariant under J. The spinor transformation associated with the transformation between the rotating Kruskal vierbein and the standard one (a rotation by $-\pi$ in the (X,θ)-plane as $\phi \mapsto \phi+\pi$) is given by

$$W^F(T,X,\theta,\phi) \mapsto e^{\frac{\phi\gamma^1\gamma^2}{2}} W^F(T,X,\theta,\phi) = W^F_s(T,X,\theta,\phi)\ . \tag{3.41}$$

Now it is easy to show that in terms of this standard Kruskal vierbein

$$W^F_s(T,X,\theta,\phi) = \psi^F_s(T,X,\theta,\phi) + \epsilon e^{-\frac{\pi\gamma^1\gamma^2}{2}} \psi^F_s(T,-X,\pi-\theta,\phi+\pi)\ . \tag{3.42}$$

This is the expression we require our W-modes on the geon to satisfy when working in this non-invariant vierbein. Now applying (3.39) to (3.42) we find positive frequency W-modes on the geon in F near $r = 2M$ given by

$$\begin{aligned} W^F(T,X,\theta,\phi) &= \psi^F(T,X,\theta,\phi) \\ &\quad +\epsilon i(-1)^j N \left[\begin{pmatrix} 0 \\ A \end{pmatrix}_{\omega k'} (T+X)^{4M\omega i} e^{-4M\omega\pi - \frac{i\pi}{2}} \left(\frac{T-X}{T+X}\right)^{\frac{1}{4}} \right. \\ &\quad \left. + \begin{pmatrix} B \\ 0 \end{pmatrix}_{\omega k'} (T-X)^{-4M\omega i} \left(\frac{T+X}{T-X}\right)^{\frac{1}{4}} \right] \frac{1}{r} \frac{(r/2M)^{1/2} e^{r/8M}}{(T^2-X^2)^{\frac{1}{4}}} Y_{m'}^{-k'}(\theta,\phi)\ . \end{aligned} \tag{3.43}$$

On continuation back to the right hand exterior in the lower half complex T-plane, these modes become

$$\begin{aligned} W^R(T,X,\theta,\phi) &= \psi^R(T,X,\theta,\phi) \\ &\quad +\epsilon(-1)^{j+1} e^{-4M\omega\pi} N \left[\begin{pmatrix} 0 \\ -A \end{pmatrix}_{\omega k'} (T+X)^{4M\omega i} \left(\frac{X-T}{T+X}\right)^{\frac{1}{4}} \right. \\ &\quad \left. + \begin{pmatrix} B \\ 0 \end{pmatrix}_{\omega k'} (X-T)^{-4M\omega i} \left(\frac{T+X}{X-T}\right)^{\frac{1}{4}} \right] \frac{1}{r} \frac{(r/2M)^{1/2} e^{r/8M}}{(X^2-T^2)^{\frac{1}{4}}} Y_{m'}^{-k'}(\theta,\phi)\ . \end{aligned} \tag{3.44}$$

We are now ready to specialise to the Boulware modes found in the previous section. For $\omega > m$ comparing (3.37) and (3.30a,3.30b), and making use of the relations (3.16)-(3.18) we find in the R region a complete set of W-modes given by

$$\begin{aligned} W^+_{\omega,k',m'}(t,r,\theta,\phi) &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}} \left(e^{2\pi M\omega} \Psi^+_{\omega,k',m'} + \epsilon e^{-2\pi M\omega} \Psi^{-,c}_{\omega,k',-m'} \right)\ , \\ W^-_{\omega,k',m'}(t,r,\theta,\phi) &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}} \left(e^{2\pi M\omega} \Psi^-_{\omega,k',m'} - \epsilon e^{-2\pi M\omega} \Psi^{+,c}_{\omega,k',-m'} \right)\ . \end{aligned} \tag{3.45}$$

(Note the similarity between (3.45) and (2.72)). These modes are delta-orthonormal with respect to the inner product (3.31). For $0 < \omega < m$, comparing (3.37) and (3.34) we find

W-modes given by

$$W_{\omega,k',m'}(t,r,\theta,\phi) = \frac{1}{\sqrt{2\cosh(4\pi M\omega)}}\left(e^{2\pi M\omega}\psi_{\omega,k',m'} + \epsilon\frac{m'}{|m'|}e^{-2\pi M\omega}\psi^{c}_{\omega,k',-m'}\right) \ . \quad (3.46)$$

again orthonormal with respect to the Kruskal inner product (3.31). It can be verified that the factors $m'/|m'|$ appearing in (3.46) cannot be absorbed into the phase factors of the Boulware modes.

On the geon, the expansion of the field in the W-modes reads

$$\begin{aligned}\Psi \quad = \quad & \sum_{k',m'}\int_0^m d\omega\left(c_{\omega,k',m'}W_{\omega,k',m'} + d^\dagger_{\omega,k',m'}W^c_{\omega,k',m'}\right) \\ & + \sum_{k',m'}\int_m^\infty d\omega\left(c_{+,\omega,k',m'}W^{+}_{\omega,k',m'} + c_{-,\omega,k',m'}W^{-}_{\omega,k',m'}\right. \\ & \qquad\qquad \left. + d^\dagger_{+,\omega,k',m'}W^{+,c}_{\omega,k',m'} + d^\dagger_{-,\omega,k',m'}W^{-,c}_{\omega,k',m'}\right) \ , \qquad (3.47)\end{aligned}$$

with the usual anticommutation relations imposed on the creation and annihilation operators. We denote by $|0_G\rangle$ the state annihilated by all the annihilation operators,

$$c_{\omega,k',m'}|0_G\rangle = d_{\omega,k',m'}|0_G\rangle = c_{\pm,\omega,k',m'}|0_G\rangle = d_{\pm,\omega,k',m'}|0_G\rangle = 0 \ , \quad \forall\omega,k',m' \ . \quad (3.48)$$

$|0_G\rangle$ is the Kruskal-like vacuum on the geon, induced by the usual Hartle-Hawking vacuum on Kruskal. Comparison of (3.35) and (3.47) gives the Bogolubov transformation

$$\begin{aligned}
a_{\omega,k',m'} &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}}\left(e^{2\pi M\omega}c_{\omega,k',m'} + \epsilon\frac{m'}{|m'|}e^{-2\pi M\omega}d^{\dagger}_{\omega,k',-m'}\right) \ , \\
b^{\dagger}_{\omega,k',m'} &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}}\left(e^{2\pi M\omega}d^{\dagger}_{\omega,k',m'} + \epsilon\frac{m'}{|m'|}e^{-2\pi M\omega}c_{\omega,k',-m'}\right) \ , \\
a_{+,\omega,k',m'} &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}}\left(e^{2\pi M\omega}c_{+,\omega,k',m'} - \epsilon e^{-2\pi M\omega}d^{\dagger}_{-,\omega,k',-m'}\right) \ , \\
a_{-,\omega,k',m'} &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}}\left(e^{2\pi M\omega}c_{-,\omega,k',m'} + \epsilon e^{-2\pi M\omega}d^{\dagger}_{+,\omega,k',-m'}\right) \ , \\
b^{\dagger}_{+,\omega,k',m'} &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}}\left(e^{2\pi M\omega}d^{\dagger}_{+,\omega,k',m'} - \epsilon e^{-2\pi M\omega}c_{-,\omega,k',-m'}\right) \ , \\
b^{\dagger}_{-,\omega,k',m'} &= \frac{1}{\sqrt{2\cosh(4\pi M\omega)}}\left(e^{2\pi M\omega}d^{\dagger}_{-,\omega,k',m'} + \epsilon e^{-2\pi M\omega}c_{+,\omega,k',-m'}\right) \ .
\end{aligned} \tag{3.49}$$

For the expectation values of Boulware mode number operators, (3.49) gives e.g.

$$\langle 0_G|a^{\dagger}_{+,\omega,k',m'}a_{+,\omega',k'',m''}|0_G\rangle = \frac{1}{(e^{8\pi M\omega}+1)}\delta(\omega-\omega')\delta_{k'k''}\delta_{m'm''} \ , \tag{3.50}$$

and a discussion similar to that given on M_- follows. Note in particular that the geon spin structure does not show up in the expectation value. The number operator expectation value does not distinguish between the state $|0_G\rangle$ on the geon and the usual Hartle-Hawking state on Kruskal, it is representative of a thermal bath at temperature $T = (8\pi M)^{-1}$.

Expressing $|0_G\rangle$ in terms of $|0_B\rangle$, we find

$$\begin{aligned}
|0_G\rangle = &\prod_{\substack{0<\omega<m\\ k',m'}} \frac{1}{(e^{-8\pi M\omega}+1)^{\frac{1}{2}}}\sum_{q=0,1}\left(\frac{\epsilon m'}{|m'|}\right)^{q} e^{-4\pi M\omega q}(a^{\dagger}_{\omega,k',m'}b^{\dagger}_{\omega,k',-m'})^{q}|0_B\rangle \\
&\times \prod_{\substack{\omega>m\\ \sigma,k',m'}} \frac{1}{(e^{-8\pi M\omega}+1)^{\frac{1}{2}}}\sum_{q=0,1}(-\epsilon)^{q}e^{-4\pi M\omega q}|q\rangle_{\sigma,+,\omega,k',m'}|q\rangle_{-\sigma,-,\omega,k',-m'} \ ,
\end{aligned} \tag{3.51}$$

where $\sigma = +1(-1)$ and

$$\begin{aligned}
|q\rangle_{+,+,\omega,k',m'} &= (a^{\dagger}_{+,\omega,k',m'})^{q}|0_B\rangle \quad |q\rangle_{+,-,\omega,k',m'} = (a^{\dagger}_{-,\omega,k',m'})^{q}|0_B\rangle \ , \\
|q\rangle_{-,+,\omega,k',m'} &= (b^{\dagger}_{+,\omega,k',m'})^{q}|0_B\rangle \quad |q\rangle_{-,-,\omega,k',m'} = (b^{\dagger}_{-,\omega,k',m'})^{q}|0_B\rangle \ .
\end{aligned} \tag{3.52}$$

A comparison of (3.51) and (2.77) shows that $|0_G\rangle$ is closely similar to the state $|0_-\rangle$ on M_-. The Boulware modes are correlated in pairs in the Hartle-Hawking-like vacuum state. The correlations now are between a particle in the right hand exterior region and an antiparticle in the same exterior with opposite m' eigenvalue. $|0_G\rangle$ does not appear thermal to generic static observers in R, but it appears thermal in the standard Hawking temperature $(8\pi M)^{-1}$ near the infinity when probed by operators that only couple to one member of each correlated pair in (3.51), such as operators that only couple to a definite sign of the angular momentum quantum number m'. In particular, number operator expectation values are thermal, as seen above, and the isometry arguments of [22] show that the experiences of any static observer become asymptotically thermal in the large $|t|$ limit.

The explicit appearance of ϵ in (3.51) shows that the non-thermal correlations in $|0_G\rangle$ reveal the geon spin structure to an observer in R. This is a phenomenon that could not have been anticipated just from the geometry of R, which in its own right has only one spin structure.

Chapter 4

Stress tensor on M_0, M_- and Minkowski space with plane boundary

4.1 Introduction

In this chapter the expectation values of the energy-momentum tensor in the Minkowski-like vacua on M_0, M_- and Minkowski space with an infinite plane boundary are presented. In section 4.2 we calculate $\langle 0|T_{\mu\nu}|0\rangle$ for the massive scalar field on all three spacetimes. The values on M_0 agree with the literature [46] while to the knowledge of the author those on M_- are new. In the massless limit these values agree with those in [22,36,47]. Various limits are discussed and qualitative differences between the massive and massless values noted. In section 4.3 we calculate $\langle 0|T_{\mu\nu}|0\rangle$ for left and right handed 2-component Weyl spinors on the spacetimes M_0 and M_-. It is shown that the values are independent of handedness but are dependent on the spin structure. Limits are discussed and the differences between $\langle 0|T_{\mu\nu}|0\rangle$ on the two spacetimes noted. Finally in section 4.4 we calculate $\langle 0|T_{\mu\nu}|0\rangle$ for massive 4-component spinors on M_0 and M_-. The results on M_0 agree with those in [48] [1]. The results on M_- are new. Again various limits are discussed. In the massless limit the results agree with those in section 4.3 (see also [23, 35]).

4.2 Massive scalar field

The stress tensor for massive scalar fields in flat spacetimes with non-trivial topology was considered by Tanaka and Hiscock in [46]. In particular they consider $\langle 0|T_{\mu\nu}|0\rangle$ on flat spacetimes with topology and product space geometry $\mathbb{R}^3 \times S^1$ (which is denoted M_0 here), $\mathbb{R}^2 \times T^2$ and $\mathbb{R}^1 \times T^3$. It is seen in [46] that the magnitude of the energy density decreases with an increasing field mass. Here we reproduce the result on M_0 and present the expectation values on M_- and on Minkowski space with a single infinite plane boundary at $x = 0$.

The energy-momentum tensor for the massive scalar field in a general 4-dimensional curved spacetime $\tilde{M}$ in our conventions is [1]

$$\begin{aligned} T_{\mu\nu} &= (1-2\xi)\phi_{;\mu}\phi_{;\nu} + \left(2\xi - \frac{1}{2}\right) g_{\mu\nu} g^{\rho\sigma}\phi_{;\rho}\phi_{;\sigma} - 2\xi\phi_{;\mu\nu}\phi \\ &\quad + \frac{1}{2}\xi g_{\mu\nu}\phi\Box\phi - \xi\left[R_{\mu\nu} - \frac{1}{2}R g_{\mu\nu} + \frac{3}{2}\xi R g_{\mu\nu}\right]\phi^2 \\ &\quad + \frac{1}{2}\left[1-3\xi\right] m^2 g_{\mu\nu}\phi^2 \ , \end{aligned} \tag{4.1}$$

which, applying the field equation $[\Box + m^2 + \xi R]\phi = 0$, may be written on a flat spacetime ($R = 0$) as

$$\begin{aligned} T_{\mu\nu} &= (1-2\xi)\phi_{;\mu}\phi_{;\nu} + \left(2\xi - \frac{1}{2}\right) g_{\mu\nu} g^{\rho\sigma}\phi_{;\rho}\phi_{;\sigma} - 2\xi\phi_{;\mu\nu}\phi \\ &\quad + 2\xi g_{\mu\nu}\phi\Box\phi + \frac{1}{2}m^2 g_{\mu\nu}\phi^2 \ , \end{aligned} \tag{4.2}$$

where ξ gives the coupling to the gravitational field ($\xi = 0$ for minimal coupling and $\xi = 1/6$ for conformal coupling). By the point splitting technique [49], where we split the points in the quadratic expressions above and take the coincidence limit at the end, the renormalized expectation value of $T_{\mu\nu}$ in vacuum state $|0\rangle$ may be written as

---

[1] Except for an extra factor of 2 here as in [48] Majorana spinors are considered.

$$\begin{aligned}\langle 0|T_{\mu\nu}|0\rangle &= \frac{1}{2}\lim_{x'\to x}\left[(1-2\xi)\nabla_\mu\nabla_{\nu'} + \left(2\xi-\frac{1}{2}\right)g_{\mu\nu}\nabla_\alpha\nabla^{\alpha'} - 2\xi\nabla_\mu\nabla_\nu\right. \\ &\quad \left. + 2\xi g_{\mu\nu}\nabla_\alpha\nabla^\alpha + \frac{1}{2}m^2 g_{\mu\nu}\right]\left(G^{(1)}_{\tilde{M}}(x,x') - G^{(1)}(x,x')\right) , \qquad (4.3)\end{aligned}$$

where $G^{(1)}_{\tilde{M}}(x,x') = \langle 0|\{\phi(x),\phi(x')\}|0\rangle = G^{+}_{\tilde{M}}(x,x') + G^{-}_{\tilde{M}}(x,x')$ is the scalar Hadamard function in $\tilde{M}$ and $G^{(1)}(x,x')$ is the Hadamard function in Minkowski space.

On Minkowski space $G^{(1)}(x,x')$ may be calculated by considering the expansion of the field in a complete set of mode solutions to the Klein Gordon equation

$$\phi(t,\mathbf{x}) = \frac{1}{(2\pi)^{3/2}}\int\frac{d^3k}{(2\omega)^{1/2}}\left(a(\mathbf{k})e^{-i(\omega t-\mathbf{k}\cdot\mathbf{x})} + a^\dagger(\mathbf{k})e^{i(\omega t-\mathbf{k}\cdot\mathbf{x})}\right) , \qquad (4.4)$$

where $\omega = \left(\mathbf{k}^2+m^2\right)^{1/2}$. Then

$$\begin{aligned}\langle 0|\{\phi(x),\phi(x')\}|0\rangle &= \frac{1}{(2\pi)^3}\int\frac{d^3k}{2\omega}e^{-i(\omega(t-t')-\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}'))} \\ &\quad + \frac{1}{(2\pi)^3}\int\frac{d^3k}{2\omega}e^{i(\omega(t-t')-\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}'))} , \qquad (4.5)\end{aligned}$$

where we have made use of the anticommutation relations for the annihilation and creation operators. The integrals may now be performed by transforming to spherical coordinates in k-space. After doing the angular integral we find

$$\begin{aligned}\langle 0|\{\phi(x),\phi(x')\}|0\rangle &= \frac{1}{(2\pi)^2 R}\int_0^\infty dk\,\frac{k}{(k^2+m^2)^{1/2}}\sin(kR)\left[e^{-i\left(k^2+m^2\right)^{1/2}(t-t')}\right. \\ &\quad \left. + e^{i\left(k^2+m^2\right)^{1/2}(t-t')}\right] , \qquad (4.6)\end{aligned}$$

where $R = |\mathbf{x}-\mathbf{x}'|$. (4.6) may be written as

$$\begin{aligned}\langle 0|\{\phi(x),\phi(x')\}|0\rangle &= \frac{-1}{8\pi^2 R}\partial_R\left[\int_{-\infty}^\infty dk\,\frac{1}{(k^2+m^2)^{1/2}}\left(e^{-i\left(\left(k^2+m^2\right)^{1/2}(t-t')-kR\right)}\right.\right. \\ &\quad \left.\left. + e^{i\left(\left(k^2+m^2\right)^{1/2}(t-t')+kR\right)}\right)\right] , \qquad (4.7)\end{aligned}$$

and the above integrals may be performed by changing coordinates to $k = m\sinh\theta$, $\omega = m\cosh\theta$ and making use of the well known integral representations for the Bessel functions (see e.g [50]). The result is (see appendix C of [51])

$$\begin{aligned} G^{(1)}(x,x') &= \frac{1}{4\pi R}\partial_R Y_0\left(m\sqrt{(t-t')^2-R^2}\right)\Theta\left((t-t')^2-R^2\right) \\ &\quad -\frac{1}{2\pi^2 R}\partial_R K_0\left(m\sqrt{R^2-(t-t')^2}\right)\Theta\left(R^2-(t-t')^2\right) \ . \end{aligned} \tag{4.8}$$

Now we note that $\partial_z F_0(z) = -F_1(z)$ for $F = K, Y$ and so [2]

$$\begin{aligned} G^{(1)}(x,x') &= \frac{m}{4\pi((t-t')^2-R^2)^{1/2}}Y_1\left(m\sqrt{(t-t')^2-R^2}\right)\Theta\left((t-t')^2-R^2\right) \\ &\quad +\frac{m}{2\pi^2\left(R^2-(t-t')^2\right)^{1/2}}K_1\left(m\sqrt{R^2-(t-t')^2}\right)\Theta\left(R^2-(t-t')^2\right) \ , \end{aligned} \tag{4.9}$$

where Y, K are Bessel functions.

The Hadamard function on the quotient spaces M_0 and M_- may be found by the method of images [35, 36],[3]

$$G^{(1)}_{M_0}(x,x') = \langle 0_0|\{\phi(x),\phi(x')\}|0_0\rangle = \sum_{n\in\mathbb{Z}}\eta^n G^{(1)}\left(x, J_0^n x'\right) \ , \tag{4.10}$$

$$G^{(1)}_{M_-}(x,x') = \langle 0_-|\{\phi(x),\phi(x')\}|0_-\rangle = \sum_{n\in\mathbb{Z}}\rho^n G^{(1)}\left(x, J_-^n x'\right) \ , \tag{4.11}$$

where $J_0 : (t,x,y,z) \mapsto (t,x,y,z+2a)$, $J_- : (t,x,y,z) \mapsto (t,-x,-y,z+a)$ and $\eta = 1,(-1)$, $\rho = 1,(-1)$ label untwisted and (twisted) fields respectively. The calculation of $\langle 0|T_{\mu\nu}|0\rangle$ is now reduced to that of finding derivatives of these Hadamard functions and applying (4.3) to (4.10) and (4.11). Next we present the results.

---

[2] Note the sign difference in the second term from that in Fullings book [2] p85 which is the reference used by Tanaka and Hiscock [46]. Further note the typographical error in [46] where the Y Bessel has been replaced by the I Bessel. This does not affect the results in [46].

[3] It can be shown that (4.10) and (4.11) may be obtained by expanding the fields in modes, obtained from an image sum, proving that they are true Green's functions, that is, they are expectation values of the operators in a quantum state.

4.2.1 M_0

For M_0, in the coincidence limit $G^{(1)}_{M_0}(x,x')$ becomes a function only of $(2na)^2$, which is positive. This means that only the K Bessel term in (4.9) contributes. Further we note that $\lim_{x'\to x}\partial_{x^\mu}G^{(1)}_{M_0}(x,x') = -\lim_{x'\to x}\partial_{x'^\mu}G^{(1)}_{M_0}(x,x')$. These observations simplify the calculations somewhat. The result is

$$\begin{aligned}\langle 0_0|T_{tt}|0_0\rangle = -\langle 0_0|T_{xx}|0_0\rangle = -\langle 0_0|T_{yy}|0_0\rangle &= -\sum_{n=1}^{\infty}\eta^n\frac{m^2}{2\pi^2(2na)^2}K_2(2mna)\ ,\\ \langle 0_0|T_{zz}|0_0\rangle &= \sum_{n=1}^{\infty}\eta^n\left[\frac{m^2}{2\pi^2(2na)^2}K_2(2mna)-\frac{m^3}{2\pi^2(2na)}K_3(2mna)\right]\ ,\end{aligned} \tag{4.12}$$

with all other terms vanishing. The results agree with [46]. The massless limit may be easily checked by noting that near $z=0$

$$K_\nu(z)=\frac{\Gamma(\nu)}{2}\left(\frac{z}{2}\right)^{-\nu}\ , \tag{4.13}$$

when the real part of ν is positive [50]. These agree with [22,47] and also with [36], the extra factor of 4 in [36] arising because the authors consider a multiplet of 2 complex massless scalar fields. The leading corrections for small mass are of the order $O\left(m^2\right)$. $\langle 0_0|T_{\mu\nu}|0_0\rangle \to 0$ exponentially in the large mass and large a limits, in contrast to the massless case where $\langle 0_0|T_{\mu\nu}|0_0\rangle$ vanishes as $O\left(a^{-4}\right)$.

4.2.2 M_-

For M_- in the coincidence limit $G^{(1)}_{M_-}(x,x')$ becomes a function only of $(x-(-1)^n x)^2 + (y-(-1)^n y)^2 + (na)^2$, which is positive. Again only the K Bessel term in (4.9) contributes. Further we note the following

$$\begin{aligned}\lim_{x'\to x}\partial_\mu G^{(1)}_{M_-}(x,x') &= -\lim_{x'\to x}\partial_{\mu'}G^{(1)}_{M_-}(x,x')\ \text{ for } \mu\in\{t,z\}\ ,\\ \lim_{x'\to x}\partial_\mu G^{(1)}_{M_-}(x,x') &= -(+)\lim_{x'\to x}\partial_{\mu'}G^{(1)}_{M_-}(x,x')\ \text{ when } n \text{ even (odd) and } \mu\in\{x,y\}\ .\end{aligned} \tag{4.14}$$

As $G^{(1)}_{M_-}(x,x') = \sum_{n\in\mathbb{Z}}\rho^n G^{(1)}\left(x,J_-^n x'\right)$, where $\rho = +1,(-1)$ labels untwisted (twisted) fields, the result here may be split into two parts where the part coming from the even terms in the sum above leads to the same expectation values as on M_0 for untwisted fields. Therefore we write $\langle 0_-|T_{\mu\nu}|0_-\rangle = \langle 0_0|T_{\mu\nu}|0_0\rangle_{(\eta=1)} + \rho\langle 0_-|T_{\mu\nu}|0_-\rangle_{\text{odd}}$ where we find

$$\begin{aligned}
\langle 0_-|T_{tt}|0_-\rangle_{\text{odd}} &= \sum_{n\in\mathbb{Z}}\left[-(4\xi-1)\frac{m^3}{4\pi^2\sigma_n}K_3(m\sigma_n)\left[1-\frac{(2na+a)^2}{\sigma_n^2}\right]\right.\\
&\qquad\left.+\left(2\xi-\frac{3}{4}\right)\frac{m^2}{\pi^2\sigma_n^2}K_2(m\sigma_n)\right]\ ,\\
\langle 0_-|T_{xx}|0_-\rangle_{\text{odd}} &= \sum_{n\in\mathbb{Z}}\left[(4\xi-1)\frac{m^3y^2}{\pi^2\sigma_n^3}K_3(m\sigma_n)-\left(2\xi-\frac{1}{2}\right)\frac{m^2}{2\pi^2\sigma_n^2}K_2(m\sigma_n)\right]\ ,\\
\langle 0_-|T_{yy}|0_-\rangle_{\text{odd}} &= \sum_{n\in\mathbb{Z}}\left[(4\xi-1)\frac{m^3x^2}{\pi^2\sigma_n^3}K_3(m\sigma_n)-\left(2\xi-\frac{1}{2}\right)\frac{m^2}{2\pi^2\sigma_n^2}K_2(m\sigma_n)\right]\ ,\\
\langle 0_-|T_{zz}|0_-\rangle_{\text{odd}} &= \sum_{n\in\mathbb{Z}}\left[\frac{m^3}{4\pi^2\sigma_n}K_3(m\sigma_n)\left[(4\xi-1)-\frac{4\xi(2na+a)^2}{\sigma_n^2}\right]\right.\\
&\qquad\left.-\left(2\xi-\frac{3}{4}\right)\frac{m^2}{\pi^2\sigma_n^2}K_2(m\sigma_n)\right]\ ,\\
\langle 0_-|T_{xy}|0_-\rangle_{\text{odd}} &= \sum_{n\in\mathbb{Z}}(1-4\xi)\frac{m^3xy}{\pi^2\sigma_n^3}K_3(m\sigma_n)\ ,
\end{aligned}\tag{4.15}$$

with $\sigma_n = \left((2x)^2+(2y)^2+(2na+a)^2\right)^{1/2}$, and the sum is over all n including $n=0$. Other components vanish. Again it is a simple matter to check the massless limit here. The results agree with those of [22] and [36] in this limit. The leading correction for small mass is $O\left(m^2\right)$, and $\langle 0_-|T_{\mu\nu}|0_-\rangle$ vanishes exponentially in the large mass and, for non zero mass, the large a limits. The difference between $\langle 0_-|T_{\mu\nu}|0_-\rangle$ on M_- and $\langle 0_0|T_{\mu\nu}|0_0\rangle$ on M_0 (with $\eta=1$) vanishes exponentially as $r^2 := x^2+y^2\to\infty$. This behaviour is qualitatively different to the massless case where the difference vanishes as $O\left(r^{-3}\right)$ [22]. The result in the massive case is to our knowledge new.

4.2.3 Minkowski space with infinite plane boundary

In this subsection we consider $\langle 0_B|T_{\mu\nu}|0_B\rangle$ in 4-dimensional Minkowski space with an infinite plane boundary at $x=0$. While the stress-energy for a massless field is well known (see e.g. [1,27]), the massive results are to our knowledge new. There is some similarity with M_- as both spaces may be considered as quotients of Minkowski space with the quotient group including a reflection in x about $x = 0$.

Again here the scalar Hadamard function is given by the method of images,

$$G_B^{(1)}(x, x') = \langle 0_B | \{\phi(x)\phi(x')\} | 0_B \rangle = G^{(1)}(x, x') + \eta G^{(1)}(x, J_B x') , \tag{4.16}$$

where $J_B : (t, x, y, z) \mapsto (t, -x, y, z)$ and $\eta = +1, (-1)$ labels Neumann (Dirichlet) boundary conditions on the plate. From (4.3) the result is

$$\begin{aligned} \langle 0_B | T_{tt} | 0_B \rangle &= -\langle 0_B | T_{yy} | 0_B \rangle = -\langle 0_B | T_{zz} | 0_B \rangle \\ &= \eta \left[\frac{(1-4\xi)m^3}{4\pi^2 |2x|} K_3(m|2x|) + \frac{(2\xi - 1)m^2}{8\pi^2 x^2} K_2(m|2x|) \right] , \\ \langle 0_B | T_{xx} | 0_B \rangle &= 0 . \end{aligned} \tag{4.17}$$

In the massless limit these agree with the literature [1]. For the massive field $\langle 0_B | T_{\mu\nu} | 0_B \rangle$ is non-zero and has nonvanishing trace for both conformal and minimal coupling. The mass breaks the conformal invariance of the field. $\langle 0_B | T_{\mu\nu} | 0_B \rangle$ vanishes exponentially as the mass and as x go to infinity, in contrast to the massless case which behaves as $O\left(x^{-4}\right)$ for the minimally coupled field and is identically 0 for conformal coupling. It is interesting to note that for conformal coupling the leading order correction for small mass is $O\left(m^2\right)$, while for minimal coupling it is $O\left(m^4\right)$. This is a qualitative difference from both M_0 and M_-.

4.3 Massless spinor field

In this section we present the expectation value of the the energy-momentum tensor for free left and right handed 2-component massless Dirac fields in the Minkowski-like vacua $|0_0\rangle$ and $|0_-\rangle$ on M_0 and M_- respectively. The corresponding values for the massless scalar field were given in [22] (and can be obtained from the results in section 4.2 in the massless limit). We find that the values depend on the spin structure but are independent of the handedness of the spinor. We begin by describing the reduction of the 4-component massive spinor formalism of chapter 2 to 2-component massless spinors.

Given the formalism set out in chapter 2 we set $m = 0$ and introduce a Weyl representation for the γ matrices

$$\gamma^0 = \begin{pmatrix} 0 & -I \\ -I & 0 \end{pmatrix} \ , \quad \gamma^i = \begin{pmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{pmatrix} \ , \tag{4.18}$$

where the σ_i are the usual Pauli matrices. With this representation the 4-component spinor ψ decouples into two 2-component spinors ψ_L and ψ_R labelled as left and right handed respectively. $\psi = \begin{pmatrix} \psi_R \\ \psi_L \end{pmatrix}$. The 4-component Lagrangian then reduces to

$$\mathcal{L}_L = \det V \frac{1}{2} i V_a^\mu \left[\psi_L^\dagger \sigma_L^a \nabla_\mu \psi_L - \left(\nabla_\mu \psi_L^\dagger \right) \sigma_L^a \psi_L \right] \ , \tag{4.19}$$

and

$$\mathcal{L}_R = \det V \frac{1}{2} i V_a^\mu \left[\psi_R^\dagger \sigma_R^a \nabla_\mu \psi_R - \left(\nabla_\mu \psi_R^\dagger \right) \sigma_R^a \psi_R \right] \ , \tag{4.20}$$

where

$$\begin{aligned} \sigma_L^a &= (I, -\sigma_x, -\sigma_y, -\sigma_z) \ , \\ \sigma_R^a &= (I, \sigma_x, \sigma_y, \sigma_z) \ . \end{aligned} \tag{4.21}$$

Further the energy-momentum tensor for the 4-component field [1],

$$T_{\mu\nu} = \frac{i}{2} \left[\bar{\psi} \gamma_{(\mu} \nabla_{\nu)} \psi - (\nabla_{(\mu} \bar{\psi}) \gamma_{\nu)} \psi \right] \ , \tag{4.22}$$

where $A_{(\mu}B_{\nu)} = 1/2(A_\mu B_\nu + A_\nu B_\mu)$, becomes the sum of left and right handed parts,

$$\begin{aligned} T^L_{\mu\nu} &= \frac{i}{2}\left[\psi^\dagger_L \sigma^L_{(\mu}\nabla_{\nu)}\psi_L - \left(\nabla_{(\mu}\psi^\dagger_L\right)\sigma^L_{\nu)}\psi_L\right] \ , && (4.23) \\ T^R_{\mu\nu} &= \frac{i}{2}\left[\psi^\dagger_R \sigma^R_{(\mu}\nabla_{\nu)}\psi_R - \left(\nabla_{(\mu}\psi^\dagger_R\right)\sigma^R_{\nu)}\psi_R\right] \ , && (4.24) \end{aligned}$$

where $\sigma^L_\mu = V^\alpha_\mu \sigma^L_\alpha$ and $\sigma^L_\alpha = \eta_{\alpha\beta}\sigma^\beta_L$. Writing (4.23) and (4.24) in the form

$$T_{\mu\nu} = \frac{i}{4}\mathrm{Tr}\left(\sigma_{(\mu}\left[\nabla_{\nu)}\psi, \psi^\dagger\right] - \sigma_{(\mu}\left[\psi, \nabla_{\nu)}\psi^\dagger\right]\right) \ , \tag{4.25}$$

where the L, R superscripts and subscripts have been dropped, and further point splitting [49] the quadratic terms in (4.25), the renormalized expectation value of $T_{\mu\nu}$, on the flat spacetimes of interest here, in the vacuum state $|0\rangle$ may now be expressed in terms of the spinor Hadamard function $S^{(1)}_{\alpha\beta\tilde{M}}(x, x') = \langle 0| \left[\psi_\alpha(x), \psi^\dagger{}_\beta(x')\right]|0\rangle$ on $\tilde{M}$ which in Minkowski space is a solution to $i\sigma^\alpha\partial_\alpha S^{(1)}(x, x') = 0$,

$$\langle 0|T_{\mu\nu}|0\rangle = \frac{i}{4}\lim_{x'\to x}\mathrm{Tr}\,\sigma_{(\mu}\left(\nabla_{\nu)} - \nabla_{\nu')}\right)\left(S^{(1)}_{\tilde{M}}(x, x') - S^{(1)}(x, x')\right) \ . \tag{4.26}$$

We may further express the spinor Hadamard function in terms of the scalar Hadamard function

$$S^{(1)}_L(x, x') = -iL^\alpha\nabla_\alpha G^{(1)}(x, x') \ , \tag{4.27}$$

$$S^{(1)}_R(x, x') = -iR^\alpha\nabla_\alpha G^{(1)}(x, x') \ , \tag{4.28}$$

where $L^\alpha = (\mathbb{I}, \sigma_x, \sigma_y, \sigma_z)$ and $R^\alpha = (\mathbb{I}, -\sigma_x, -\sigma_y, -\sigma_z)$, and so

$$\begin{aligned} \langle 0|T^L_{\mu\nu}|0\rangle &= \frac{1}{8}\lim_{x'\to x} \mathrm{Tr}\left[\left(\sigma^L_\mu \nabla_\nu + \sigma^L_\nu \nabla_\mu\right) - \left(\sigma^L_\mu \nabla_{\nu'} + \sigma^L_\nu \nabla_{\mu'}\right)\right] \\ &\quad \times L^\rho \nabla_\rho \left(G^{(1)}_{\tilde{M}}(x,x') - G^{(1)}(x,x')\right) \ , \end{aligned} \tag{4.29}$$

$$\begin{aligned} \langle 0|T^R_{\mu\nu}|0\rangle &= \frac{1}{8}\lim_{x'\to x} \mathrm{Tr}\left[\left(\sigma^R_\mu \nabla_\nu + \sigma^R_\nu \nabla_\mu\right) - \left(\sigma^R_\mu \nabla_{\nu'} + \sigma^R_\nu \nabla_{\mu'}\right)\right] \\ &\quad \times R^\rho \nabla_\rho \left(G^{(1)}_{\tilde{M}}(x,x') - G^{(1)}(x,x')\right) \ . \end{aligned} \tag{4.30}$$

$G^{(1)}(x,x')$ may be obtained as the massless limit of (4.9),

$$G^{(1)}(x,x') = -\frac{1}{[2\pi^2\left((t-t')^2 - (x-x')^2 - (y-y')^2 - (z-z')^2\right)]} \ , \tag{4.31}$$

which satisfies $\partial^\alpha \partial_\alpha G^{(1)}(x,x') = 0$. We find therefore that the 2-component spinor Hadamard function in Minkowski space is

$$\begin{aligned} S^{(1)}(x,x') &= -\frac{i}{\pi^2[(t-t')^2 - (r-r')^2]^2} \\ &\quad \times \left[\begin{array}{cc} (t-t') - \lambda(z-z') & \lambda(-(x-x') + i(y-y')) \\ \lambda(-(x-x') - i(y-y')) & (t-t') + \lambda(z-z') \end{array}\right] \ , \end{aligned} \tag{4.32}$$

where we have introduced the parameter $\lambda = 1(-1)$ for left-handed (right-handed) spinors respectively. The Hadamard functions on M_0 and M_- are given by the method of images, but care must be taken with the local frames with respect to which the spinors are expressed. We shall work throughout in a vierbein aligned along the usual Minkowski coordinate axes. This has the advantage of making covariant and partial derivatives coincide. As this vierbein is invariant under J_0 the calculation on M_0 is reduced to a straightforward calculation of derivatives of the Hadamard function. The Minkowski vierbein however is not invariant under J_- and more care must be taken on M_-.

4.3.1 M_0

On M_0, the vierbein aligned along Minkowski coordinate axes is invariant under J_0 and the Hadamard function is simply given by the method of images. We therefore have

$$S^{(1)}_{M_0}(x,x') = \sum_{n\in\mathbb{Z}} \eta^n S^{(1)}\left(x, J_0^n(x')\right) \ , \tag{4.33}$$

where again $\eta = 1, (-1)$ for spinors with periodic (antiperiodic) boundary conditions with respect to this vierbein. In this expression $G^{(1)}(x,x')$ is the Minkowski Hadamard function (4.31), and (x,x') on the left of the equality labels points in M_0 while on the right it labels points in Minkowski space. To find $\langle 0_0|T_{\mu\nu}|0_0\rangle$ we apply (4.29) and (4.30) to (4.33). The result is

$$\langle 0_0|T^L_{tt}|0_0\rangle = -\langle 0_0|T^L_{xx}|0_0\rangle = -\langle 0_0|T^L_{yy}|0_0\rangle = \frac{1}{3}\langle 0_0|T^L_{zz}|0_0\rangle = \frac{1}{8\pi^2}\sum_{n=1}^{\infty}\frac{\eta^n}{n^4a^4} \ , \tag{4.34}$$

with all other components vanishing. The values for $\langle 0_0|T^R_{\mu\nu}|0_0\rangle$ are the same. The remaining summation may be calculated explicitly using the result [40]

$$\sum_{l'=1}^{\infty}\frac{\eta^{l'}}{l'^4} = \left(-\frac{7}{8}\right)^{\frac{1-\eta}{2}}\frac{\pi^4}{90} \ , \tag{4.35}$$

and so

$$\langle 0_0|T^L_{tt}|0_0\rangle = -\langle 0_0|T^L_{xx}|0_0\rangle = -\langle 0_0|T^L_{yy}|0_0\rangle = \frac{1}{3}\langle 0_0|T^L_{zz}|0_0\rangle = \frac{\pi^2}{720a^4}\left(-\frac{7}{8}\right)^{\frac{1-\eta}{2}} \ . \tag{4.36}$$

Expectation values are non-zero in contrast to Minkowski space, and are -2 times those for the massless scalar field (see e.g [22] or the massless limits of the results in section 4.2.1). The stress tensor is diagonal and invariant under all isometries and it is independent of the handedness of the spinor but dependent on the spin structure. The $\eta = -1$ spin structure is energetically preferred.

4.3.2 M_-

On M_-, the standard Minkowski vierbein is not invariant under J_-. Care must therefore be taken in applying the method of images to expressions written with respect to this vierbein. In an invariant vierbein the spinor Hadamard function on M_- would be given directly by the method of images, that is

$$S^{(1)}_{M_-}(x, x') = \sum_{n \in \mathbb{Z}} \rho^n S^{(1)}\left(x, J_-^n x'\right) \ , \tag{4.37}$$

with $\rho = +1, (-1)$, however as we choose to work in the Minkowski vierbein the image expression is different. Consider vierbein (2.8) which is invariant under J_-. (2.8) rotates by π in the (x, y)-plane as $z \to z + a$. The transformation from this vierbein to the standard Minkowski one is clearly the corresponding rotation by $-\pi$. The associated transformation on the 2-component spinor Hadamard function is

$$S^{(1)}_{SM}(x, x') = e^{-\frac{\pi\sigma_x\sigma_y}{2a}z} S^{(1)}_{RM}(x, x') e^{\frac{\pi\sigma_x\sigma_y}{2a}z'} \ , \tag{4.38}$$

where the $R(S)$ subscript denotes the rotating (standard) vierbein respectively. Therefore on M_- in the standard vierbein

$$S^{(1)}_{SM_-}(x, x') = \sum_{n \in \mathbb{Z}} \rho^n S^{(1)}_{SM_-}\left(x, J_-^n x'\right) e^{-\frac{n\pi\sigma_x\sigma_y}{2}} \ . \tag{4.39}$$

In terms of the scalar Hadamard function this translates to using the following expression in (4.29) and (4.30)

$$G^{(1)}_{M_-}(x, x') = \sum_{n \in \mathbb{Z}} \rho^n G^{(1)}\left(x, J_-^n x'\right) e^{-\frac{n\pi\sigma_x\sigma_y}{2}} \ . \tag{4.40}$$

Here $\rho = +1, (-1)$ labels periodic (antiperiodic) spinors with respect to the vierbein which rotates by π as $z \to z + a$. ρ thus labels the two inequivalent spin structures on M_- [23]. Applying (4.29) and (4.30) to (4.40), the result is $\langle 0_- | T^L_{\mu\nu} | 0_- \rangle = \langle 0_0 | T^L_{\mu\nu} | 0_0 \rangle_{(\eta=-1)} +$

$\rho\langle 0_-|T^L_{\mu\nu}|0_-\rangle_{\mathrm{odd}}$, where

$$\begin{aligned}\langle 0_-|T^L_{xz}|0_-\rangle_{\mathrm{odd}} &= -\frac{4}{\pi^2}\sum_{n\in\mathbb{Z}}\frac{(-1)^n a(2n+1)y}{\{(2x)^2+(2y)^2+(2n+1)^2a^2\}^3}\ ,\\ \langle 0_-|T^L_{yz}|0_-\rangle_{\mathrm{odd}} &= \frac{4}{\pi^2}\sum_{n\in\mathbb{Z}}\frac{(-1)^n a(2n+1)x}{\{(2x)^2+(2y)^2+(2n+1)^2a^2\}^3}\ ,\end{aligned} \tag{4.41}$$

with all other components vanishing. The values for right handed spinors are the same.

The stress tensor is invariant under all continuous isometries of M_-, as it must be by construction, but now there are non-zero shear components, $\langle 0_-|T_{xz}|0_-\rangle$ and $\langle 0_-|T_{yz}|0_-\rangle$, which change sign under isometries that reverse the spatial orientation. These components do not depend on the handedness but they have the opposite signs in the two spin structures. The spatial orientation determined by the spin structure can thus be detected from the shear part of the stress tensor.

As on M_0, the summations may be evaluated in terms of elementary functions. The diagonal terms are given by the values on M_0 in the $\eta = -1$ spin structure there, while the non-diagonal terms may be evaluated using the calculus of residues. The results are best expressed in terms of the orthonormal frame $\{dt, dr, \omega^{\hat\phi}, dz\}$ defined by

$$\begin{aligned} x &= r\cos\phi\ ,\\ y &= r\sin\phi\ ,\end{aligned} \tag{4.42}$$

with $\omega^{\hat\phi} = r d\phi$. We find

$$\begin{aligned}\langle 0_-|T_{tt}|0_-\rangle = -\langle 0_-|T_{rr}|0_-\rangle &= -\langle 0_-|T_{\hat\phi\hat\phi}|0_-\rangle = \frac{1}{3}\langle 0_-|T_{zz}|0_-\rangle = -\left(\frac{7}{8}\right)\frac{\pi^2}{720a^4}\ ,\\ \langle 0_-|T_{\hat\phi z}|0_-\rangle &= -\frac{\rho\pi^2}{64a^4}\frac{d}{dq}\left(\frac{\sinh q}{q\cosh^2 r}\right)\ ,\end{aligned} \tag{4.43}$$

where $q = \frac{\pi}{a}r$. For $q > 0$ we find $\frac{d}{dq}\left(\frac{\sinh q}{q\cosh^2 r}\right) < 0$ and so $\langle 0_-|T_{\hat\phi z}|0_-\rangle$ picks up the sign of ρ. The shear part in either spin structure vanishes at $r = 0$ and tends to 0 exponentially as $r \to \infty$. $\langle 0_-|T_{\mu\nu}|0_-\rangle$ thus reduces to $\langle 0_0|T_{\mu\nu}|0_0\rangle$ in the $\eta = -1$ spin structure at large r.

	M_0, $\eta = 1$	M_0, $\eta = -1$	M_-
$\langle T_{00} \rangle$	$\dfrac{\pi^2}{720a^4}$	$-\dfrac{7\pi^2}{5760a^4}$	$-\dfrac{7\pi^2}{5760a^4}$
$\langle T_{rr} \rangle = \langle T_{\hat{\phi}\hat{\phi}} \rangle$	$-\dfrac{\pi^2}{720a^4}$	$\dfrac{7\pi^2}{5760a^4}$	$\dfrac{7\pi^2}{5760a^4}$
$\langle T_{zz} \rangle$	$\dfrac{\pi^2}{240a^4}$	$-\dfrac{7\pi^2}{1920a^4}$	$-\dfrac{7\pi^2}{1920a^4}$
$\langle T_{\hat{\phi}z} \rangle$	0	0	$-\dfrac{\rho\pi^2}{64a^4}\dfrac{d}{dq}\left(\dfrac{\sinh q}{q\cosh^2 q}\right)$

Table 4.1: The nonvanishing components of $\langle T_{\mu\nu} \rangle$ for a left-handed two-component spinor in the Minkowski-like vacua on M_0 and M_- in the orthonormal frame $\{dt, dr, \omega^{\hat{\phi}}, dz\}$. The values for a right-handed spinor are identical.

Numerical evidence however suggests that there is a range for r where the shear part is in fact the dominant part of the stress-energy.

We summarise the results in table 4.1. Note the difference in the results between M_0 and M_-. Thus we see that quantum fields can probe the topology. Any observer who is sensitive to the energy-momentum tensor can tell the difference between the two vacuum states.

4.4 Massive spinor field

In this section we repeat the calculations above for a massive Dirac field. The result on M_0 was given in a recent paper [48] where it is shown that the magnitude of energy density decreases with increasing field mass, as for the scalar field. We present the results for M_0 and M_-.

The energy-momentum tensor for the massive Dirac field is given in (4.22). By a calculation analogous to that leading from (4.23) to (4.26) the renormalized expectation value of the energy-momentum tensor in the state $|0\rangle$ may be written

$$\langle 0|T_{\mu\nu}|0\rangle = \frac{i}{4} \lim_{x' \to x} \operatorname{Tr} \gamma_{(\mu} \left(\nabla_{\nu)} - \nabla_{\nu')}\right) \left(S^{(1)}_{\tilde{M}}(x, x') - S^{(1)}(x, x')\right) , \qquad (4.44)$$

where $S^{(1)}_{\alpha\beta}(x,x') = \langle 0| \left[\psi_\alpha(x), \bar{\psi}_\beta(x')\right] |0\rangle$ is the massive Dirac field Hadamard function. $S^{(1)}(x,x')$ satisfies the homogeneous Dirac equation and may be expressed in terms of the scalar Hadamard function by $S^{(1)}(x,x') = -(i\gamma^\rho\nabla_\rho + m)G^{(1)}(x,x')$. Hence

$$\begin{aligned}\langle 0|T_{\mu\nu}|0\rangle &= \frac{1}{8}\lim_{x'\to x}\mathrm{Tr}[(\gamma_\mu\nabla_\nu + \gamma_\nu\nabla_\mu) - (\gamma_\mu\nabla_{\nu'} + \gamma_\nu\nabla_{\mu'})] \\ &\quad \times\gamma^\rho\nabla_\rho\left(G^{(1)}_{\tilde{M}}(x,x') - G^{(1)}(x,x')\right) \ . \end{aligned} \tag{4.45}$$

Again the Hadamard function on M_0 and M_- may be found by the method of images, with the same care taken with the local Lorentz frames as discussed for the massless field in the previous section. Again we shall work throughout with a vierbein aligned along the usual Minkowski coordinate axes.

4.4.1 M_0

Recall that on M_0

$$G^{(1)}_{M_0}(x,x') = \sum_{n\in\mathbb{Z}} \eta^n G^{(1)}\left(x, J_0^n x'\right) \ . \tag{4.46}$$

Here again we note that in the coincidence limit $G^{(1)}_{M_0}(x,x')$ becomes a function only of $(2na)^2$, which is positive, and so only the K Bessel term in (4.9) contributes. Note also that $\lim_{x'\to x}\partial_{x^\mu}G^{(1)}_{M_0}(x,x') = -\lim_{x'\to x}\partial_{x'^\mu}G^{(1)}_{M_0}(x,x')$. With these observations (4.45) reduces to

$$\langle 0_0|T_{\mu\nu}|0_0\rangle = \lim_{x'\to x}\nabla_\mu\nabla_\nu\left(G^{(1)}_{M_0}(x,x') - G^{(1)}(x,x')\right) \ , \tag{4.47}$$

and we find for the non-zero expectation values

$$\begin{aligned}\langle 0_0|T_{tt}|0_0\rangle = -\langle 0_0|T_{xx}|0_0\rangle = -\langle 0_0|T_{yy}|0_0\rangle &= \sum_{n=1}^{\infty}\eta^n\frac{2m^2}{\pi^2(2na)^2}K_2(2mna) \ , \\ \langle 0_0|T_{zz}|0_0\rangle = \sum_{n=1}^{\infty}\eta^n&\left[-\frac{2m^2}{\pi^2(2na)^2}K_2(2mna) + \frac{2m^3}{\pi^2(2na)}K_3(2mna)\right] \ , \end{aligned} \tag{4.48}$$

where $\eta = +1, (-1)$ labels periodic (antiperiodic) spinors with respect to the standard Minkowski vierbein.

The results are -4 times those of the massive scalar field (4.12). The factor of -1 is due to the different statistics while the factor of 4 is due to degrees of freedom. Thus, as with the scalar field, the leading corrections for small mass are of the order $O\left(m^2\right)$, and $\langle 0_0|T_{\mu\nu}|0_0\rangle \to 0$ exponentially in the large mass and large a limits, in contrast to the massless case where $\langle 0_0|T_{\mu\nu}|0_0\rangle$ vanishes as $O\left(a^{-4}\right)$. The massless limit agrees with twice the massless 2-component expectation values (4.34) (see also [23, 36, 47]) as expected.

4.4.2 M_-

Again we choose to work here in the Minkowski vierbein which is not invariant under J_-. A similar discussion therefore follows to that in section 4.3.2 when applying the method of images to the Hadamard functions. For 4-component spinors the transformation on the Hadamard function from the rotating vierbein (2.8) to the standard Minkowski vierbein (a rotation by $-\pi$ in the (x, y)-plane) is

$$S^{(1)}_{SM}(x, x') = e^{\frac{\pi\gamma^1\gamma^2}{2a}z} S^{(1)}_{RM}(x, x') e^{-\frac{\pi\gamma^1\gamma^2}{2a}z'} \ , \tag{4.49}$$

where the R, (S) subscript denotes the rotating (standard) vierbein respectively. Therefore, on M_- in the standard vierbein we obtain

$$S^{(1)}_{SM_-}(x, x') = \sum_{n\in\mathbb{Z}} \rho^n S^{(1)}_{SM_-}\left(x, J_-^n x'\right) e^{\frac{n\pi\gamma^1\gamma^2}{2}} \ , \tag{4.50}$$

where $\rho = +1, (-1)$ labels periodic (antiperiodic) spinors with respect to the vierbein which rotates by π as $z \longrightarrow z + a$. In terms of the scalar Hadamard function this translates to using in (4.45) the expression

$$G^{(1)}_{M_-}(x, x') = \sum_{n\in\mathbb{Z}} \rho^n G^{(1)}\left(x, J_-^n x'\right) e^{\frac{n\pi\gamma^1\gamma^2}{2}} \ . \tag{4.51}$$

In the coincidence limit $G^{(1)}_{M_-}(x, x')$ becomes a function only of $(x - (-1)^n x)^2 + (y - (-1)^n y)^2 + (na)^2$, which is positive, so that again only the K Bessel term in (4.9) contributes. Further we note again the relations (4.14). We now therefore just apply (4.45) to (4.51). The calculation is made easier by splitting the sum in (4.51) into odd and even terms. The even terms lead to the expectation values on M_0 in the twisted spin structure there and $\langle 0_-|T_{\mu\nu}|0_-\rangle = \langle 0_0|T_{\mu\nu}|0_0\rangle_{(\eta=-1)} + \rho\langle 0_-|T_{\mu\nu}|0_-\rangle_{\mathrm{odd}}$ with

$$\begin{aligned} \langle 0_-|T_{xz}|0_-\rangle_{\mathrm{odd}} &= -\sum_{n\in\mathbb{Z}}(-1)^n \frac{m^3 y(2na+a)}{\pi^2\sigma_n^3} K_3(m\sigma_n) \ , \\ \langle 0_-|T_{yz}|0_-\rangle_{\mathrm{odd}} &= \sum_{n\in\mathbb{Z}}(-1)^n \frac{m^3 x(2na+a)}{\pi^2\sigma_n^3} K_3(m\sigma_n) \ , \end{aligned} \tag{4.52}$$

where $\sigma_n = \left((2x)^2 + (2y)^2 + (2na+a)^2\right)^{1/2}$.

We see that the spinor expectation values on M_- are not -4 times those of the scalar field. In particular it is interesting to note that for the scalar field the only non-zero off-diagonal term is $\langle 0_-|T_{xy}|0_-\rangle$, while for the spinor field this term is 0 but $\langle 0_-|T_{xz}|0_-\rangle$ and $\langle 0_-|T_{yz}|0_-\rangle$ are non-zero. Note also that these two terms for the spinor field change sign under a change of spin structure. In the massless limit the results agree with twice those of the massless 2-component spinor field found in section 4.3. The leading order corrections for small mass are $O\left(m^2\right)$. The non-diagonal terms, and therefore the difference from the values on M_0 (with $\eta = -1$), are vanishing at $x = y = 0$, and they tend exponentially to zero in the limits of large mass, large a and large $x^2 + y^2$.

Chapter 5

Particle detector models

5.1 Introduction

In this chapter I present a number of new results and extensions on the theory of particle detector models within quantum field theory in curved spacetimes [14, 27]. I shall begin in section 5.2 with a review of the history and current literature on such models. The quantity of literature on this subject is quite vast and as such my review will not aim to be complete, but I will aim to cover all papers relevant for the new results presented in the subsequent chapters. Our main focus here will be with a recent paper by Schlicht [25]. Schlicht presents a model which is able to handle time-dependent responses in Minkowski space, that is detectors following trajectories which are not stationary in the sense of Letaw [52]. This is true also of the model of DeWitt [27]; however, to the knowledge of the author, Schlicht is the first to point out the importance of the regularization of the correlation function in the calculations. In particular, as will be discussed in more detail below, Schlicht shows that the usual $i\epsilon$-prescription is inadequate in this context. He then introduces an alternative regularization by considering the monopole detector of [27] as the zero size limit of a detector with spatial extent, and reproduces the usual thermal response [14, 27] on a uniformly accelerated trajectory.

We have two main aims. The first is to extend the model of Schlicht in a number of directions. In particular we extend it to Minkowski space in arbitrary dimension, to spacetimes built as quotients of Minkowski space under discrete isometry groups, and to the non-linearly coupled scalar field and the massless Dirac field. We also discuss briefly the detector coupled to the massive scalar field. The second aim is to consider the response of such detectors in various specific spacetimes and along various specific trajectories of interest. In particular we consider the responses of inertial and uniformly accelerated detectors in d-dimensional Minkowski space with a single infinite plane boundary, on a particular conical spacetime [53] (and a higher dimensional generalisation) and on the quotient spaces M_0 and M_- introduced in chapter 2. We also consider static detectors on the $\mathbb{RP}^3$ geon and comoving detectors on $\mathbb{RP}^3$ de Sitter space via those in the associated global embedding Minkowski spaces (GEMS).

5.2 Background

Particle detector models in the context of quantum field theory in curved spacetime began to be discussed in the mid 1970's. At that time one of the most important questions within the theory was, and remains to be, "what is a suitable definition of a particle in a general curved spacetime". In Minkowski space quantum field theory particles are defined with great input from the invariance group, the Poincare group. In particular a particle mode solution to a field equation in Minkowski space is one which is positive frequency with respect to the usual Minkowski time coordinate t. Under Poincare transformations positive frequency solutions transform to positive frequency solutions and so the concept of particle is the same for every inertial observer, in the sense that all inertial observers agree on the number of particles present. Further the Minkowski vacuum state, defined as the state with no particles present, is invariant under the Poincare group. The problem in a general curved spacetime however becomes clear. General relativity, in particular general covariance, teaches us that there is no notion of a preferred time coordinate and so no preferred notion of positive frequency. Further the Poincare group is not a symmetry group in a general curved spacetime, and so the definition of particle in Minkowski space cannot be extended there. As a possible resolution to the problem it was suggested by DeWitt and others that a possible operational definition of a particle could be that a particle is something that a particle detector detects (although note the circularity in the definitions if we define a particle detector to be something that detects a particle).

The original detector considered by Unruh [14] is that of a particle in a box coupled to a quantum field in a curved background manifold. The detection of a particle is indicated by an excitation of the detector by the field, and it is shown that this may correspond to both absorption or emission of a field quantum. An inertial detector in Minkowski space is not excited in the Minkowski vacuum but may be de-excited by the possibility of spontaneous emission. Unruh [14] was the first to show that a uniformly accelerated detector in Minkowski space responds as if immersed in a thermal bath at temperature $T = a/(2\pi)$ (In natural units) with the associated Planckian spectrum, which is now well known as the Unruh effect. What an accelerated detector sees as an absorption of a field quantum an inertial observer regards as the emission of one by the detector. Further Unruh also showed in [14] that the response of a static detector close to a Schwarzschild black hole is closely related to the case of the accelerated detector in Minkowski space and is thermal, while a geodesic detector near the horizon will not see this thermal flux of particles.

DeWitt's model [27] [1] is a simplification of Unruh's. He considers a quantum mechanical particle with many energy levels linearly coupled to a massless scalar field via a monopole moment operator. This is the model that appears most in the literature, and the one we will use throughout this chapter. Again the response of a uniformly accelerated detector in Minkowski space, where the detector is switched on in the infinite past and off in the infinite future is seen to be thermal and Planckian.

A good early review article which discusses particle detector models is that of Sciama, Candelas and Deutsch [55]. This is quite a thorough article on the relation between quantum field theory, horizons and thermodynamics. They discuss the uniformly accelerated detector from the point of view of the balance between the thermal fluctuations of the field along the trajectory and those of the detector itself. They find that "when the detector accelerates, the correlations in the zero-point fluctuations of the field *along the detector's world-line* no longer precisely balance its own zero-point fluctuations. The detector is consequently excited." They also discuss the reason why the effect is exactly thermal, due to the Gaussian nature of the processes. Further they consider the response in the asymptotic regions ($r \to 2M, \infty$) in the Kruskal manifold, in the Boulware, Unruh and Hartle-Hawking vacuum states, and discuss the thermal response of the inertial detector in 4-dimensional de Sitter space. All for detections carried out over the entire wordline of the detector.

---

[1] Note that the first published paper where such a detector is discussed seems to be that of Candelas and Sciama [54].

Since these original papers, a large number of papers have been written on many different aspects of these models. Takagi [31, 56, 57] considers the response of a linearly coupled monopole detector for a massive scalar field in d-dimensional Minkowski space. For the uniformly accelerated detector he notes for the first time the curious phenomena that the response for a scalar field in even dimensional spacetime has a spectrum which includes a Planckian factor whereas for odd dimensions a Fermi factor replaces the Planck one. This is often called the "apparent statistics inversion" highlighting the fact that the statistics inversion is just in the power spectrum of the detector and does not point to any change of statistics for the quantum field itself. As Unruh shows [58], the Fermi factor in the case of odd dimension results from the integral over all modes the detector is sensitive to, with each mode having a Planckian population as expected for the scalar field. In [31,57]. Takagi also shows that the response of a uniformly accelerated detector is dependent on the mass of the field. He does not find an explicit expression for the response function but does prove that the response as the mass tends to infinity tends to 0 in contrast with the thermal massless result. Further Takagi [31, 59] extends the model of DeWitt to a detector coupled to the massless Dirac field via the scalar quantity $\bar{\psi}\psi$ and also to a detector coupled to the electromagnetic field. In all cases he shows that the response for the uniformly accelerated detector satisfies the KMS condition [60, 61], and is thus thermal in this sense. Takagi's paper [31] offers a good review of the Unruh effect up until 1986, along with a good discussion of the KMS condition in the context of detector responses.

In a nice paper by Letaw [52] the stationary worldlines in Minkowski space are defined by the criterion that the particle detector responses along them are time independent. That is the Wightman function along them is invariant under translations in proper time τ. Further these worldlines are shown to be the orbits of the timelike Killing vector fields. The stationary worldlines are classified into six types (only three of which, inertial, uniform linear acceleration and uniform circular motion being well known) and responses on a representative trajectory from each class are considered. In all cases except the inertial one, the response is non-zero. An interesting case is that of uniform circular motion. The response is non-zero even though, as pointed out in an earlier paper by Letaw and Pfautsch [62], the Bogolubov transformation between the two relevant coordinate systems is trivial. Further as the Minkowski vacuum and that adapted to the circular motion are shown to coincide, the response of a detector in circular motion moving through the vacuum state adapted to it is also non-zero [63]. Davies et al [63] clarify the situation by considering a detector in uniform circular motion in Minkowski space with an infinite cylindrical boundary and also in a compactified Minkowski space. It is seen that the response of the detector vanishes if the boundary is at a radius less than the speed-of-light surface (that is, no corotating points within the spacetime have a velocity greater than c). It is only in this case that the rotating vacuum is unambiguously defined. The same conclusions were found earlier in $2+1$-dimensions for a detector in a cylindrical cavity in [64]. See also [65] for a corresponding discussion in a general axisymmetric static curved spacetime and a comment on the electromagnetic and Dirac cases. The related issue of defining a Hartle-Hawking vacuum (i.e. invariant under the isometries and regular everywhere) on the Kerr spacetime has also been discussed. A theorem by Kay and Wald [16] proves the non-existence of any stationary Hadamard state (and therefore any Hartle-Hawking state) on Kerr. However Ottewill and Winstanley, for example, [66] discuss various attempts to define a thermal state with most (but not all) of the properties of the Hartle-Hawking state. In a very recent paper Duffy and Ottewill [67] have shown that a state with the defining features of the Hartle-Hawking vacuum can be constructed when the region outside the speed-of-light surface is removed by introducing a mirror.

These results show that one must be cautious in how one interprets the particle detector as detecting particles, as discussed in [62] and for example by Sriramkumar and Padmanabhan in [68]. The response of a uniformly accelerated detector in the Minkowski vacuum and the number of Rindler particles there, found by canonically quantizing the field, agree, as of course does the response of an inertial detector and the number of Minkowski particles in the Minkowski vacuum. However the response of a circular motion detector and the number of "rotating particles", found by canonically quantizing the field with respect to a rotating coordinate system do not agree, the first being non-zero while the latter being 0. An alternative viewpoint is expressed by De Lorenci, De Paola and Svaiter [69–71] who propose an alternative canonical quantization and rotating vacuum state. We shall not discuss circular motion detectors further in this chapter.

A number of early papers discussed detectors coupled to the scalar field with more complicated couplings than the linear coupling considered by DeWitt. For example Hinton [72, 73] considers detectors non-linearly coupled to the field and those coupled to derivatives of the field. He shows, as expected with different interaction Lagrangians, that the responses for these different couplings differ. He then makes an attempt to introduce a classification scheme for such detectors. More recently Sriramkumar has shown [74, 75] that, for a uniformly accelerated detector non-linearly coupled to the scalar field in a d-dimensional Minkowski space, when d is even the response includes the Planckian factor for all couplings, while for d odd there is a Fermi factor when the power of the coupled field is odd and a Planckian factor when this power is even. Detectors which couple to components of the energy momentum tensor have been considered by Padmanabahn and Singh [76]. Ford and Roman [77] further answer a challenge set by Padmanabhan and Singh by constructing a detector which is able to measure the stress tensor of a quantized scalar field by non-gravitational means. The prescription requires a detector coupled to the derivative of the field and switched on and off rapidly.

In all the above papers the detector is that of a monopole detector. That is, the detector is coupled to the field at a point along the detector's trajectory. A step closer to a more realistic model is to consider, as Unruh does in his original paper [14], a detector extended in space. Of the two models considered by Unruh, one is not fully relativistic while the other has infinite extent. In [78] the authors define the concept of rigid detectors, where rigidity is given in the detector's rest frame. Their detector model has a truly finite spatial extent. The main example they use is that of the detector in uniform circular motion in Minkowski space. The analysis is complicated here by the effects of radiation of the walls of the detector itself. Once these are properly taken into account, the discrepancies found in [52, 62] between the detector response and a Bogolubov transformation analysis are reconciled. In this chapter we shall not consider such finite extent detectors, although we will make some more comments on them in section 5.3.1. The model we consider will instead be that of a monopole detector but considered as a limit of an infinitely extended one (see [25, 26]).

In most of the previous papers (with the exception of [52, 73]), only responses taken over the entire detector worldline are considered. Another step towards a more realistic detection process is to consider detectors switched on only for a finite time interval. The first paper to consider such a response for an inertial and uniformly accelerated detector in Minkowski space appears to be that of Svaiter and Svaiter [79]. Here the switching of the detector is performed instantaneously. Even in the inertial case the detector is excited due to the switching. In the limit of large time interval the usual results are recovered. For both the inertial and uniformly accelerated detectors the probability of detection is seen to have a logarithmic divergence, but the transition rate (the proper time derivative of the detection probability) at any time after the switching is finite, although unbounded close to the switching. It is worth noting that an earlier paper by Grove [80] also considers a finite time detector. His main interest is in the detection of the negative energy flux outside a moving mirror and the complicated situation necessarily hides the conclusions mentioned here, and half way through his calculation he moves to a detector switched on at $-\infty$. In [81] the authors prove that the logarithmic divergences in the probability seen by Svaiter and Svaiter are due to the instantaneous switching. They consider instead a continuous switching window function. The probability is then seen to be finite, and the usual results are recovered when the time of detection goes to infinity. For a more extensive discussion on this and the consideration of different continuous switching functions see [82]. All these finite time calculations are done for the massless scalar field. It is worth noting in a discussion of finite time detectors the papers [83, 84] where the Unruh effect for finite time observers is considered from the point of view of the "thermal time hypothesis" (this has also recently been extended to observers in de Sitter space [85]). Here the authors are able to associate a temperature with such observers, and even to inertial observers with a finite lifetime. It would be interesting to relate their results to the responses of finite time particle detectors. The relation is not so straightforward however as the two approaches have quite different theoretical foundations. Further the temperature introduced in [83] depends on the lifetime of the observer as well as on the proper time along the trajectory. The approach is therefore not causal in the sense that the temperature experienced at a given time depends on how long the observer will live in his future. The particle detector calculations on the other hand are causal in that the response only depends on the past motion of the detector and not on what will happen in his future. We shall not pursue the relation between these approaches further here.

Our main reference in this chapter will be Schlicht [25, 26], who is concerned with the time dependent response of a monopole detector linearly coupled to the massless scalar field travelling along non-stationary trajectories (those which are not the orbits of any timelike Killing vector field) in Minkowski space. The focus of this paper is the regularization of the correlation function that appears in the expression for the detector transition rate. In particular Schlicht shows that for a detector uniformly accelerated in Minkowski space the usual $i\epsilon$ regularization of the Wightman function leads to a time dependent, and hence presumably physically inappropriate, result. He then argues for an alternative regularization by considering the monopole detector as the limit of a spatially extended detector. The usual result is reproduced in the case of the uniformly accelerated detector. Further a numerical calculation of the response of a detector following a non-stationary worldline that interpolates between inertial and uniformly accelerated motion is presented. The response interpolates between zero and the Plankian response in a way that appears physically reasonable. It is worth noting here that some papers mention Schlicht's paper [25] as a reference for finite time detectors (in particular [84, 86, 87]). Although [25] can easily be adapted to this case it is no more a paper on it, than the original ones e.g. [27, 52] because all calculations are done for detectors which are switched on at $\tau_0 = -\infty$.

The layout of the rest of this chapter is as follows. In section 5.3 we first briefly review the formalism for the monopole detector and extend Schlicht's regularization to a Minkowski space of arbitrary dimension. We reproduce the expected results in the case of inertial and uniformly accelerated detectors. In contrast to Schlicht we shall consider both detectors switched on in the infinite past and detectors that are switched on at a finite time. In the finite time case we consider an instantaneous switching of the detector, no logarithmic divergences are encountered as we shall work throughout with the transition rate rather than the response function. Further we comment on the case of a detector coupled to a massive scalar field and present the analogue of Schlicht's correlation function there. The section ends with a reformulation of Schlicht's regularization in terms of a point-like detector and a momentum cut-off. The advantage of this view is that it makes the extension to curved spacetimes, such as Friedman-Robertson-Walker cosmologies more straightforward. Further it is shown that a similar view may be taken of a regularization obtained from that of Schlicht by modifying the detector shape profile.

In section 5.4 we consider a Schlicht-type detector on some spacetimes built as quotient spaces of Minkowski space under certain discrete isometry groups. We use the formalism of automorphic field theory on Minkowski space as discussed by Banach and Dowker [35, 36]. That is, we consider quantum field theory on these spaces as quantum field theory on Minkowski space under certain restrictions imposed by the quotient group. We are in particular interested in responses on the spacetimes M_0 and M_- introduced in section 2.2. We will present an analytic expression for the response of the inertial detector on M_0 and M_-

and for the uniformly accelerated detector on M_0, in all cases with motion in the x-direction. The time dependent response a uniformly accelerated detector on M_- is also considered, giving some analytic arguments for its general behaviour as well as discussing its numerical evaluation. This response is related to that of a static detector in the single exterior region of the $\mathbb{RP}^3$ geon. We also consider 4-dimensional Minkowski space with an infinite plane boundary and inertial and uniformly accelerated detectors with motion parallel and perpendicular to the boundary. This situation has relevance to the investigations of quantum inequality bounds and the detection of negative energy [88–95]. Ford, Svaiter and Lyra [96] and more recently Davies and Ottewill [97] have considered an inertial detector travelling parallel to the boundary, with Dirichlet boundary conditions, and switched on at a finite time, with a number of different switching functions. It is shown that the presence of the boundary, presumably for reasons related to the negative vacuum energy, reduces the excitations that result from the finite time switching.[2] We find that for the motions not parallel to the boundary the picture is considerably more subtle. Finally we consider detectors on a 4-dimensional conical spacetime and on some higher dimensional generalizations. The 4-dimensional spacetime may be considered as the spacetime outside a particular idealized cosmic string, of infinite length and zero radius (see e.g. [98]). The higher dimensional generalizations have relevance to the responses of a static detector in the exterior of the $\mathbb{RP}^3$ geon and comoving detectors in $\mathbb{RP}^3$ de Sitter space via their global embedding Minkowski spaces, as will be discussed further in sections 5.7 and 5.9.

In section 5.5 we extend Schlicht's formalism to a non-linear coupling of the scalar field, first in Minkowski space (section 5.5.1) and then in the quotient spaces (section 5.5.2). We comment briefly on responses in various specific situations.

In section 5.6 we extend the formalism further to the more realistic case of a detector coupled to the massless Dirac field in Minkowski space (section 5.6.1). In 4-dimensional Minkowski space we recover the usual result for the transition rate and the power spectrum of the Dirac noise as defined by Takagi in (e.g. [31]) for a uniformly accelerated detector. That is a power spectrum, with Fermi factor, and response function characteristic of a thermal response at the Unruh temperature (1.2). The Dirac detector is futher extended to the quotient spaces in section 5.6.2 and the transition rate and power spectrum are discussed for a uniformly accelerated detector on M_0 and M_-. On M_- our main aim is to address the question as to whether a uniformly accelerated detector can distinguish the two different spin structures [23]. We will find that at least in the case of any motion in the (x, y)-plane no such distinction arises.

---

[2] This conclusion was also reached earlier by Grove [80] with a more complicated analysis.

Section 5.7 begins with an introduction to the literature on relating the Hawking effect in black hole spacetimes to the Unruh effect in global embedding Minkowski spaces (or GEMS). It then presents an embedding of the $\mathbb{RP}^3$ geon spacetime in a 7-dimensional Minkowski space with identifications, as a quotient space of the embedding of the Kruskal manifold in 7-dimensional Minkowski space discussed by Chamblin and Gibbons [99]. The response of a uniformly accelerated detector in the embedding space, relevant to that of a static detector on the $\mathbb{RP}^3$ geon, is then discussed.

Section 5.8 considers a detector coupled linearly to a conformally coupled massless scalar field in a conformally flat spacetime. A regularization of the correlation function in the conformal vacuum is introduced which is different from that obtained by applying a naive $i\epsilon$ prescription. The expression obtained is conformally related to the correlation function found in Minkowski space in section 5.3.1.

Finally section 5.9 considers detectors in de Sitter and $\mathbb{RP}^3$ de Sitter space [100, 101]. In the case of a comoving observer we show that the regularization given in section 5.8 may also arise from the consideration of a detector with spatial extent. The usual thermal result for comoving and uniformly accelerated detectors in de Sitter space is recovered. We then discuss a comoving detector, where motion is orthogonal to the distinguished foliation in $\mathbb{RP}^3$ de Sitter space [100]. We first consider the response from a direct calculation in the $\mathbb{RP}^3$ de Sitter space and then from a relevant calculation for the associated worldline in its embedding space, motivated by the GEMS scheme. The embedding space is a 5-dimensional Minkowski space with identifications, related to the usual 5-dimensional embedding of 4-dimensional de Sitter space. These last calculations should be useful in assessing the validity and determining the relation between particle detector calculations in the global embedding Minkowski spaces and those in the original spacetimes of interest. The results provide support to the view that the GEMS approach captures, to a considerable extent, properties of quantum fields.

Appendices B,D,F,G contain details of some of the integrations required in sections 5.3-5.9. Appendix C provides details of the correlation function and responses in 2-dimensional Minkowski space. Appendix E provides more details for the automorphic field detector.

5.3 Causal detector with linear coupling

5.3.1 Massless scalar field in d-dimensional Minkowski space

In this section we briefly review the model point-like particle detectors of DeWitt [27]. We discuss the use of spatial sampling functions to regularize the correlation function and extend the regularization in [25], using a Lorentzian sampling function, to d-dimensional Minkowski space M, $d \geq 2$.

DeWitt's monopole detector

The model is that of a monopole detector, moving along a prescribed classical trajectory in M and coupled to a massless scalar field ϕ. The field has Hamiltonian H_ϕ, and satisfies the massless Klein-Gordon equation. The free field operator is expanded in terms of a standard complete set of orthonormal solutions to the field equation as

$$\phi(t, \mathbf{x}) = \frac{1}{(2\pi)^{(d-1)/2}} \int \frac{d^{d-1}k}{(2\omega)^{1/2}} \left(a(\mathbf{k}) e^{-i(\omega t - \mathbf{k}\cdot\mathbf{x})} + a^\dagger(\mathbf{k}) e^{i(\omega t - \mathbf{k}\cdot\mathbf{x})} \right) \ , \tag{5.1}$$

where $(t, x_1, x_2, \ldots, x_{d-1})$ are usual Minkowski coordinates and, in the massless case, $\omega = |\mathbf{k}|$. The field is quantized by imposing, for the creation and annihilation operators, the usual commutation relations

$$\left[a(\mathbf{k}), a^\dagger(\mathbf{k})\right] = \delta^{d-1}(\mathbf{k} - \mathbf{k}') \ . \tag{5.2}$$

The Minkowski vacuum $|0\rangle$ is the state annihilated by all the annihilation operators.

The detector is a quantum mechanical system with a set of energy eigenstates $\{|0_D\rangle, |E_i\rangle\}$. It moves along a prescribed classical trajectory $t = t(\tau)$, $\mathbf{x} = \mathbf{x}(\tau)$, where τ is the detector's proper time, and it couples to the scalar field via the interaction Hamiltonian

$$H_{\text{int}} = cm(\tau)\phi(\tau) \ , \tag{5.3}$$

where c is a (small) coupling constant and $m(\tau)$ is the detector's monopole moment operator [27]. The evolution of $m(\tau)$ is given by

$$m(\tau) = e^{iH_D\tau} m(0) e^{-iH_D\tau} \ . \tag{5.4}$$

As the interaction takes place at a point along the trajectory at any given time, the detector is known as a point-like detector.

Suppose that at time τ_0 the detector and field are in the product state $|0, E_0\rangle = |0\rangle|E_0\rangle$, where $|E_0\rangle$ is a detector state with energy E_0. We want to know the probability that at a later time $\tau_1 > \tau_0$ the detector is found in state $|E_1\rangle$, regardless of the final state of the field. We are interested in both excitations, $E_1 > E_0$ and de-excitations, $E_1 < E_0$. We find the required expression in the interaction picture (see e.g. [102]) where all field operators satisfy the free field equations and the time evolution is shared between operators and states. In this picture the time evolution of the product states is governed by the interaction Hamiltonian

$$i\frac{d}{d\tau}|\psi(\tau)\rangle = cm(\tau)\phi(\tau)|\psi(\tau)\rangle \ . \tag{5.5}$$

The amplitude for the transition from state $|0, E_0\rangle$ at $\tau = \tau_0$ to $|\psi, E_1\rangle$ at $\tau = \tau_1$, by the usual interaction picture theory is

$$\langle\psi, E_1|0, E_0\rangle = \langle\psi, E_1|T \exp\left[-i\int_{\tau_0}^{\tau_1} d\tau\, H_{\text{int}}(\tau)\right]|0, E_0\rangle \ , \tag{5.6}$$

where T is the time ordering operator. To first order in perturbation theory the expression reads

$$\begin{aligned}\langle\psi, E_1|0, E_0\rangle &= ic\langle\psi, E_1|\int_{\tau_0}^{\tau_1} d\tau\, m(\tau)\phi(\tau)|0, E_0\rangle \\ &= ic\langle E_1|m(0)|E_0\rangle\int_{\tau_0}^{\tau_1} d\tau\, e^{i(E_1-E_0)\tau}\langle\psi|\phi(\tau)|0\rangle\ . \end{aligned} \tag{5.7}$$

The transition probability to all possible states of the field is given by squaring (5.7) and summing over the complete set $\{|\psi\rangle\}$ of final field states, with the result

$$\sum_{\psi}|\langle\psi, E_1|0, E_0\rangle|^2 = c^2|\langle E_1|m(0)|E_0\rangle|^2\int_{\tau_0}^{\tau_1} d\tau\int_{\tau_0}^{\tau_1} d\tau' e^{-i(E_1-E_0)(\tau-\tau')}\langle 0|\phi(\tau)\phi(\tau')|0\rangle\ . \tag{5.8}$$

This expression has two parts. The sensitivity $c^2|\langle E_1|m(0)|E_0\rangle|^2$ depends only on the internal details of the detector and is not considered hereafter. The "response function"

$$F_{\tau_0,\tau_1}(\omega) = \int_{\tau_0}^{\tau_1} d\tau\int_{\tau_0}^{\tau_1} d\tau' e^{-i\omega(\tau-\tau')}\langle 0|\phi(\tau)\phi(\tau')|0\rangle\ , \tag{5.9}$$

where $\omega = E_1 - E_0$ ($\omega > 0$ for excitations and $\omega < 0$ for de-excitations), does not depend on the internal details of the detector and so is common for all such detectors.

We now follow Schlicht [25] and change coordinates to $u = \tau$, $s = \tau - \tau'$ for $\tau' < \tau$ and $u = \tau'$, $s = \tau' - \tau$ for $\tau' > \tau$. Then

$$F_{\tau_0,\tau_1}(\omega) = 2\int_{\tau_0}^{\tau_1} du\int_{0}^{u-\tau_0} ds\, Re\left(e^{-i\omega s}\langle 0|\phi(u)\phi(u-s)|0\rangle\right)\ , \tag{5.10}$$

because $\langle 0|\phi(\tau')\phi(\tau)|0\rangle = \langle 0|\phi(\tau)\phi(\tau')|0\rangle^*$ (a consequence of ϕ being a self adjoint operator and the axioms of inner products). To obtain an expression for the "transition rate", we differentiate with respect to τ_1 and obtain

$$\dot{F}_{\tau_0,\tau}(\omega) = 2\int_{0}^{\tau-\tau_0} ds\, Re\left(e^{-i\omega s}\langle 0|\phi(\tau)\phi(\tau-s)|0\rangle\right)\ , \tag{5.11}$$

where we have written $\tau_1 = \tau$. The transition rate is clearly causal in the sense that it does not depend on the state of motion of the detector after time τ but only on times $\tau_0 < \tau' < \tau$ which label the past motion of the detector. If the correlation function $\langle 0|\phi(\tau)\phi(\tau - s)|0\rangle$ is invariant under translations in τ this expression may be simplified to

$$\dot{F}_{\tau_0,\tau}(\omega) = \int_{-\tau+\tau_0}^{\tau-\tau_0} ds\ e^{-i\omega s} \langle 0|\phi(s)\phi(0)|0\rangle\ . \tag{5.12}$$

The transition rate, which is proportional to the derivative of the probability of the transition with respect to the switch off time τ, may be written as

$$\dot{F}_{\tau_0,\tau}(\omega) = \lim_{\delta\tau \to 0} \frac{F_{\tau+\delta\tau,\tau_0} - F_{\tau,\tau_0}}{\delta\tau}\ . \tag{5.13}$$

From (5.13) it is clear that the transition rate is not related to the response of a detector turned on at τ and off at $\tau + \delta\tau$ in the limit as $\delta\tau$ goes to zero. Rather the transition rate compares detector responses in two different ensembles of detectors, one switched off at τ and the other at $\tau + \delta\tau$ in this limit. $F_{\tau_0,\tau}$, being proportional to a probability is strictly non-negative. $\dot{F}_{\tau_0,\tau}$, however, may a priori be negative, and we shall see examples where it indeed is.

The above formalism for a monopole detector is standard in the literature (see e.g [1]) and in particular is the model first introduced by DeWitt [27]. The correlation function $\langle 0|\phi(x)\phi(x')|0\rangle$ in (5.11) is the positive frequency Wightman function, which can be obtained from the expansion (5.1),

$$\langle 0|\phi(x)\phi(x')|0\rangle = \frac{1}{(2\pi)^{d-1}} \int \frac{d^{d-1}k}{2\omega} e^{-i\omega(t-t')+i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}')}\ . \tag{5.14}$$

The integrals for $d > 2$ may be performed by moving to hyperspherical coordinates (polar coordinates and spherical coordinates for $d = 3$ and $d = 4$ respectively). The $|\mathbf{k}|$ integral requires regularization due to ultraviolet divergences. The usual regularization [1], often known in the literature as the $i\epsilon$-prescription, is to introduce the exponential cut-off $e^{-\epsilon|\mathbf{k}|}$

in the high frequency Minkowski modes. The resulting expression is (see appendix B)

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{\Gamma[d/2-1]}{4\pi^{d/2}} \frac{1}{\left[i^2(t(\tau)-t(\tau')-i\epsilon)^2 + |\mathbf{x}(\tau)-\mathbf{x}(\tau')|^2\right]^{d/2-1}} \ , \tag{5.15}$$

for $d > 2$.

For $d = 2$ it is well known that the massless scalar field theory suffers from infrared divergences [103]: The formal expression for the correlation function is (see appendix C)

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = -\frac{1}{4\pi}\ln((t(\tau)-t(\tau')-i\epsilon)^2 - (x(\tau)-x(\tau'))^2) + C \ , \tag{5.16}$$

where C is an ill-defined constant. We may still consider the particle detector in 2-dimensions dropping the constant divergent term in the correlation function. Provided we include a temporal window function which vanishes at infinity and consider an asymptotic (infinite time) detection the constant terms in the correlation function do not contribute to the response (see appendix C), this point was made by Takagi in [31].

The fundamental observation of reference [25] is that if we now use (5.15) in (5.11) on a uniformly accelerated worldline with acceleration $1/\alpha$ and proper time τ,

$$\begin{aligned} t &= \alpha \sinh(\tau/\alpha) \ , \\ x &= \alpha \cosh(\tau/\alpha) \ , \end{aligned} \tag{5.17}$$

with the detector switched on in the infinite past, $\tau_0 = -\infty$, we obtain a time dependent and apparently unphysical result, instead of the expected time independent thermal result (see e.g. [1, 27]). Schlicht shows this in 4 dimensions by specific numerical and analytic calculations, and the general d case follows similarly. His solution is to propose a different regularization of the correlation function, by considering the monopole detector as the limit of a spatially extended one. We next discuss Schlicht's regularization and related alternatives.

Alternative regularizations

In [25] Schlicht introduces a regularization of the Wightman function by considering the monopole detector as the limit of a detector extended in space[3]. An argument for the physical reasonableness of the regularization is that a spatially extended model should provide a more realistic model of a real particle detector. A related attempt to introduce a regularization via a spatially extended detector was made and discussed in some depth by Takagi [31]. Takagi however employs formal limiting arguments in a way that leads to the usual $i\epsilon$ prescription. The justification of these arguments has been criticised in [25].

Intuitively the most physically reasonable model is one which remains rigid in the detector's proper reference frame. In Minkowski space a reasonable choice for this reference frame is that built from Fermi coordinates (see e.g. [104] for a thorough discussion on the construction of Fermi coordinates). This is the choice adopted by Schlicht. Having chosen to work in a Fermi coordinate frame there are many ways in which we could model the extended detector. For one such model of a detector which cannot resolve distances shorter than ϵ we can replace the field value at a point, $\phi(\tau)$ in the interaction Hamiltonian (5.3), by a field average over a patch of the spatial hypersurface, of characteristic size ϵ and centered on the worldline. For example we could consider

$$\phi(\tau) = \int_{\epsilon\times\ldots\times\epsilon} d^{d-1}\xi \, \phi(x(\tau,\xi)) \ , \tag{5.18}$$

where (τ, ξ) are the Fermi coodinates, the integral is over all $-\epsilon/2 < \xi_i < \epsilon/2$ for all i and $x(\tau,\xi)$ describes the transformation between the Minkowski and Fermi coordinates. This model detector has a truly finite spatial extent. The effect of the spatial averaging is to introduce a cut-off at small distances. The unregularized integral expression (5.14) for the Wightman function is recovered from $\langle 0|\phi(\tau)\phi(\tau')|0\rangle$ with $\phi(\tau)$ given by (5.18) in the limit as ϵ goes to 0.

Of course we are not limited to considering the spatial averaging (5.18). Other possibilities arise, for example, if we smear the field operator with a suitable window function of characteristic length ϵ,

---

[3] See also [78] for a discussion of detector models with spatial extent.

$$\phi(\tau) = \int d^{d-1}\xi \, W_\epsilon(\xi)\phi(x(\tau,\xi)) \ , \tag{5.19}$$

which is the procedure considered in [25]. The box averaging in (5.18) is equivalent to a window function which is the product of step functions. In effect this again introduces a cut-off at short distances of the order of ϵ in size. We may choose a window function with infinite support, such as that used in [25], if the window function decreases sufficiently rapidly at large distances. Or we may consider a detector of truly finite extent with the use of a window function with compact support. We require the window function to be normalised as

$$\int d^{d-1}\xi \, W_\epsilon(\xi) = 1 \ , \tag{5.20}$$

so that smearing a constant function will return that constant value. We also require the window function to approximate the $(d-1)$-dimensional Dirac δ function, so that in the limit as ϵ tends to 0, (5.19) formally gives the field value $\phi(x(\tau))$. The window function hence staisfies $W_\epsilon(\xi) \approx 0$ for $|\xi| >> \epsilon$ and $W_\epsilon(\xi) \propto \epsilon^{-(d-1)}$ for $|\xi| << \epsilon$.

Schlicht's correlation function in d dimensions

We now extend Schlicht's regularization of the Wightman function [25] to d-dimensional Minkowski space. Although a large number of different window functions could be considered, the one chosen in [25] seems to be the easiest for obtaining a closed expression for the Wightman function on an arbitrary trajectory. Further, as we shall see in section 5.3.3, the regularization introduced by this window function is similar to that of the usual $i\epsilon$ regularization except for the important difference that the cut-off of high frequency modes is as seen in the reference frame of the detector, whereas the usual $i\epsilon$ procedure introduces a cut-off of the high frequency modes as seen by inertial observers.

The d-dimensional analogue of the window function considered in [25] is

$$W_\epsilon(\xi) = \frac{\Gamma[d/2]}{\pi^{d/2}} \frac{\epsilon}{(\xi^2+\epsilon^2)^{d/2}} \ . \tag{5.21}$$

(5.21) is sometimes referred to as a Lorentzian window or sampling function. It approximates a $(d-1)$-dimensional Dirac δ function as it scales as

$$W_\epsilon(\xi) = \frac{1}{\epsilon^{d-1}} W_1(\xi/\epsilon) \ , \tag{5.22}$$

and

$$\int d^{d-1}\xi \, W_\epsilon(\xi) = 1 \ . \tag{5.23}$$

The advantage of this window function is that in the calculation of the correlation function $\langle 0|\phi(\tau)\phi(\tau')|0\rangle$ all integrals may be done explicitly.

Now using (5.19) and (5.21), we find

$$\begin{aligned} \langle 0|\phi(\tau)\phi(\tau')|0\rangle &= \frac{1}{(2\pi)^{d-1}} \int \frac{d^{d-1}k}{2\omega} \int d^{d-1}\xi \, W_\epsilon(\xi) e^{-i(\omega t(\tau,\xi) - \mathbf{k}\cdot\mathbf{x}(\tau,\xi))} \\ &\quad \times \int d^{d-1}\xi' \, W_\epsilon(\xi') e^{i(\omega t(\tau',\xi') - \mathbf{k}\cdot\mathbf{x}(\tau',\xi'))} \ . \end{aligned} \tag{5.24}$$

Consider then the integral over ξ in (5.24),

$$g_\epsilon(\mathbf{k};\tau) = \int d^{d-1}\xi \, W_\epsilon(\xi) e^{-i(k\cdot x(\tau,\xi))} \ , \tag{5.25}$$

where we have written $(t,\mathbf{x})$, $(\omega,\mathbf{k})$,as a d-vectors x and k respectively. The transformation to Fermi coordinates is

$$x(\tau,\xi) = x(\tau) + \xi_1 e_{\xi_1}(\tau) + \cdots + \xi_{d-1} e_{\xi_{d-1}}(\tau) \ , \tag{5.26}$$

where $\dot{x}(\tau), e_{\xi_1}(\tau), \ldots, e_{\xi_{d-1}}(\tau)$ form an orthonormal basis which is Fermi transported along the worldline. Now

$$\begin{aligned} g_\epsilon(\mathbf{k};\tau) &= \int d^{d-1}\xi \, W_\epsilon(\xi) e^{-i(k\cdot x(\tau,\xi))} \\ &= e^{-i(k\cdot x(\tau))} \int d^{d-1}\xi \, W_\epsilon(\xi) e^{-i(\xi_1 k\cdot e_{\xi_1}(\tau) + \cdots + \xi_{d-1} k\cdot e_{\xi_{d-1}}(\tau))} \ . \end{aligned} \tag{5.27}$$

Writing $\bar{\mathbf{k}} := (k \cdot e_{\xi_1}(\tau), \ldots, k \cdot e_{\xi_{d-1}}(\tau))$, we have

$$g_\epsilon(\mathbf{k};\tau) = \frac{\epsilon\Gamma(d/2)}{\pi^{d/2}} e^{-i(k\cdot x(\tau))} \int d^{d-1}\xi \, \frac{1}{(\xi^2+\epsilon^2)^{d/2}} e^{-i\bar{\mathbf{k}}\cdot\xi} \ . \tag{5.28}$$

The integration for $d > 2$ may be performed by moving to hyperspherical coordinates in ξ space (see appendix B), while that for $d = 2$ may be done directly (see appendix C). The result is

$$g_\epsilon(\mathbf{k};\tau) = e^{-i(k\cdot x(\tau))} e^{-\epsilon|\bar{\mathbf{k}}|} = e^{-i(k\cdot x(\tau))} e^{-\epsilon\left[(k\cdot e_{\xi_1}(\tau))^2+\cdots+(k\cdot e_{\xi_{d-1}}(\tau))^2\right]^{(1/2)}} \ . \tag{5.29}$$

Now we use the same arguments as Schlicht. We multiply the expression

$$k = (k\cdot\dot{x})\dot{x} - (k\cdot e_{\xi_1})e_{\xi_1} - \cdots - (k\cdot e_{\xi_{d-1}})e_{\xi_{d-1}} \tag{5.30}$$

by k, giving

$$(k\cdot\dot{x})^2 = (k\cdot e_{\xi_1})^2 + \cdots + (k\cdot e_{\xi_{d-1}})^2 = (\omega\dot{t} - \mathbf{k}\cdot\dot{\mathbf{x}})^2 \ . \tag{5.31}$$

As $k\cdot\dot{x}$ is the product of a lightlike and a timelike vector, both future-pointing, it is postive. Hence

$$g_\epsilon(\mathbf{k};\tau) = e^{-i(k\cdot x(\tau))} e^{-\epsilon k\cdot\dot{x}} \ , \tag{5.32}$$

and so

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{1}{(2\pi)^{d-1}} \int \frac{d^{d-1}k}{2\omega} e^{-i\omega(t-t'-i\epsilon(\dot{t}+\dot{t}'))+i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))} \ . \tag{5.33}$$

For $d > 2$ the integral in (5.33) may be performed again by moving to hyperspherical coordinates (as shown in appendix B). The final result is

$$\begin{gathered} \langle 0|\phi(\tau)\phi(\tau')|0\rangle = \tfrac{\Gamma[d/2-1]}{4\pi^{d/2}} \tfrac{1}{A^{d/2-1}} \ , \\ A = \left[i^2(t(\tau)-t(\tau')-i\epsilon(\dot{t}(\tau)+\dot{t}(\tau')))^2 + (\mathbf{x}(\tau)-\mathbf{x}(\tau')-i\epsilon(\dot{\mathbf{x}}(\tau)+\dot{\mathbf{x}}(\tau')))^2\right] \ . \end{gathered} \tag{5.34}$$

For $d = 2$,

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{1}{4\pi}\int \frac{dk}{|k|} e^{-i|k|(t-t'-i\epsilon(\dot{t}+\dot{t}'))+ik(x-x'-i\epsilon(\dot{x}+\dot{x}'))} \ . \tag{5.35}$$

The integral is performed as that leading to (5.16). The correlation function is thus (see appendix C)

$$\begin{aligned} \langle 0|\phi(\tau)\phi(\tau')|0\rangle &= -\frac{1}{4\pi}\ln[(t(\tau)-t(\tau')-i\epsilon(\dot{t}(\tau)+\dot{t}(\tau'))^2 \\ &\quad -(x(\tau)-x(\tau')-i\epsilon(\dot{x}(\tau)+\dot{x}(\tau'))^2] + C \ , \end{aligned} \tag{5.36}$$

where C is an ill-defined constant.

We note here that following a similar calculation to that given above the usual $i\epsilon$ regularization of the correlation function (i.e. (5.15) and (5.16)) may also be obtained by considering a spatially smeared field. The difference from the above regularization being that the smearing is done in the Minkowski reference frame, that is always with respect to inertial observers, and not in the Fermi frame. This model detector is thus not rigid in its rest frame.

Inertial detector in d dimensions

First we consider the transition rate (5.12) with the correlation function (5.34) for a detector following an inertial trajectory. On Minkowski space the existence of boost Killing vectors implies that the response of the detector will be independent of velocity v. We consider therefore the trajectory $t = \tau$, $\mathbf{x} =$ constant, where $-\infty < \tau < \infty$. If the detector is switched on in the infinite past, $\tau_0 = -\infty$, the transition rate for $d > 2$ is

$$\dot{F}_\tau(\omega) = \frac{\Gamma[d/2-1]}{4\pi^{d/2}}\int_{-\infty}^{\infty} ds\, \frac{e^{-i\omega s}}{[i^2(s-2i\epsilon)^2]^{d/2-1}} \ . \tag{5.37}$$

Note that we obtain the same expression in this case if we use instead the usual correlation function (5.15) i.e. with the $i\epsilon$ regularization. The integral may be done by residues and the result as $\epsilon \to 0$ is

$$\dot{F}_\tau(\omega) = \frac{\Gamma[d/2-1](-\omega)^{d-3}}{2\pi^{d/2-1}(d-3)!}\Theta(-\omega) \ , \tag{5.38}$$

where $\Theta(-\omega)$ is the Heaviside step function. When $d = 2$ there are added complications due to the constant terms, one of which is infinite, in the correlation function (5.36), but these terms do not contribute to the transition rate provided we include a temporal window function into the integral (see appendix C). We find

$$\dot{F}_\tau(\omega) = -\frac{1}{\omega}\Theta(-\omega) \ . \tag{5.39}$$

The results given by (5.38) and (5.39) are as expected on Minkowski space. The transition rate vanishes for $\omega > 0$ indicating that an inertial detector is not excited by the Minkowski vacuum. For $\omega < 0$ the transition rate is non-zero due to the possibility of spontaneous emission by the detector.

Following Takagi's analysis of the uniformly accelerated detector [31] we may relate the transition rate given by (5.38) and (5.39) to a local density of states. On Minkowski space or a quotient of Minkowski space with non-compact spatial sections and mode solutions

$$u_{\mathbf{k}}(t,\mathbf{x}) = \frac{e^{-i|\mathbf{k}|t}}{(2|\mathbf{k}|)^{1/2}} h_{\mathbf{k}}(\mathbf{x}) \ , \tag{5.40}$$

the local density of states is defined to be [105]

$$\rho(\omega,\mathbf{x}) = \int d^{d-1}k \, |h_{\mathbf{k}}(\mathbf{x})|^2 \delta(|\mathbf{k}| - \omega) \ . \tag{5.41}$$

Due to the normalization of the modes

$$\rho(\omega) = \int d^{d-1}x \, \rho(\omega,\mathbf{x}) = \int d^{d-1}k \, \delta(|\mathbf{k}| - \omega) \ , \tag{5.42}$$

and so if the space is homogeneous then $\rho(\omega,\mathbf{x})$ is independent of $\mathbf{x}$ and is then the density of states per unit volume. The usual Minkowski density of states is obtained by taking

$h_{\mathbf{k}}(\mathbf{x}) = \frac{e^{i\mathbf{k}\cdot\mathbf{x}}}{(2\pi)^{(d-1)/2}}$, and then

$$\rho_M(\omega, \mathbf{x}) = \int \frac{d^{d-1}k}{(2\pi)^{d-1}}\, \delta(|\mathbf{k}| - \omega) \ . \tag{5.43}$$

For $d > 2$ the integral may be done by transforming to hyperspherical coordinates. We find

$$\rho_M(\omega, \mathbf{x}) = \frac{\omega^{d-2}}{2^{d-2}\pi^{(d-1)/2}\Gamma((d-1)/2)}\Theta(\omega) \ . \tag{5.44}$$

By making use of the properties of Gamma functions it is easy to show that

$$\dot{F}_\tau(\omega) = -\frac{\pi}{\omega}\Theta(-\omega)\rho_M(|\omega|, \mathbf{x}) \ . \tag{5.45}$$

This relation also holds for $d = 2$ as shown in appendix C.

For a detector that is switched on at a finite proper time τ_0, the transition rate, for $d > 2$, is given by (5.38) plus the correction term

$$C_{\Delta\tau}(\omega) = -\frac{\Gamma[d/2-1]}{2\pi^{d/2}} \int_{\Delta\tau}^{\infty} ds\, \frac{\cos(\omega s)}{(i^2 s^2)^{d/2-1}} \ , \tag{5.46}$$

where $\Delta\tau = \tau - \tau_0$. We note that $C_{\Delta\tau}(\omega)$ is manifestly finite for all $\Delta\tau > 0$. The integration may be performed by successive integrations by parts. In the case $d = 4$ we recover the result of Svaiter and Svaiter [79],

$$C_{\Delta\tau}(\omega) = \frac{\cos(\omega\Delta\tau)}{2\pi^2\Delta\tau} + \frac{|\omega|}{2\pi^2}\mathrm{si}(|\omega|\Delta\tau) \ , \tag{5.47}$$

where si is a shifted sine integral [106],

$$\mathrm{si}(z) = \int_0^z dt\, \frac{\sin t}{t} - \frac{\pi}{2} \ . \tag{5.48}$$

There are regions of the $|\omega|\Delta\tau$ parameter space where $C_{\Delta\tau}(\omega)$ and therefore $\dot{F}_\tau(\omega)$ is negative (recall that for $\omega > 0$ the asymptotic response is 0 and so $C_{\Delta\tau}(\omega)$ is the total response there). A plot of (5.47) can be found in [79].

Finally we consider the limit of the response as $\Delta\tau \to 0$. If we take this limit directly, (5.47) shows there is a divergence. It is interesting however to see whether or not keeping $\epsilon \neq 0$ regularizes this divergence. That is, is the spatial smearing enough to regularize this ultraviolet divergence? We restrict ourselves here to 4 dimensions. The transition rate for this finite time detection is

$$\begin{aligned}
\dot{F}_\tau(\omega,\epsilon) &= -\frac{1}{4\pi^2}\int_{-\Delta\tau}^{\Delta\tau} ds\, \frac{e^{-i\omega s}}{(s-2i\epsilon)^2}\ , \\
&= -\frac{1}{4\pi^2}\left[\frac{e^{i\omega\Delta\tau}}{(-\Delta\tau-2i\epsilon)} - \frac{e^{-i\omega\Delta\tau}}{(\Delta\tau-2i\epsilon)}\right] + \frac{i\omega}{4\pi^2}\int_{-\Delta\tau}^{\Delta\tau} ds\, \frac{e^{-i\omega s}}{(s-2i\epsilon)}\ , \\
&= -\frac{1}{4\pi^2}\left[\frac{e^{i\omega\Delta\tau}}{(-\Delta\tau-2i\epsilon)} - \frac{e^{-i\omega\Delta\tau}}{(\Delta\tau-2i\epsilon)}\right] + \frac{i\omega}{4\pi^2}e^{2\epsilon\omega}\left[\mathrm{E_i}(-i\omega\Delta\tau-2\epsilon\omega)\right. \\
&\quad \left. -\mathrm{E_i}(i\omega\Delta\tau-2\epsilon\omega)\right]\ , \qquad (5.49) \\
&= -\frac{1}{4\pi^2}\left[\frac{e^{i\omega\Delta\tau}}{(-\Delta\tau-2i\epsilon)} - \frac{e^{-i\omega\Delta\tau}}{(\Delta\tau-2i\epsilon)}\right] + \frac{i\omega}{4\pi^2}e^{2\epsilon\omega}\left[\mathrm{Ci}(\omega\Delta\tau-2i\epsilon\omega)\right. \\
&\quad \left. -i\,\mathrm{si}(\omega\Delta\tau-2i\epsilon\omega) - \mathrm{Ci}(\omega\Delta\tau+2i\epsilon\omega) - i\,\mathrm{si}(\omega\Delta\tau+2i\epsilon\omega)\right]\ , \qquad (5.50)
\end{aligned}$$

where we have now indicated the explicit dependence on ϵ. The second line follows by integration by parts, the third from [50], $\mathrm{E_i}$ is the exponential integral and Ci and si are cosine and shifted sine integrals respectively [106]. Taking the $\epsilon \to 0$ limit of (5.50) we recover (5.47)+$|\omega|\Theta(-\omega)/(2\pi)$ as expected. Suppose however we keep $\epsilon \neq 0$ and take the $\Delta\tau \to 0$ limit of (5.50). We see clearly from (5.49) that in this limit the transition rate vanishes. Thus the spatial smearing is enough to regularize the transition function in this limit. Similar conclusions hold in the 2-dimensional case, though we shall not spell out the calculations here. Similar investigations (though not the exact case considered here) of the $\Delta\tau$ and ϵ limits of the response function on inertial and uniformly accelerated worldlines were made in [82].

Uniformly accelerated detector in d dimensions

Next we consider the transition rate for a spatially extended detector whose centre follows the uniformly accelerated worldline (5.17). Inserting the worldline (5.17) into (5.34) gives

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{\Gamma[d/2-1]}{i^{d-2}(4\pi)^{d/2}}\frac{1}{\left(\alpha\sinh\left(\frac{\tau-\tau'}{2\alpha}\right)-i\epsilon\cosh\left(\frac{\tau-\tau'}{2\alpha}\right)\right)^{d-2}} \ . \tag{5.51}$$

As the correlation function is invariant under translations in τ the transition rate is given by

$$\dot{F}_\tau(\omega) = \frac{\Gamma[d/2-1]}{i^{d-2}(4\pi)^{d/2}}\int_{-\Delta\tau}^{\Delta\tau} ds\, \frac{e^{-i\omega s}}{\left(\alpha\sinh\left(\frac{s}{2\alpha}\right)-i\epsilon\cosh\left(\frac{s}{2\alpha}\right)\right)^{d-2}} \ . \tag{5.52}$$

In the case of a detector switched on in the infinite past, $\tau_0 = -\infty$, the integral here may be done by residues (see Appendix D). The transition rate is independent of τ along the trajectory and is given by

$$\begin{aligned}\dot{F}_\tau(\omega) = & \qquad \frac{\pi}{2^{d-2}\pi^{(d-1)/2}\alpha^{d-3}\Gamma((d-1)/2)} \\ & \times \begin{cases} \frac{\alpha\omega}{(e^{2\pi\omega\alpha}-1)}\prod_{k=1}^{(d-4)/2}\left(\left(\frac{d-2}{2}-k\right)^2+\alpha^2\omega^2\right) & d \ \text{even} \\ \frac{1}{(e^{2\pi\omega\alpha}+1)}\prod_{k=1}^{(d-3)/2}\left(\left(\frac{d-2}{2}-k\right)^2+\alpha^2\omega^2\right) & d \ \text{odd}\end{cases} \ ,\end{aligned} \tag{5.53}$$

where for $d=3$ and $d=4$ the products $\prod_{k=1}^{(d-3)/2}\left(\left(\frac{d-2}{2}-k\right)^2+\alpha^2\omega^2\right)$ and $\prod_{k=1}^{(d-4)/2}\left(\left(\frac{d-2}{2}-k\right)^2+\alpha^2\omega^2\right)$ in (5.53) are both 1. The corresponding result for $d=2$ is (see appendix C)

$$\dot{F}_\tau(\omega) = -\frac{1}{\omega(e^{2\pi\alpha\omega}-1)} \ . \tag{5.54}$$

The transition rate (5.53) and (5.54) is as in the literature [74]. It is thermal with characteristic temperature $T = 1/(2\pi\alpha)$ in the sense that is satisfies the KMS condition

$$\dot{F}_\tau(\omega) = e^{-\omega/T}\dot{F}_\tau(-\omega) \ , \tag{5.55}$$

at that temperature (see e.g [31]). Further (5.53) contains the expected "apparent" statistics inversion as we go from odd to even dimensions.

Takagi has shown [31] that the transition rate is related to the local density of Rindler states where we consider Rindler mode solutions in the definition (5.41). The relation reads explicitly

$$\dot{F}_\tau(\omega) = \frac{\pi}{\omega}\frac{\rho_R(\omega)}{\left(e^{2\pi\alpha\omega}-1\right)} \ . \tag{5.56}$$

More details can be found in [31].

The correction term for finite time detections is

$$C_{\Delta\tau}(\omega) = -\frac{\Gamma[d/2-1]}{i^{d-2}(4\pi)^{d/2}\alpha^{d-2}}\int_{\Delta\tau}^{\infty} ds\, \frac{\cos(\omega s)}{\sinh^{d-2}\left(\frac{s}{2\alpha}\right)} \ . \tag{5.57}$$

The correction term is even in ω and so breaks the KMS condition (5.55). The transition rate for a detector accelerated for only a finite time is thus not thermal in this sense. The effect of using a smoother switching for finite time detections for inertial and uniformly accelerated motions was investigated in [81, 82].

5.3.2 Massive scalar field

In this section we compute the correlation function for the smeared field operator (5.19) for a massive scalar field.

The detector model is as in section 5.3.1. The field is expanded in modes as in (5.1) but now with $\omega = \left(\mathbf{k}^2 + m^2\right)^{1/2}$, where m is the field mass. The transition rate for the detector is given by (5.11), where $\phi(\tau)$ is the smeared field operator (5.19). The correlation function is given by the expression (5.24) and we are again led to consider $g_\epsilon(\mathbf{k};\tau)$ as in (5.25). The calculation of $g_\epsilon(\mathbf{k};\tau)$ is identical to that leading to (5.32) and so we find $g_\epsilon(\mathbf{k};\tau) = e^{-i(k\cdot x(\tau))}e^{-\epsilon k\cdot\dot{x}}$ and

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{1}{(2\pi)^{d-1}}\int \frac{d^{d-1}k}{2\omega}\, e^{-i\omega(t-t'-i\epsilon(\dot{t}+\dot{t}'))+i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))} \ , \tag{5.58}$$

now with $\omega = \left(\mathbf{k}^2 + m^2\right)^{1/2}$.

The integrals here may again be done by moving to hyperspherical coordinates in k-

space. We restrict ourselves now to the case $d = 4$ (although the arbitrary d case follows similarly). After performing the angular integrals, we find

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{1}{(2\pi)^2 R}\int_0^\infty dk\,\frac{k}{\left(k^2+m^2\right)^{1/2}}\sin(kR)e^{-i\left(k^2+m^2\right)^{1/2}(t-t'-i\epsilon(\dot{t}+\dot{t}'))}\ , \quad (5.59)$$

where $R = \sqrt{(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))^2}$. This may be written as

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{-1}{8\pi^2 R}\partial_R\int_{-\infty}^\infty dk\,\frac{1}{\left(k^2+m^2\right)^{1/2}}e^{-i\left(\left(k^2+m^2\right)^{1/2}(t-t'-i\epsilon(\dot{t}+\dot{t}'))-kR\right)}\ . \quad (5.60)$$

We now change variables by $k = m\sinh\theta$, so that $\omega = m\cosh\theta$ and

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{-1}{8\pi^2 R}\partial_R\int_{-\infty}^\infty d\theta\, e^{im(R\sinh\theta-(t-t'-i\epsilon(\dot{t}+\dot{t}'))\cosh\theta)}\ . \quad (5.61)$$

For the detector trajectory we consider only timelike worldlines, $(t-t') > (\mathbf{x}-\mathbf{x}')$. We distinguish two cases. Firstly, for $(t-t') > 0$, we make the substitution $(t-t'-i\epsilon(\dot{t}+\dot{t}')) = \sqrt{\lambda}\cosh\theta_0$, $R = \sqrt{\lambda}\sinh\theta_0$, with $\lambda = (t-t'-i\epsilon(\dot{t}+\dot{t}'))^2 - R^2$. We find

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{-1}{8\pi^2 R}\partial_R\int_{-\infty}^\infty d\theta\, e^{-im\sqrt{\lambda}\cosh(\theta_0-\theta)}\ . \quad (5.62)$$

Now we note

$$K_0(z) = \frac{1}{2}\int_{-\infty}^\infty dt\, e^{-z\cosh t}\ , \quad (5.63)$$

valid for $Re(z) > 0$ [50], where K_0 is a modified Bessel function, and we may show for a timelike worldline that $Im(\sqrt{\lambda}) < 0$. Hence

$$\begin{aligned} \langle 0|\phi(\tau)\phi(\tau')|0\rangle &= \frac{-1}{4\pi^2 R}\,\partial_R\left[K_0\left(im\sqrt{\lambda}\right)\right] \\ &= -\frac{im}{4\pi^2\sqrt{\lambda}}\,K_1\left(im\sqrt{\lambda}\right)\ , \end{aligned} \quad (5.64)$$

as $\partial_z K_0(z) = K_1(z)$. The case $(t-t') < 0$ is similar except we make the change of variables $(t-t'-i\epsilon(\dot{t}+\dot{t}')) = -\sqrt{\lambda}\cosh\theta_0$, $R = \sqrt{\lambda}\sinh\theta_0$ again we may use the integral representation of K_0 and we obtain the same result (5.64).

The massless limit is easily checked. We have near $z = 0$, $K_1(z) = 1/z$ [50] and so the correlation function (5.64) agrees with that of Schlicht in this limit. Had we used the field operator without smearing we would have obtained (5.60) with $\epsilon = 0$. The usual regularization procedure, as with the massless case, would then be to introduce a cut-off in the high frequency modes by $t \rightarrow t - i\eta$, where η is small. The result would be (5.64) but with $\lambda = (t - t' - i\eta)^2 - |\mathbf{x} - \mathbf{x}'|^2$.

5.3.3 The regularization as an ultraviolet cut-off

In this section we show that the regularization discussed in sections 5.3.1 and 5.3.2 may be viewed as the introduction of an ultraviolet cut-off into the integral expression for the correlation function.

Suppose we begin with a normal monopole detector model like that of DeWitt. The transition rate is given by (5.11), and the correlation function is

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{1}{(2\pi)^{d-1}} \int \frac{d^{d-1}k}{2\omega}\, e^{-i\omega(t(\tau)-t(\tau'))+i\mathbf{k}\cdot(\mathbf{x}(\tau)-\mathbf{x}(\tau'))} \; . \tag{5.65}$$

This expression requires regularization. The usual $i\epsilon$ procedure is to introduce an ultraviolet cut-off in the modes by including in the integrand the factor $e^{-\epsilon\omega}$. This means that high frequency modes as seen by an inertial observer are exponentially suppressed. As discussed in section 5.3.1 this leads to an unphysical time-dependent result in the case where the uniformly accelerated detector is switched on in the infinite past. Let us instead consider an alternative ultraviolet cut-off where it is the high frequency modes as seen by the detector which are suppressed. Motivated by the fact that, as Unruh puts it [14] “A particle detector will react to states which have positive frequency with respect to the detectors proper time, not with respect to any universal time.”. Here the frequency is the time component of the d-momentum. We want therefore this frequency component in the rest frame of the detector, that is the frame which is Fermi-Walker transported along the trajectory with basis vectors $\{u(\tau), \mathbf{e}_{\xi_1}(\tau), \ldots, \mathbf{e}_{\xi_{d-1}}(\tau)\}$ given the d-momentum $(\omega, k_{x_1}, \ldots, k_{x_{d-1}})$ in the usual Minkwoski frame. The two frames are of course linked by a Lorentz transformation, and a simple calculation reveals that the frequency component in the Fermi-Walker frame at time τ is $\omega'(\tau) = \omega\dot{t}(\tau) - \mathbf{k}\cdot\dot{\mathbf{x}}(\tau)$. Therefore if we introduce an ultraviolet cut-off in these frequencies at τ and τ' into (5.65), that is we add a factor of $e^{-\epsilon(\omega(\dot{t}(\tau)+\dot{t}(\tau'))-\mathbf{k}\cdot(\dot{\mathbf{x}}(\tau)+\dot{\mathbf{x}}(\tau')))}$ and compare with (5.33) we see that the resulting correlation function is that of Schlicht.

Another way to introduce this cut-off is to define operationally the frequency as seen by the moving detector by $i\frac{d\phi}{d\tau} = \omega'\phi$, where τ is the detector's proper time. A mode $\phi \propto e^{-i\omega t+i\mathbf{k}\cdot\mathbf{x}}$ with frequency ω in the standard Minkowski frame thus has frequency $\omega' = \omega\dot{t}(\tau) - \mathbf{k}\cdot\dot{\mathbf{x}}(\tau)$ as seen by the detector.

We have obtained an appropriate regularization of the correlation function without introducing a detector with infinite spatial extent, that is without smearing the field operator in the interaction Hamiltonian. We will be using this reasoning as the definition of a regularization in de Sitter space in section 5.9.

5.3.4 An alternative detector form

In this section we discuss the effect of an alternative shape for the detector. In particular we discuss a Gaussian detector profile, instead of (5.21). We show that again the regularization may be viewed as a cut-off in the high frequency modes as seen by the detector.

The model is that of the previous sections with the smeared field operator in the interaction Hamiltonian given by (5.19). However now we consider a Gaussian detector shape

$$W_\epsilon(\xi) = \frac{1}{\epsilon^{d-1}(2\pi)^{(d-1)/2}} e^{-\frac{\xi^2}{2\epsilon^2}} \ . \tag{5.66}$$

This shape function also approximates a $(d-1)$-dimensional δ function, since it scales as

$$W_\epsilon(\xi) = \frac{1}{\epsilon^{d-1}} W_1(\xi/\epsilon) \ , \tag{5.67}$$

and satisfies

$$\int d^{d-1}\xi \, W_\epsilon(\xi) = 1 \ . \tag{5.68}$$

The associated correlation function again takes the form (5.24). Following the arguments after (5.24), we arrive at

$$g_\epsilon(\mathbf{k};\tau) = \frac{1}{\epsilon^{d-1}(2\pi)^{(d-1)/2}} e^{-i(k\cdot x(\tau))} \int d^{d-1}\xi \, e^{-\frac{\xi^2}{2\epsilon^2}} e^{-i\bar{\mathbf{k}}\cdot\xi} \ . \tag{5.69}$$

The integral may be done by transforming to hyperspherical coordinates as described in Appendix B. We find

$$\begin{aligned} g_\epsilon(\mathbf{k};\tau) &= e^{-i(k\cdot x(\tau))} e^{-\frac{\epsilon^2|\bar{\mathbf{k}}|^2}{2}} \ , \\ &= e^{-i(k\cdot x(\tau))} e^{-\frac{\epsilon^2|k\cdot\dot{x}(\tau)|^2}{2}} \ , \end{aligned} \tag{5.70}$$

and the correlation function becomes

$$\begin{aligned} \langle 0|\phi(\tau)\phi(\tau')|0\rangle &= \frac{1}{(2\pi)^{d-1}} \int \frac{d^{d-1}k}{2\omega} \, e^{-i\omega(t-t')+i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}')} \\ &\quad \times e^{-\frac{\epsilon^2|\omega\dot{t}(\tau)-\mathbf{k}\cdot\dot{x}(\tau)|^2}{2}} e^{-\frac{\epsilon^2|\omega\dot{t}(\tau')-\mathbf{k}\cdot\dot{x}(\tau')|^2}{2}} \ . \end{aligned} \tag{5.71}$$

Comparing (5.71) with the discussion of section 5.3.3, we see that the regularization is again equivalent to introducing an ultraviolet cut-off in the modes as seen by the moving detector: only now the cut-off at time τ takes the form $e^{-\frac{\epsilon^2\omega'^2}{2}}$ where ω' is the frequency of the Minkowski modes as seen from the Fermi-Walker transported frame. This result is expected to be useful in any thorough investigation of the dependence (or otherwise) of the response on the shape of the detector. It is clear that for a detector with extent ϵ the response will be dependent on the shape, and my intuition is that this dependence will disappear in the $\epsilon \to 0$ limit, that is in this limit all shape functions will lead to the same response. This however is still an open question.

5.4 Linearly coupled scalar field detector on quotients of Minkowski space

In this section we adapt the detector model of section 5.3.1 to spacetimes built as quotients M/Γ of Minkowski space under certain discrete symmetry groups Γ. In particular we calculate responses on M_0, M_-, Minkowski space with an infinite plane boundary and certain conical spacetimes.

We notice immediately that there is a problem in directly applying the methods of section 5.3.1 to these spacetimes. As these quotient spaces do not have infinite spatial sections in all directions, it does not directly make sense to consider a detector with infinite spatial extent as used in (5.19). We shall argue however that we may introduce a detector similar to that of section 5.3.1 by working with automorphic fields on M [35,36]. That is, we consider fields on the quotient spaces as fields on M satisfying certain transformation properties under the action of the discrete group.

5.4.1 The automorphic field detector

Consider Minkowski space M in d dimensions, and consider the quotient space M/Γ where Γ is some discrete isometry group. $|\Gamma|$ may be infinite (as indeed is the case on M_0 and M_-) which will mean that some of the following expressions remain formal in those cases. We will find however that these formalities do not interfere as in any calculations done the infinities and the formally vanishing normalization factors will cancel to give finite results.

The automorphic field $\hat{\phi}$ is constructed from the ordinary field ϕ as the sum

$$\hat{\phi}(x) := \frac{1}{\left(\sum_{\gamma\in\Gamma} p(\gamma)^2\right)^{1/2}} \sum_{\gamma\in\Gamma} p(\gamma)\phi(\gamma^{-1}x) \ , \tag{5.72}$$

where $p(\gamma)$ is a representation of Γ in $SL(\mathbb{R}) \simeq \{1,-1\}$. The normalization in (5.72) has been chosen so that, at equal times (see appendix E)

$$\left[\hat{\phi}(x), \dot{\hat{\phi}}(x')\right] = i\delta^{(d-1)}(x - x') + \text{image terms} \ . \tag{5.73}$$

The two point function for the automorphic field is then given by the method of images as

$$\langle 0|\hat{\phi}(x)\hat{\phi}(x')|0\rangle = \sum_{\gamma\in\Gamma} p(\gamma)\langle 0|\phi(x)\phi(\gamma^{-1}x')|0\rangle \ , \tag{5.74}$$

where $\langle 0|\phi(x)\phi(x')|0\rangle$ is the usual correlation function on Minkowski space.

As a model of a particle detector on M/Γ, we introduce on M a detector linearly coupled to the automorphic field by

$$H_{\text{int}} = cm(\tau)\hat{\phi}(\tau) \ , \tag{5.75}$$

with[4]

$$\hat{\phi}(\tau) = \int d^{d-1}\xi \, W_\epsilon(\xi)\hat{\phi}(x(\tau, \xi)) \ . \tag{5.76}$$

One might ask why we have not included in (5.75) image terms under Γ (that is one term for each image of the detector). There are two obvious ways in which such terms could be included. Firstly we could consider each image term in the sum to be weighted by the representation $p(\gamma)$. In certain situations, such as on Minkowski space with an infinite plane boundary with Dirichlet boundary conditions this would however lead to the detector and its image terms cancelling each other to give a vanishing interaction Hamiltonian. Alternatively we could consider image terms without the representation weights. Then each image term would be equal to that in (5.75) and so we would obtain the same results with an overall (possibly infinite) normalisation factor, which can be absorbed in the coefficient c. We therefore work with (5.75).

From (5.75), a discussion analogous to that in section 5.3.1 leads to the transition rate

---

[4] If we considered here $\hat{\phi}(\tau) = \hat{\phi}(x(\tau))$ with the usual $i\epsilon$ regularization, again we would find as Schlicht does in Minkowski space an unphysical result for the response of the uniformly accelerated detector on these spacetimes. This is most easily seen by considering that the $\gamma = I$ term (where I is the identity element) in (5.74) is that found in Minkowski space.

(to first order in perturbation theory)

$$\dot{F}_{\tau}(\omega) = 2\int_0^{\infty} ds\, Re\left(e^{-i\omega s}\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau - s)|0\rangle\right) \ , \tag{5.77}$$

where the correlation function for the automorphic field in (5.77), $\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau-s)|0\rangle$, is given by the method of images applied to (5.34) (see appendix E). Explicitly we have

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = \sum_{\gamma\in\Gamma} p(\gamma)\langle 0|\phi(\tau)\phi(\gamma^{-1}\tau')|0\rangle \ . \tag{5.78}$$

5.4.2 Scalar detector on M_0

Recall that M_0 is a quotient of 4-dimensional Minkowski space where the quotienting group Γ is that generated by the isometry $J_0 : (t, x, y, z) \mapsto (t, x, y, z + 2a)$. The transition rate for our detector is given by (5.77) with

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = \sum_{n\in\mathbb{Z}} \eta^n \langle 0|\phi(\tau)\phi(J_0^n\tau')|0\rangle \ , \tag{5.79}$$

where $\eta = +1, (-1)$ are the representations of Γ, labelling untwisted (twisted) fields respectively.

Inertial detector on M_0

Consider first a detector following the inertial trajectory

$$\begin{aligned} t = \tau(1-v^2)^{-1/2} \ , \quad & z = \tau v(1-v^2)^{-1/2} \ , \\ x = x_0 \ , \qquad & y = y_0 \ , \end{aligned} \tag{5.80}$$

where velocity $-1 < v < 0$, $-\infty < \tau < \infty$ is the detector's proper time and x_0, y_0 are constants. Substituting the trajectory into (5.79) and then (5.77), we find that the transition rate for a detector switched on in the infinite past reads

$$\dot{F}_\tau(\omega) = -\frac{1}{4\pi^2} \sum_{n=-\infty}^{\infty} \eta^n \int_{-\infty}^{\infty} ds \, \frac{e^{-i\omega s}}{(s-2i\epsilon)^2 + 4nav(s-2i\epsilon)(1-v^2)^{-1/2} - (2na)^2} \ . \tag{5.81}$$

The integral may be done by residues. The result is

$$\dot{F}_\tau(\omega) = -\frac{(1-v^2)^{1/2}}{4\pi a} \sum_{n=-\infty}^{\infty} \eta^n \frac{\sin(2\omega n a(1-v^2)^{-1/2})}{n} e^{\frac{2\omega n a v i}{(1-v^2)^{1/2}}} \Theta(-\omega) \ . \tag{5.82}$$

As on Minkowski space the transition rate vanishes for $\omega > 0$ while it is non-zero for $\omega < 0$ although the rate of spontaneous emission is altered from the Minkowski rate due to the non-trivial topology. The transition rate depends on velocity v due to the absence of a boost Killing vector in the z direction on M_0.

Consider now the limit as $v \to 0$. By the isometries of M_0 this gives the response of an inertial detector that may have arbitrary velocity in the x or y directions. Then

$$\dot{F}_\tau(\omega) = \left(-\frac{\omega}{2\pi} + \frac{1}{2\pi a} \sum_{n=1}^{\infty} \eta^n \frac{1}{n} \sin(-2n\omega a) \right) \Theta(-\omega) \ . \tag{5.83}$$

In the case of an untwisted field, $\eta = 1$, the summation in (5.83) is recognized as the Fourier series of the 2π-periodic function that on the interval $(0, 2\pi)$ takes the form $f(-2\omega a) = \frac{1}{2}(\pi + 2\omega a)$ (see e.g. [40]). We hence find

$$\dot{F}_\tau(\omega) = \frac{\left(\left[\frac{-\omega a}{\pi}\right] + \frac{1}{2}\right)}{2a} \Theta(-\omega) \ , \tag{5.84}$$

where $[x]$ denotes the integer part of x. We plot $\pi \dot{F}_\tau / |\omega|$ against $|\omega| a/\pi$ for $\omega < 0$ in figure 5.1. For a twisted field, $\eta = -1$, we note $(-1)^n \sin(nx) = \sin(n(x+\pi))$ and find

$$\dot{F}_\tau(\omega) = \frac{\left[\frac{-\omega a}{\pi} + \frac{1}{2}\right]}{2a} \Theta(-\omega) \ . \tag{5.85}$$



Figure 5.1: Transition rate of an inertial detector on M_0 for an untwisted scalar field. We have taken $\omega < 0$ and plotted $\pi \dot{F}_\tau(\omega)/|\omega|$ against $|\omega|a/\pi$.

Uniformly accelerated detector

If we consider a detector following the worldline of uniform acceleration (5.17) we obtain, again for the detector switched on at $\tau_0 = -\infty$,

$$\dot{F}_\tau(\omega) = \frac{\omega}{2\pi(e^{2\pi\alpha\omega} - 1)} \left(1 + \sum_{n=1}^{\infty} \eta^n \frac{\sin\left[2\alpha\omega \operatorname{arc\,sinh}\left(na/\alpha\right)\right]}{na\omega\sqrt{1 + n^2a^2/\alpha^2}} \right) , \tag{5.86}$$

where the integral in the transition rate has again been done by residues. The result agrees with that obtained by Louko and Marolf in [22]. The response is independent of τ. Moreover it is thermal in the sense that it satisfies the KMS condition (5.55). There is however a break from the purely Planckian form found on Minkowski space due to the image sum.

5.4.3 Minkowski space with an infinite plane boundary

In this subsection we consider a detector on d-dimensional Minkowski space[5], with Minkowski coordinates $(t, x_1, \ldots, x_{d-1})$, with an infinite boundary at $x_1 = 0$. Detectors on this spacetime (in particular in the case of 4-dimensions) have been considered by a number of authors (see e.g. [107, 108])[6]. However there has not been any presentations (as far as the author is aware) of the time dependent response for an inertial or uniformly accelerated observer who approaches the boundary from infinity. The results in this section are interesting also as a preliminary calculation before the response of detectors on M_- is considered. Many of the features seen here will also be observed there, but in a simpler context as there is no compact direction and therefore the image sum is finite. Further we will see in section 5.9.2 that the response of a uniformly accelerated detector on 4-dimensional Minkwoski space with an infinite plane boundary is very closely related to the response of an inertial detector coupled to a conformal scalar field in $\mathbb{RP}^3$ de Sitter spacetime [100].

---

[5] Throughout this section we consider $d > 2$.

[6] The case of Dirichlet boundary conditions for $d = 4$ has been the focus of some study recently on the response of detectors to negative energy [97]. The authors of [97] considered the response of a finite time detector travelling inertially parallel to the boundary. They found that the negative energy outside the boundary (4.41) has the effect of decreasing the excitations which are present even in Minkowski space due to the switching of the detector.

Again the discussion of section 5.4.1 follows through and the transition rate is given by (5.77) where now the automorphic correlation function is

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = \sum_{n=0,1} \beta^n \langle 0|\phi(\tau)\phi(J_b^n\tau')|0\rangle \ , \tag{5.87}$$

where $J_b : (t, x_1, x_2, \ldots, x_{d-1}) \mapsto (t, -x_1, x_2, \ldots, x_{d-1})$ and $\beta = +1, (-1)$, which label Neumann and (Dirichlet) boundary conditions respectively. We note here that on 4-dimensional Minkowski space with boundary at $x = 0$ the renormalized expectation values $\langle 0|T_{\mu\nu}|0\rangle$ of the energy-momentum tensor in the vacuum state induced by the Minkowski vacuum were given in section 4.2.3. For the minimally coupled massless scalar field they reduce to

$$\langle 0|T_{tt}|0\rangle = -\langle 0|T_{yy}|0\rangle = -\langle 0|T_{zz}|0\rangle = \beta\frac{1}{16\pi^2x^4} \ , \ \ \langle 0|T_{xx}|0\rangle = 0 \ . \tag{5.88}$$

Inertial detector

Firstly we consider an inertial detector with motion parallel to the boundary. Due to the isometries of the spacetime we may consider, without loss of generality, the trajectory $t = \tau$, $x_1 = \lambda$, $x_i = 0$ for $1 < i \leq d-1$ where λ, the distance from the boundary, is constant. The transition rate contains two terms. The first one comes from the first term in the boundary space correlation function (5.87) and equals the corresponding transition rate on Minkowski space. For a detector switched on in the infinite past this is given by (5.38). The second term, comes from the image term in (5.87), and is given by

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\Gamma(d/2-1)}{4\pi^{d/2}} \int_{-\infty}^{\infty} ds \, \frac{e^{-i\omega s}}{[i^2(s-2i\epsilon)^2 + (2\lambda)^2]^{d/2-1}} \ . \tag{5.89}$$

The integral in (5.89) can be evaluated by residues. Specialising to 4 dimensions, the total transition rate including the Minkowski part is found to be

$$\dot{F}_\tau(\omega) = \left(-\frac{\omega}{2\pi} - \frac{\beta}{4\pi\lambda} \sin(2\omega\lambda) \right) \Theta(-\omega) \ . \tag{5.90}$$

In figure 5.2 we plot $\dot{F}_\tau(\omega)/|\omega|$ against $|\omega|\lambda$ for Neumann boundary conditions. On the boundary the rate is twice that in Minkowski space, while far from the boundary the rate becomes that on Minkowski space. For Dirichlet boundary conditions the transition rate vanishes on the boundary as expected. Our results agree with those in [107].



Figure 5.2: Transition rate for inertial detector moving parallel to the boundary, as a function of distance λ from the boundary. We have taken $\omega < 0$ and plotted $\dot{F}_\tau(\omega)/|\omega|$ against $|\omega|\lambda$ for Neumann boundary conditions.

As on Minkowski space the response can be related to the local density of states and the difference from the Minkowski result can be attributed to the difference in the density of states. For example in the case of 4-dimensional Minkowski space with Dirichlet boundary conditions on the boundary we have mode solutions to the Klein Gordon equation of the form $u_{\mathbf{k}}(\mathbf{x}) = \frac{e^{-i\omega t}}{(4\pi^3\omega)^{1/2}} e^{i(k_y y + k_z z)} \sin(k_x x)$. From (5.41) the density of states is thus

$$\rho_B(\omega, \mathbf{x}) = \int \frac{d^3k}{2\pi^3} \sin^2(k_x x)\delta(|\mathbf{k}| - \omega) \ . \tag{5.91}$$

The integral may be done by moving to spherical coordinates, with the result

$$\rho_B(\omega, \mathbf{x}) = \frac{\omega^2}{2\pi^2}\left(1 - \frac{\sin(2\omega x)}{2\omega x}\right)\Theta(\omega) \ . \tag{5.92}$$

Hence $\dot{F}_\tau(\omega) = -\frac{\pi}{\omega}\Theta(-\omega)\rho_B(|\omega|, \mathbf{x})|_{x=\lambda}$.

Inertial detectors with motion parallel to the boundary and switched on at a finite time are considered in [96, 97].

Next, restricting ourselves to $d = 4$ again, we consider an inertial detector approaching the boundary from infinity following the worldline

$$\begin{aligned} t &= \tau(1 - v^2)^{-1/2} \ , \quad & x &= \tau v(1 - v^2)^{-1/2} \ , \\ y &= y_0 \ , & z &= z_0 \ , \end{aligned} \tag{5.93}$$

with $-1 < v < 0$ and $-\infty < \tau < 0$. We expect the response in this case to be dependent on proper time τ, as the boundary breaks the translation invariance of Minkowski space in the x-direction.

Again the transition rate is in two parts. The Minkowski part (the $n = 0$ term) will lead again to the Minkowski space rate (5.38). For a detector switched on the the infinite past this part reads

$$\dot{F}_{M\tau}(\omega) = -\frac{\omega}{2\pi}\Theta(-\omega) \ , \tag{5.94}$$

where $\Theta(-\omega)$ is a step function, and M denotes that this is the Minkowski term. The Minkowski term of course is independent of τ. It is the image term in the correlation function which leads to a τ-dependent result. The image part of the transition rate is

$$\dot{F}_{B\tau}(\omega) = -\frac{\beta}{2\pi^2}(1 - v^2) \lim_{\epsilon \to 0} \int_0^\infty ds \, Re\left(\frac{e^{-i\omega s}}{(s - 2i\epsilon)^2 - v^2(s - 2\tau)^2}\right) \ . \tag{5.95}$$

The integral may be evaluated, using some contour arguments, in terms of sine and cosine integrals (see appendix F). We find

$$\begin{aligned} \dot{F}_\tau(\omega) &= -\frac{\beta}{2\pi^2(b+c)}(-\mathrm{Ci}(b|\omega|)\cos(b|\omega|) - \mathrm{si}(b|\omega|)\sin(b|\omega|) \\ &\quad +\mathrm{Ci}(c|\omega|)\cos(c|\omega|) + \mathrm{si}(c|\omega|)\sin(c|\omega|) + 2\pi\sin(b\omega)\Theta(-\omega)) \ , \end{aligned} \tag{5.96}$$

where Ci, si are the cosine and shifted sine integrals [106], $b = 2v\tau/(1+v)$ and $c = 2v\tau/(1-v)$.

In figure 5.3 we plot $\dot{F}_\tau(\omega)/|\omega|$ against $|\omega|\tau$ for $\omega > 0$, Neumann boundary conditions and for various values of v. We note that for $\omega > 0$ the transition rate is non-zero, in contrast to the response of an inertial detector travelling parallel to the boundary, and diverges as the boundary is reached. For $\omega < 0$ recall that the Minkowski part of the transition rate is non zero. We plot the total rate for $\omega < 0$ in figure 5.4. In both cases the response depends on the velocity, as expected since there is no boost isometry in the x-direction. Further in both cases we may show that the divergence at $x = 0$ goes as $1/\tau$ and so is weaker than that in the energy expectation values (5.88). It can also be verified that the transition rate dies off at $\tau = -\infty$ as $O(1/\tau^3)$. Further numerical evidence suggests that the divergences persist for a detector that is switched on at a finite time.



Figure 5.3: Transition rate for inertial detector approaching boundary with Neumann boundary conditions and $\omega > 0$. $\dot{F}_\tau(\omega)/|\omega|$ is plotted against $|\omega|\tau$ for $v = -1/2$ (lower curve near the axis), $v = -1/3$ and $v = -1/4$ (upper curve).



Figure 5.4: Transition rate for inertial detector approaching boundary with Neumann boundary conditions and $\omega < 0$. $\dot{F}_\tau(\omega)/|\omega|$ is plotted against $|\omega|\tau$ for $v = -1/2$ (lower curve), $v = -1/3$ and $v = -1/4$ (upper curve).

Uniformly accelerated detectors

Consider now a uniformly accelerated detector with acceleration parallel to the boundary and switched on in the infinite past. We may consider without loss of generality the worldline

$$\begin{aligned} t &= \alpha \sinh(\tau/\alpha) \ , \\ x_1 &= \lambda \ , \\ x_2 &= \alpha \cosh(\tau/\alpha) \ , \\ x_i &= 0 \ , \qquad\qquad 2 < i \leq d-1 \ . \end{aligned} \tag{5.97}$$

The response again is in two parts. The first term in (5.87) leads to the thermal transition rate an accelerated detector on Minkowski space (5.53). Restricting to four dimensions, the boundary part of the transition rate is

$$\dot{F}_{B\tau}(\omega) = -\frac{\beta}{4\pi^2}\int_{-\infty}^{\infty} ds \, \frac{e^{-i\omega s}}{\left(4\left(\alpha\sinh(\frac{s}{2\alpha}) - i\epsilon\cosh(\frac{s}{2\alpha})\right)^2 - (2\lambda)^2\right)} \ . \tag{5.98}$$

The integral can be done by residues. The result is

$$\dot{F}_{B\tau}(\omega) = \frac{\beta}{4\pi}\frac{\alpha}{\lambda(\alpha^2+\lambda^2)^{1/2}}\frac{1}{(e^{2\pi\omega\alpha}-1)}\sin(2\omega\alpha\text{arcsinh}(\lambda/\alpha)) \tag{5.99}$$

which agrees with [107]. The response is thermal in the sense that it satisfies the KMS condition at temperature $T = (2\pi\alpha)^{-1}$. In [107] the authors further show that the response is related to the local density of Rindler states here.

Now let us consider the uniformly accelerated worldline (5.17). The acceleration is now perpendicular to the boundary. We begin by considering the detector switched on in the infinite past. The Minkowski part of the correlation function again leads to the thermal response (5.53). The image term on worldline (5.17) gives

$$\langle 0|\phi(\tau)\phi(J_B\tau')|0\rangle = \frac{\beta\Gamma[d/2-1]}{4\pi^{d/2}\left((4\alpha^2+4\epsilon^2)\cosh^2\left(\frac{\tau+\tau'}{2\alpha}\right)\right)^{d/2-1}} \ , \tag{5.100}$$

where again $\beta = 1, (-1)$ labels Neumann (Dirichlet) boundary conditions. It may be argued by the dominated convergence theorem that the ϵ can be dropped when calculating the transition rate. The geometrical reason is that the worldline and its image under J_B are totally spacelike separated, and so the correlation function required in the transition rate contains no divergences in the integration region.

The image term of the transition rate is thus

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\Gamma[d/2-1]}{2\pi^{d/2}}\int_0^\infty ds\,\frac{\cos(\omega s)}{\left(2\alpha\cosh\left(\frac{2\tau-s}{2\alpha}\right)\right)^{d-2}} \ . \tag{5.101}$$

We see immediately that this part of the transition rate is even in ω and hence the boundary term breaks the KMS condition (5.55). In this sense the response is non-thermal and non-Planckian.

Consider now the four-dimensional case. When $\tau = 0$ we can do the integral in (5.101) analytically, with the result

$$\dot{F}_{B0}(\omega) = \frac{\beta\omega}{4\pi\sinh(\omega\pi\alpha)} \ . \tag{5.102}$$

For general τ we may compute the integral numerically for different values of α, τ and ω. We have done this on Maple. For $\alpha = 1$ we find the total transition rate (including the thermal part) displayed in figure 5.5. Note that at many times τ the image part dominates



Figure 5.5: Transition rate for uniformly accelerated detector approaching boundary with Neumann boundary conditions, for $\alpha = 1$, $\omega = 1$ (upper curve), $\omega = 1.5$ and $\omega = 2$ (lower curve).

the Minkowski part. Further we can prove analytically that the image part of the transition rate is given by

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\omega\cos(2\tau\omega)}{2\pi\sinh(\omega\pi\alpha)} + B_\tau(\omega) \ , \tag{5.103}$$

where the function $B_\tau(\omega)$ is bounded in absolute value by $\frac{\beta}{2\pi^2\alpha}e^{-\frac{2\tau}{\alpha}}$ (see appendix G). Therefore for large but finite τ the image part of the transition rate is found not to tend to 0 but instead is periodic in τ with period π/ω.[7] This is a property only of the transition rate of a detector which is turned on in the infinite past.

Considering now a detector switched on at finite time τ_0 (which recall is the more realistic situation). The image part of the transition rate is given by (5.101) with the upper limit of the integral replaced by $\tau - \tau_0$. In figure 5.6 we plot this image part of the transition rate only, when the switching of the detector is instantaneous for, $\tau_0 = -15$, $\alpha = 1$ and $\omega = 1$. We see that when the detector is switched on only for a finite time the transition rate is periodic for some time however falls off to the usual thermal response at late τ. In appendix G we present an analytic proof of this, i.e. that the boundary part of the response vanishes in the limit as $\tau \to \infty$ when the detector is switched on at a finite time.

Recall that in the case of instantaneous switching it was found that even for the inertial

---

[7] In appendix G we prove that for arbitrary dimension the image part of the transition rate consists of a term periodic in τ with period π/ω plus a term bounded in absolute value by (constant)$e^{-\frac{(d-2)\tau}{2\alpha}}$.



Figure 5.6: Image term of the transition rate of a uniformly accelerated detector approaching the boundary with Neumann boundary conditions for a detector switched on at $\tau_0 = -15$, for $\alpha = 1$, $\omega = 1$.

detector in Minkowski space the response function includes a logarithmic divergence [79]. The transition rate however, although altered from the infinite time case, is finite for all non-zero finite time detections. Further in [81] it was shown that the divergence in the response rate is due to the instantaneous switching: if the detector is switched on smoothly, no divergence occurs. It is interesting then to briefly investigate the effect of smooth switching on the results obtained above. We introduce therefore a smooth window function in time τ into the transition rate (5.11), that is we consider the rate

$$\dot{F}_{\tau,\tau_0}(\omega) = 2\int_0^\infty ds\, W(s,\tau-\tau_0) Re\left(e^{-i\omega s}\langle 0|\phi(\tau)\phi(\tau-s)|0\rangle\right) \ , \tag{5.104}$$

where $W(s,\tau-\tau_0)$ is a smooth window function with characteristic length $\tau-\tau_0$, ie $W(s,\tau-\tau_0) \approx 1$ for $s << \tau-\tau_0$ and $W(s,\tau-\tau_0) \approx 0$ for $s >> \tau-\tau_0$. In particular we consider exponential and Gaussian switching functions

$$W_1(s,\tau-\tau_0) = e^{-\frac{|s|}{\tau-\tau_0}} \ , \tag{5.105}$$

$$W_2(s,\tau-\tau_0) = e^{-\frac{s^2}{2(\tau-\tau_0)^2}} \ . \tag{5.106}$$

The effect of these window functions on the response of a uniformly accelerated detector in Minkowski space was investigated in [82]. Here we will only consider the effect on the image part of the transition rate on Minkowski space with boundary. Substituting the image term

(5.100) and one of the window functions (5.105), (5.106) into the transition rate (5.104), we may calculate the integral numerically. In figures 5.7 and 5.8 we plot the transition rate in 4-dimensions for a sample of the parameters and with exponential and Gaussian switching respectively. The numerical results suggest that in all cases of finite time detection the image part of the transition rate tends to 0 as the detection time tends to infinity. That is, the transition rate tends to that on Minkowski space in this limit, as expected as in this limit the detector recedes infinitely far from the boundary. It is an interesting result that this was not the case for a detector switched on in the infinite past.



Figure 5.7: Image term of the transition rate of a uniformly accelerated detector approaching boundary with Neumann boundary conditions for a detector switched on at $\tau_0 = -15$ with an exponential switching function, for $\alpha = 1$, $\omega = 1$.



Figure 5.8: Image term of the transition rate of a uniformly accelerated detector approaching boundary with Neumann boundary conditions for a detector switched on at $\tau_0 = -15$ with a Gaussian switching function, for $\alpha = 1$, $\omega = 1$.

5.4.4 Conical singularity and generalisations

In this section we consider the response of a uniformly accelerated detector following trajectory (5.17) on the quotient space of Minkowski space under the group generated by the involution $J_{c_2} : (t, x, y, z) \mapsto (t, -x, -y, z)$. Further we consider the higher dimensional generalisation of this spacetime constructed as the quotient of d-dimensional Minkowski space under the involution $J_{c_k} : (t, x_1, x_2, \dots x_{d-1}) \mapsto (t, -x_1, -x_2, - \dots - x_k, x_{k+1}, \dots, x_{d-1})$ where $1 < k < d$. For reasons discussed later the response on these higher dimensional spaces will be relevant when we consider the response of static detectors on the $\mathbb{RP}^3$ geon in section 5.7 and inertial detectors on $\mathbb{RP}^3$ de Sitter space in section 5.9.

These spacetimes are conifolds [109, 110]. As quotients of Minkowski space under an involution with fixed points they are flat away from these $(d-k)$-dimensional hypersurfaces of fixed points but may be considered to have a distributional curvature on them (see e.g [111, 112]). The spacetime M/J_{c_2} is sometimes referred to as a conical spacetime as it has a conical singularity at $x_1 = x_2 = 0$. Transforming to cylindrical coordinates by $x_1 = r\cos\phi$, $x_2 = r\sin\phi$, the isometry takes the form $J_{c_2} : (t, r, \phi, x_3, \dots x_{d-1}) \mapsto (t, r, \phi + \pi, x_3, \dots, x_{d-1})$ and the metric reads

$$ds^2 = dt^2 - dr^2 - r^2 d\phi^2 - (dx_3)^2 - \cdots - (dx_{d-1})^2 \ , \tag{5.107}$$

where $dr^2 + r^2 d\phi^2$ with the identification $(r, \phi) \sim (r, \phi + \pi)$ is the metric on a cone with deficit angle π. In 4 dimensions M/J_{c_2} may be considered as the spacetime outside an idealized, cosmic string with “gravitational mass per unit length” $\mu = 1/8$ (see [113]).

First we note that for an inertial or uniformly accelerated detector whose motion is in any direction x_i with $k < i < d$, the response on these spacetimes will be the same as that of a detector at rest or accelerating parallel to the boundary on Minkowski space with boundary (where λ in (5.90) and (5.99) is now given by the shortest distance of the detector to the hypersurface of fixed points $\lambda = \left((x_1)^2 + (x_2)^2 + \cdots + (x_k)^2\right)^{1/2}$). This can be clearly seen by directly comparing the correlation functions in both cases.

Consider now a particle detector uniformly accelerated with trajectory (5.17) in the spacetime M/J_{c_k}. Again as J_{c_k} is an involution the correlation function consists of two terms,

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = \langle 0|\phi(\tau)\phi(\tau')|0\rangle + \beta\langle 0|\phi(\tau)\phi(J_{c_k}\tau')|0\rangle \ , \tag{5.108}$$

where $\beta = +1, (-1)$ label the two possibilities for the representations of the group in the automorphic field expansion (5.72). Consider first a detector switched on at $\tau_0 = -\infty$. The first term, when substituted into the transition rate (5.77) on the worldline (5.17), leads to the thermal response in Minkowski space (5.53). The image term is

$$\beta\langle 0|\phi(\tau)\phi(J_{c_k}\tau')|0\rangle = \frac{\beta\Gamma[d/2-1]}{4\pi^{d/2}\left(4\alpha^2\cosh^2\left(\frac{2\tau-s}{2\alpha}\right)+C_k\right)^{d/2-1}} \ , \tag{5.109}$$

where $C_k = \sum_{m=2}^{k}(2x_m)^2$. As the trajectory and its image are totally spacelike separated we have dropped the regularization in the above expression. The transition rate for this part is then

$$\dot{F}_{I\tau(\omega)} = \frac{\beta\Gamma[d/2-1]}{2\pi^{d/2}}\int_0^\infty ds\, \frac{\cos(\omega s)}{\left(4\alpha^2\cosh^2\left(\frac{2\tau-s}{2\alpha}\right)+C_k\right)^{d/2-1}} \ . \tag{5.110}$$

(5.110) is even in ω and so the image term breaks the KMS condition and the response is non-thermal and non-Planckian. Further with a similar calculation to that of appendix G we may prove that (5.110) consists of a term periodic in τ with period π/ω and a term which decays exponentially as $\tau \to \infty$. The qualitative behaviour is therefore similar to that of the uniformly accelerated detector on M with boundary, investigated in section 5.4.3. As in section 5.4.3 it can be shown also here that for a detector switched on at $\tau_0 > -\infty$ the image part of the transition rate tends to 0 as $\tau \to \infty$.

For a detector that accelerates towards the surface of fixed points of the involution, that is in the direction $r = ((x_1)^2 + (x_2)^2 + \cdots + (x_k)^2)^{1/2}$, the response is identical to that on Minkowski space with boundary with acceleration perpendicular to the boundary. In sections 5.7 and 5.9 we shall plot (5.110) for some specific values of d and k numerically.

We end this section with a comment on more general cosmic string spacetimes. The methods used above could be easily applied to a larger class of idealized cosmic string spacetimes for which the metric is (5.107), with the identification $(r, \phi) \sim (r, \phi + \pi/n)$ where $n \in \mathbb{Z}$ (and thus a deficit angle of $\pi(2-1/n)$), as in these cases the correlation function may be given by a mode sum. Detectors with motion parallel to such cosmic strings have been considered by [114, 115]. Their main conclusions, which agree with ours here where they overlap, are that the detector does respond to the presence of the string in a manner which depends on its distance from the string. Our results above and in the previous subsection add to the discussion, as we have been able to show the behaviour of detectors when motion is perpendicular to the string for the specific case of $n = 1$. Numerical evaluations of the transition rate for any n could be done in a similar way, but we shall not pursue this further here. It is important to note however this class of cosmic strings does not include realistic cosmic strings of the GUT scale where the deficit angle is $\approx 10^{-5}$.

5.4.5 Scalar detector on M_-

Finally let us consider M_-. Recall that M_- is a quotient of Minkowski space (or of M_0, it being a double cover of M_-) under the map $J_- : (t, x, y, z) \mapsto (t, -x, -y, z + a)$. Again we may use the method of images to find the correlation function for the automorphic field, with the result

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = \sum_{n \in \mathbb{Z}} \langle 0|\phi(\tau)\phi(J_- \tau')|0\rangle \ . \tag{5.111}$$

The transition rate is given by (5.77).

Inertial detector

Considering a detector following the inertial trajectory (5.93). The transition rate may again be split into two parts. The first comes from the M_0 part in the image sum (the even n terms in (5.111)) and will lead to the same response as on M_0 (5.84). The other part (due to odd n terms in (5.111)) is similar to the boundary part of such a detector on M with boundary, giving in the transition rate the contribution

$$
\begin{aligned}
\dot{F}_{I\tau(\omega)} &= -\frac{1}{2\pi^2}(1-v^2)\sum_{n=-\infty}^{\infty}\frac{1}{C_n+B_n}(-\mathrm{Ci}(B_n|\omega|)\cos(B_n|\omega|) \\
&\quad -\mathrm{si}(B_n|\omega|)\sin(B_n|\omega|)+\mathrm{Ci}(C_n|\omega|)\cos(C_n|\omega|)+\mathrm{si}(C_n|\omega|)\sin(C_n|\omega|) \\
&\quad +2\pi\sin(B_n\omega)\Theta(-\omega))\ , \qquad (5.112)
\end{aligned}
$$

where

$$
B_n = \frac{-4\tau v^2+(16\tau^2v^2+4(1-v^2)^2((2y_0)^2+(2na+a)^2))^{1/2}}{2(1-v^2)}\ , \qquad (5.113)
$$

$$
C_n = \frac{+4\tau v^2+(16\tau^2v^2+4(1-v^2)^2((2y_0)^2+(2na+a)^2))^{1/2}}{2(1-v^2)}\ . \qquad (5.114)
$$

In contrast to the analogous result on M with boundary, there is no divergence here at $x=0$ as there is no obstruction there and the inertial detector on M_- carries through $x=0$ smoothly. Note that on M_- the energy-momentum tensor expectation values are finite over the whole spacetime (see e.g. [22] or the massless limits of the results found in section 4.2.2), while on M with boundary they diverge at $x=0$. For $\omega>0$ the M_0 part of the transition rate vanishes while the image part is odd in τ. Further, numerical evaluations of the sum indicate that it is non-zero. Here therefore we have an example of a spacetime and trajectory with no pathologies at all where the total transition rate is negative for some values of proper time τ. The response depends on the velocity as $t\partial_x + x\partial_t$ is not a Killing vector on M_-.

Uniformly accelerated detector

Consider a detector following the uniformly accelerated worldline (5.17). Again the correlation function is in two parts. The part coming from the M_0 part of the image sum is given by the corresponding response on M_0 ((5.86) with $\eta=-1$). This part satisfies the KMS condition and so in this sense is thermal. The "image" part is then somewhat similar to the boundary part found in the case of M with boundary. The transition rate for this term is

$$\dot{F}_{I\tau}(\omega) = \frac{1}{2\pi^2}\sum_{n=-\infty}^{\infty}\int_0^{\infty} ds\, \frac{\cos(\omega s)}{4\alpha^2\cosh^2\left(\frac{2\tau-s}{2\alpha}\right)+4y_0^2+a^2(2n-1)^2}\ , \tag{5.115}$$

where y_0 is the y coordinate of the detector. As on Minkowski space with boundary this image part of the transition rate is even in ω and so breaks the KMS condition. The transition rate is thus non-thermal and non-Planckian. With a similar calculation to that done in appendix G we find

$$\dot{F}_{I\tau}(\omega) = \sum_{n=-\infty}^{\infty}\left(\frac{\alpha\cos(2\tau\omega)\sin\left(\alpha\omega\operatorname{arccosh}\left(\frac{c_n}{2\alpha^2}\right)\right)}{\pi(c_n^2-4\alpha^4)^{1/2}\sinh(\omega\pi\alpha)}+B_{n,\tau}(\omega)\right)\ , \tag{5.116}$$

where $c_n = 2\alpha^2+4y_0^2+(2na-a)^2$, and each $B_{n,\tau}(\omega)$ is bounded by $1/(2\pi^2\alpha)e^{-2\tau/\alpha}$. Further we show in appendix G that the sum of $B_{n,\tau}(\omega)$ over n is bounded by a function which exponentially decays as $\tau\to\infty$. We see as with the detector on M with boundary for large τ the response becomes periodic in τ with period π/ω.

We may investigate the general case here numerically. An analytic result is easy to find in the case when $\tau = 0$, with the result

$$\dot{F}_{I0}(\omega) = \frac{\alpha}{2\pi}\sum_{n=-\infty}^{\infty}\frac{\sin\left(\alpha\omega\operatorname{arccosh}\left(\frac{c_n}{2\alpha^2}\right)\right)}{(c_n^2-4\alpha^4)^{1/2}\sinh(\omega\pi\alpha)}\ . \tag{5.117}$$

For a detector switched on instantaneously at a finite time $\tau_0 > -\infty$ an analytic calculation (see appendix G) shows that the difference between the response on M_- and that on M_0 dies off as $\tau\to\infty$, as would be expected far away from $x=0$. It is an interesting point that this is not the case for the detector switched on the infinite past.

This clarifies and adds to the discussion on particle detectors given in [22].

5.5 Causal detector with non-linear coupling

In this section we extend the detector model to non-linear couplings. The aim is to see whether the formalism adapts easily to this case. The non-linear case is also interesting in that it is a step towards the Dirac detector considered in section 5.6.

5.5.1 Minkowski space

The detector model is similar to that section 5.3 except in that the interaction Hamiltonian is taken to be

$$H_{\text{int}} = cm(\tau)\phi^n(\tau) \ , \tag{5.118}$$

where n is an integer greater than 1. Again we assume the field is in its ground state and detector in a state with energy E_0 at $\tau = \tau_0$, and we seek the probability that the detector is found in state $|E_1\rangle$ at later time $\tau_1 > \tau_0$. This probability is found to be, to first order in perturbation theory

$$\sum_{\psi} |\langle\psi, E_1|0, E_0\rangle|^2 = c^2|\langle E_1|m(0)|E_0\rangle|^2 \int_{\tau_0}^{\tau_1} d\tau \int_{\tau_0}^{\tau_1} d\tau' e^{-i\omega(\tau-\tau')}\langle 0|\phi^n(\tau)\phi^n(\tau')|0\rangle \ , \tag{5.119}$$

with $\omega = E_1 - E_0$. The response function is

$$F_\tau(\omega) = \int_{\tau_0}^{\tau_1} d\tau \int_{\tau_0}^{\tau_1} d\tau' e^{-i\omega(\tau-\tau')}\langle 0|\phi^n(\tau)\phi^n(\tau')|0\rangle \ , \tag{5.120}$$

and the transition rate is given by

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds\, Re\left(e^{-i\omega s}\langle 0|\phi^n(\tau)\phi^n(\tau - s)|0\rangle\right) \ , \tag{5.121}$$

where we have used the property $\langle 0|\phi^n(\tau')\phi^n(\tau)|0\rangle = \langle 0|\phi^n(\tau)\phi^n(\tau')|0\rangle^*$, which follows from ϕ being a self adjoint operator, and have taken $\tau_0 = -\infty$ and $\tau_1 = \tau$. Finite time detectors are easily considered by changing the upper limit in (5.121) to $\tau - \tau_0$.

For this non-linear coupling there are added issues not present in the linear case due to the interaction Lagrangian involving the product of the field taken at the same point. As a result the usual field theoretic divergence here must be dealt with in a suitable manner. We explain here the procedure on Minkowski space. The additional subtleties that arise for automorphic fields will be addressed in section 5.5.2 (see [108] for a discussion on this point).

Instead of the interaction Hamiltonian (5.118), we are in fact led to consider a normal ordered version, $H_{\text{int}} = cm(\tau) : \phi^n(\tau) :$. Then in (5.121) $\langle 0|\phi^n(\tau)\phi^n(\tau')|0\rangle$ is replaced by $\langle 0| : \phi^n(\tau) :: \phi^n(\tau') : |0\rangle$ and by Wick's theorem we obtain

$$\langle 0| : \phi^n(\tau) :: \phi^n(\tau') : |0\rangle = n!\langle 0|\phi(\tau)\phi(\tau')|0\rangle^n \ . \tag{5.122}$$

Alternatively using the original Hamiltonian (5.118) we are also led to (5.122) by expanding the $2n$-point function in terms of 2-point functions and replacing any divergent quantities with their renormalized expectation values, such as replacing $\langle 0|\phi^2(x)|0\rangle$ with $\langle 0| : \phi^2(x) : |0\rangle$. These renormalized expectation values vanish on Minkowski space.

Putting (5.122) into (5.121), we are led to a very similar expression for the transition rate as in the linear case. Further it is clear from a direct calculation that if we consider the uniformly accelerated worldline and the usual correlation function (5.15) in (5.122) we obtain a τ-dependent expression. Further a numerical evaluation of the transition rate for $n = 2$ shows that the transition rate is also τ-dependent even in the $\epsilon \to 0$ limit. We see no reason why this would not also be the case for general n. This suggests that as for the linear coupling we need to seek out a different regularization of the correlation function. We therefore consider a smeared field operator (5.19) in the interaction Hamiltonian (5.118). The transition rate is still given by (5.121) but now $\phi(\tau)$ is the smeared field operator, and we are led to an expression analogous to (5.122) with no added complications due to the smearing. The transition rate therefore depends on $\langle 0|\phi(\tau)\phi(\tau')|0\rangle$ and so using the smeared field operator (5.19) with the Lorentzian profile function (5.21) we are clearly led to the regularization (5.34) for this correlation function.

Consider again a uniformly accelerated detector. We have already seen that Schlicht's correlation function on this worldline is given, for $d > 2$, by (5.51). Substituting (5.51) into transition rate (5.121) we obtain

$$\dot{F}_\tau(\omega) = \left(-\frac{\Gamma[d/2-1]}{i^{d-2}(4\pi)^{d/2}}\right)^n n! \lim_{\epsilon\to 0}\int_{-\infty}^{\infty} ds\, \frac{e^{-i\omega s}}{\left(\alpha\sinh\left(\frac{s}{2\alpha}\right) - i\epsilon\cosh\left(\frac{s}{2\alpha}\right)\right)^{n(d-2)}}. \tag{5.123}$$

The integral may be performed by contour methods (see appendix B). The result is

$$\begin{aligned}\dot{F}_\tau(\omega) = \quad & \left(\frac{\Gamma[d/2-1]}{4\pi^{d/2}}\right)^n n! \frac{2\pi}{2^{d-2}\Gamma[m]\alpha^{m-1}} \\ \times & \begin{cases} \frac{\alpha\omega}{(e^{2\pi\omega\alpha}-1)} \prod_{l=1}^{m/2-1}\left(\left(\frac{m}{2}-l\right)^2+\alpha^2\omega^2\right) & m \text{ even} \\ \frac{1}{(e^{2\pi\omega\alpha}+1)} \prod_{l=1}^{(m-1)/2}\left(\left(\frac{m}{2}-l\right)^2+\alpha^2\omega^2\right) & m \text{ odd}\end{cases} \quad , \end{aligned} \tag{5.124}$$

where $m = n(d-2)$, and for $m = 4$ the product over l is set to 1. The response is independent of τ and satisfies the KMS condition (5.55) at the temperature $T = (2\pi\alpha)^{-1}$. It also contains the expected statistics inversions for different values of dimensions and couplings and agrees with the literature (e.g [74]).

5.5.2 Automorphic fields

For the non-linearly coupled detectors on quotients of Minkowski space, we consider detectors coupled to automorphic fields in Minkowski space as in section 5.4. We specialize in this section to the case $n = 2$ (though the case of higher order will follow in an analogous way).

The automorphic field is defined as previously by

$$\hat{\phi}(x) = \frac{1}{\left(\sum_{\gamma\in\Gamma} p(\gamma)^2\right)^{1/2}} \sum_{\gamma\in\Gamma} p(\gamma)\phi(\gamma^{-1}x)\ . \tag{5.125}$$

We consider a monopole detector coupled to the automorphic field via the interaction Hamiltonian

$$H_{\text{int}} = cm(\tau)\hat{\phi}^2(\tau)\ , \tag{5.126}$$

with

$$\hat{\phi}(\tau) = \int d^{d-1}\xi\, f(\xi)\hat{\phi}(x(\tau,\xi)) \ , \tag{5.127}$$

where ξ has the same meaning as in the previous sections. Following the arguments of the previous sections, the transition rate is found to be

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds\, Re\left(e^{-i\omega s}\langle 0|\hat{\phi}^2(\tau)\hat{\phi}^2(\tau - s)|0\rangle\right) \ . \tag{5.128}$$

We need to address the infinities in (5.128). We first note that (5.125), (5.127) and the usual expansion of the Minkowski field operator ϕ in terms of annihilation and creation operators leads to Wick's theorem for $\hat{\phi}$ in the form

$$\begin{aligned}\langle 0|\hat{\phi}(\tau_1)\hat{\phi}(\tau_2)\hat{\phi}(\tau_3)\hat{\phi}(\tau_4)|0\rangle \quad = \quad & \langle 0|\hat{\phi}(\tau_1)\hat{\phi}(\tau_4)|0\rangle\langle 0|\hat{\phi}(\tau_2)\hat{\phi}(\tau_3)|0\rangle \\ & +\langle 0|\hat{\phi}(\tau_1)\hat{\phi}(\tau_3)|0\rangle\langle 0|\hat{\phi}(\tau_2)\hat{\phi}(\tau_4)|0\rangle \\ & +\langle 0|\hat{\phi}(\tau_1)\hat{\phi}(\tau_2)|0\rangle\langle 0|\hat{\phi}(\tau_3)\hat{\phi}(\tau_4)|0\rangle \ . \end{aligned} \tag{5.129}$$

Setting $\tau_1 = \tau_2 = \tau$ and $\tau_3 = \tau_4 = \tau'$ yields

$$\langle 0|\hat{\phi}^2(\tau)\hat{\phi}^2(\tau')|0\rangle = 2\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle^2 + \langle 0|\hat{\phi}^2(\tau)|0\rangle\langle 0|\hat{\phi}^2(\tau')|0\rangle \ . \tag{5.130}$$

The divergent term in (5.130) is the second term.

We remove the divergence by normal ordering the interaction Hamiltonian $H_{\text{int}} = cm(\tau) : \hat{\phi}^2(\tau) :$ with respect to the Minkowski vacuum. This renormalization procedure is equivalent to subtracting off the divergent Minkowski terms. In (5.130) $\langle 0|\hat{\phi}^2(\tau)\hat{\phi}^2(\tau')|0\rangle$ is thus replaced by

$$\langle 0| : \hat{\phi}^2(\tau) :: \hat{\phi}^2(\tau') : |0\rangle = 2\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle^2 + \langle 0| : \hat{\phi}^2(\tau) : |0\rangle\langle 0| : \hat{\phi}^2(\tau') : |0\rangle \ , \tag{5.131}$$

where

$$\begin{aligned}\langle 0| :\hat{\phi}^2(\tau): |0\rangle &= \langle 0|\hat{\phi}^2(\tau)|0\rangle - \langle 0|\phi^2(\tau)|0\rangle \ , \\ &= \lim_{\tau'\to\tau}\left[\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle - \langle 0|\phi(\tau)\phi(\tau')|0\rangle\right] \ , \\ &= \sum_{\gamma\in\Gamma\setminus\{I_d\}} p(\gamma)\langle 0|\phi(\tau)\phi(\gamma^{-1}\tau)|0\rangle \ . \end{aligned} \tag{5.132}$$

In the last equality in (5.132) we have used the method of images expression (5.74) for the correlation function of the automorphic field.

Consider now, as an example, a uniformly accelerated detector, following the trajectory (5.17), on 4-dimensional Minkowski space with boundary at $x = 0$. We have already seen in section 5.4.3 that the two point function in this case takes the form

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = -\frac{1}{16\pi^2}\frac{1}{\left(\alpha\sinh(\frac{\tau-\tau'}{2\alpha}) - i\epsilon\cosh(\frac{\tau-\tau'}{2\alpha})\right)^2} + \frac{\beta}{16\pi^2\alpha^2}\frac{1}{\cosh^2(\frac{\tau+\tau'}{2\alpha})} \ . \tag{5.133}$$

Further from (5.132) we have

$$\langle 0| :\hat{\phi}^2(\tau): |0\rangle = \frac{\beta}{16\pi^2\alpha^2}\frac{1}{\cosh^2(\frac{\tau}{\alpha})} \ , \tag{5.134}$$

where the $\epsilon \to 0$ limit has been taken before doing the transition rate integral as can be argued by the dominated convergence theorem. Substituting (5.133) and (5.134) into (5.131) and then (5.128) we have a transition rate with 4 terms. The first term comes from the Minkowski parts of the image sum and so leads to the usual thermal spectrum. The other three terms however break the KMS condition leading to a non-thermal response, due to the presence of the boundary. We shall not discuss the calculation further here. For more details of the response function for finite time observations in this case see [108].

5.6 Causal detector for the Dirac field

5.6.1 Minkowski space

In this section we extend the causal detector to the massless Dirac field in 4-dimensional Minkowski space. The detector is still a many-level quantum mechanical system with free Hamiltonian H_D. However now the detector moves through a massless Dirac field ψ (with free Hamiltonian H_ψ) in Minkowski space to which it is coupled via the interaction Hamiltonian

$$H_{\text{int}} = cm(\tau)\bar{\psi}(\tau)\psi(\tau) \ , \tag{5.135}$$

where $\bar{\psi} = \psi^\dagger\gamma^0$, and $\psi(\tau) = \psi(x(\tau))$. The equation of motion for the free field ψ is the massless free Dirac equation $i\gamma^\mu\partial_\mu\psi = 0$. We choose a basis of solutions and expand the field in terms of this basis. We work throughout with the standard representation of γ matrices,

$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \ , \quad \gamma^i = \begin{pmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{pmatrix} \ , \tag{5.136}$$

where σ_i are the Pauli matrices, $\sigma_1 = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, $\sigma_2 = \left(\begin{smallmatrix} 0 & -i \\ i & 0 \end{smallmatrix}\right)$ and $\sigma_3 = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$. Then

$$\psi(t, \mathbf{x}) = \sum_{s=1,2} \int \frac{d^3k}{(2(2\pi)^3)^{\frac{1}{2}}} \left[b_s(\mathbf{k})u(\mathbf{k}, s)e^{-i\omega t + i\mathbf{kx}} + d_s^\dagger(\mathbf{k})v(\mathbf{k}, s)e^{i\omega t - i\mathbf{kx}} \right] \ , \tag{5.137}$$

where

$$u(\mathbf{k}, 1) = \begin{pmatrix} 1 \\ 0 \\ \frac{k_z}{\omega} \\ \frac{k_+}{\omega} \end{pmatrix} \ , \qquad u(\mathbf{k}, 2) = \begin{pmatrix} 0 \\ 1 \\ \frac{k_-}{\omega} \\ \frac{-k_z}{\omega} \end{pmatrix} \ , \tag{5.138}$$

and

$$v(\mathbf{k},1) = \begin{pmatrix} \frac{k_z}{\omega} \\ \frac{k_+}{\omega} \\ 1 \\ 0 \end{pmatrix} , \qquad v(\mathbf{k},2) = \begin{pmatrix} \frac{k_-}{\omega} \\ \frac{-k_z}{\omega} \\ 0 \\ 1 \end{pmatrix} , \tag{5.139}$$

with $k_\pm = k_x \pm ik_y$. The modes in the expansion (5.137) are expressed in terms of a standard Minkowski vierbein, aligned along Minkowski coordinate axes, and they are suitably orthonormal with respect to the usual Dirac inner product in Minkowski space,

$$\langle \psi_1, \psi_2 \rangle = \int d^3x\, \psi_1^\dagger \psi_2 \ . \tag{5.140}$$

The free field is then quantized in the usual manner, imposing the usual anticommutation relations on the annihilation/creation operators,

$$\left\{ b_s(\mathbf{k}), b^\dagger_{s'}(\mathbf{k}) \right\} = \delta_{ss'}\delta(\mathbf{k}-\mathbf{k}') \ , \tag{5.141}$$

and similarly for the d operators, with all mixed anticommutators vanishing.

We assume again that at time τ_0 the full interacting field is in the product state $|0, E_0\rangle = |0\rangle|E_0\rangle$. Working in the interaction picture we find, to first order in perturbation theory, that the probability that at a later time $\tau_1 > \tau_0$ the detector is found in state $|E_1\rangle$ is

$$\begin{aligned} \sum_\Psi |\langle \Psi, E_1|0, E_0\rangle|^2 \quad &= \quad c^2 |\langle E_1|m(0)|E_0\rangle|^2 \int_{\tau_0}^{\tau_1} d\tau \int_{\tau_0}^{\tau_1} d\tau' e^{-i\omega(\tau-\tau')} \\ &\quad \times \langle 0|\bar\psi(\tau)\psi(\tau)\bar\psi(\tau')\psi(\tau')|0\rangle \ , \end{aligned} \tag{5.142}$$

with $\omega = E_1 - E_0$. Once again we shall concentrate on the response function part. With the same change of coordinates as in section 5.3.1, this part may be written as

$$F_{\tau_1,\tau_0}(\omega) = 2\int_{\tau_0}^{\tau_1} du \int_0^{u-\tau_0} ds\, Re(e^{-i\omega s}\langle 0\bar\psi(u)\psi(u)\bar\psi(u-s)\psi(u-s)|0\rangle) \ . \tag{5.143}$$

We have used the relation $\langle 0|\bar{\psi}(\tau)\psi(\tau)\bar{\psi}(\tau')\psi(\tau')|0\rangle = \langle 0|\bar{\psi}(\tau')\psi(\tau')\bar{\psi}(\tau)\psi(\tau)|0\rangle^*$, which follows from the axioms of inner products and $(\bar{\psi}(\tau')\psi(\tau'))^\dagger = \bar{\psi}(\tau')\psi(\tau')$. Finally, as in the scalar case, we let the detector be switched on in the infinite past and differentiate with respect to $\tau_1 = \tau$ to get the transition rate, with the result

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds\, Re\left(e^{-i\omega s}\langle 0|\bar{\psi}(\tau)\psi(\tau)\bar{\psi}(\tau - s)\psi(\tau - s)|0\rangle\right) \ . \tag{5.144}$$

Further here we find

$$\langle 0|\bar{\psi}(\tau)\psi(\tau)\bar{\psi}(\tau')\psi(\tau')|0\rangle = \mathrm{Tr}((S_M^+(\tau,\tau'))^2) \ , \tag{5.145}$$

where Tr is the trace and $S_M^+(\tau,\tau') = \langle 0|\psi(\tau)\bar{\psi}(\tau')|0\rangle$ is the positive frequency Wightman function which is related to the positive frequency scalar Wightman function by (see e.g [1])

$$S_M^+(\tau,\tau') = i\gamma^\mu\partial_\mu G_M^+(\tau,\tau') \ . \tag{5.146}$$

To prove expression (5.145) a straigtforward, but rather messy, calculation was used where the expansion of the field (5.137) was substituted directly into the correlation function $\langle 0|\bar{\psi}(\tau_1)\psi(\tau_2)\bar{\psi}(\tau_3)\psi(\tau_4)|0\rangle$ and use was made of the anticommutation relations (5.141), setting $\tau_1 = \tau_2 = \tau$, $\tau_3 = \tau_4 = \tau'$ at the end. We note here that all expressions for the response are independent of the vierbein used to express the ψ field. This is due to the form of H_{int} (5.135) which is a Lorentz scalar. In the case of a massive Dirac field in Minkowski space (5.145) contains a second term proportional to $\mathrm{Tr}(S_M^+(\tau,\tau'))\mathrm{Tr}(S_M^-(\tau',\tau))$. Here in the massless case this term does not enter as $\mathrm{Tr}(S_M^+(\tau,\tau')) = 0$ for any worldline.

Consider the uniformly accelerated worldline (5.17). Again a numerical calculation shows that if we use the usual correlation function, i.e. with the $i\epsilon$ regularization, for the scalar field (5.15) in the expressions above we will get a τ-dependent result for the transition rate (5.144) even in the $\epsilon \to 0$ limit. We are thus led once again to consider an alternative setup where we use a smeared form for the field operator in the interaction Hamiltonian. That is we consider

$$\psi(\tau) = \int d^3\xi\, W_\epsilon(\xi) S(\tau,\xi)\psi(x(\tau,\xi)) \ , \tag{5.147}$$

with the same definitions for $W_\epsilon(\xi)$ and ξ as in section 5.3.1. In contrast to the scalar case, here we include $S(\tau,\xi)$, which is the spinor transformation associated with the transformation from the Minkowski vierbein to one adapted to the Fermi-Walker coordinates. However we may now argue that in this case $S(\tau,\xi)$ may be dropped. Firstly we note that the metric written in Fermi Walker coordinates is [25]

$$ds^2 = \left(1 + 2\left(\dot{t}\ddot{x} - \dot{x}\ddot{t}\right)\xi + \left(\ddot{x}^2 - \ddot{t}^2\right)\xi^2\right)d\tau^2 - d\xi^2 \ . \tag{5.148}$$

Constant τ spatial sections are therefore flat. It then follows that the transformation from Minkowski vierbein to that adapted to these Fermi coordinates will be independent of ξ, as Fermi Walker transport along these spatial sections in a non rotating vierbein will be trivial. It therefore follows that $S(\tau,\xi)$ may be taken outside the integral in (5.147). Further it follows from the form of H_{int} (5.135) that as S may be taken outside the integral it may be dropped completely. Therefore on M we may work throughout with expressions written with respect to the standard Minkowski vierbein and with

$$\psi(\tau) = \int d^3\xi\, W_\epsilon(\xi)\psi(x(\tau,\xi)) \ , \tag{5.149}$$

as the expression for a smeared field operator. By arguments similar to those that lead to (5.144) and (5.145) the transition rate is again given by

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds\, Re\left(e^{-i\omega s}\langle 0|\bar{\psi}(\tau)\psi(\tau)\bar{\psi}(\tau-s)\psi(\tau-s)|0\rangle\right) \ , \tag{5.150}$$

and we find

$$\langle 0|\bar{\psi}(\tau)\psi(\tau)\bar{\psi}(\tau')\psi(\tau')|0\rangle = \text{Tr}(\langle 0|\psi(\tau)\bar{\psi}(\tau')|0\rangle^2) \ , \tag{5.151}$$

where $\psi(\tau)$ is now the smeared field (5.149).

Now suppose we consider again the Lorentzian profile function

$$W_\epsilon(\xi) = \frac{1}{\pi^2}\frac{\epsilon}{(\xi^2+\epsilon^2)^2} \ . \tag{5.152}$$

The spinor correlation function is then given by

$$\begin{aligned}\langle 0|\psi(\tau)\bar{\psi}(\tau')|0\rangle &= \frac{1}{2(2\pi)^3}\sum_{s=1,2}\int d^3k\, u(\mathbf{k},s)u^\dagger(\mathbf{k},s)\gamma^0 \\ &\quad \times \int d^3\xi\, W_\epsilon(\xi)e^{-ik\cdot x(\tau,\xi)}\int d^3\xi'\, W_\epsilon(\xi')e^{ik\cdot x(\tau,\xi')} \ . \end{aligned}\tag{5.153}$$

The integrals over ξ and ξ' in (5.153) are the same as those in (5.24). Proceeding as with (5.24), we thus have

$$\langle 0|\psi(\tau)\bar{\psi}(\tau')|0\rangle = \frac{1}{2(2\pi)^3}\sum_{s=1,2}\int d^3k\, u(\mathbf{k},s)u^\dagger(\mathbf{k},s)\gamma^0 e^{-i\omega(t-t'-i\epsilon(\dot{t}+\dot{t}'))+i\mathbf{k}(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))} \ . \tag{5.154}$$

Further from (5.138) we have

$$\sum_{s=1,2} u(\mathbf{k},s)u^\dagger(\mathbf{k},s)\gamma^0 = \begin{pmatrix} 1 & 0 & -\frac{k_z}{\omega} & \frac{-k_x+ik_y}{\omega} \\ 0 & 1 & \frac{-k_x-ik_y}{\omega} & \frac{k_z}{\omega} \\ \frac{k_z}{\omega} & \frac{k_x-ik_y}{\omega} & -1 & 0 \\ \frac{k_x+ik_y}{\omega} & -\frac{k_z}{\omega} & 0 & -1 \end{pmatrix} \ . \tag{5.155}$$

If we now compare (5.154) with the scalar field expression (5.33), with $d=4$, we find we have

$$S_M^+(\tau,\tau') = \langle 0|\psi(\tau)\bar{\psi}(\tau')|0\rangle = i\gamma^\mu\partial_\mu\langle 0|\phi(\tau)\bar{\phi}(\tau')|0\rangle \ , \tag{5.156}$$

where $\langle 0|\phi(\tau)\bar{\phi}(\tau')|0\rangle$ is the scalar field correlation function (5.33) and the partial derivative acts on the $(t,\mathbf{x})$ but NOT on the $(\dot{t},\dot{\mathbf{x}})$. Therefore we find, from (5.34), that the spinor correlation function is given by

$$\begin{aligned} S^+_M(\tau,\tau') &= \frac{i}{4\pi^2}\frac{1}{[\tilde{t}^2-\tilde{x}^2-\tilde{y}^2-\tilde{z}^2]^2} \\ &\times \begin{bmatrix} 2\tilde{t} & 0 & -2\tilde{z} & 2(i\tilde{y}-\tilde{x}) \\ 0 & 2\tilde{t} & -2(i\tilde{y}+\tilde{x}) & 2\tilde{z} \\ 2\tilde{z} & -2(i\tilde{y}-\tilde{x}) & 2\tilde{t} & 0 \\ 2(i\tilde{y}+\tilde{x}) & -2\tilde{z} & 0 & 2\tilde{t} \end{bmatrix} , \end{aligned} \tag{5.157}$$

where $\tilde{a} = (a(\tau) - a(\tau') - i\epsilon(\dot{a}(\tau) + \dot{a}(\tau')))$. From this it is easy to show that

$$\mathrm{Tr}(S^+_M(\tau,\tau')^2) = -\frac{1}{\pi^4}\frac{1}{((t-t'-i\epsilon(\dot{t}+\dot{t}'))^2-(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))^2)^3} \ . \tag{5.158}$$

Inertial detector

First we consider the response of a Dirac field detector following the inertial worldline (5.93) in Minkowski space. From (5.158), (5.150) and (5.151) the transition rate is found to be

$$\dot{F}_\tau(\omega) = -\frac{1}{\pi^4}\lim_{\epsilon\to 0}\int_{-\infty}^{\infty} ds\, \frac{e^{-i\omega s}}{(s-2i\epsilon)^6} \ . \tag{5.159}$$

The integral can be done by residues, with the result

$$\dot{F}_\tau(\omega) = -\frac{\omega^5}{60\pi^3}\Theta(-\omega) \ . \tag{5.160}$$

Consider also the "power spectrum" of the Dirac noise as defined by Takagi in [31]. The noise $g(\tau,\tau')$ is defined by

$$g(\tau,\tau') = S(\tau)S^+_M(\tau,\tau')S(\tau')^{-1} \ , \tag{5.161}$$

where $S(\tau) = S(\tau,\xi)$ as given in (5.147). $S(\tau)$ is the spinor transformation which takes care of the Fermi-Walker transport so that $S(\tau)\psi(\tau)$ does not rotate with respect to the detector's proper reference frame. The definition for the power spectrum on a stationary worldline, where $S^+_M(\tau,\tau')$ depends on τ and τ' only through $\tau-\tau'$, is

$$P(\omega) = \frac{1}{4}\mathrm{Tr}\,\gamma^0 \int_{-\infty}^{\infty} d\tau\, e^{-i\omega\tau} g(\tau)\ . \tag{5.162}$$

On the inertial trajectory (5.93), the transformation to the Fermi frame is trivial and we find

$$\begin{aligned} P(\omega) &= \frac{i}{2\pi^2}\int_{-\infty}^{\infty} ds\, \frac{e^{-i\omega s}}{(s-2i\epsilon)^3} \\ &= \frac{\omega^2}{2\pi}\Theta(-\omega)\ . \end{aligned} \tag{5.163}$$

We note that the power spectrum is $-\omega$ times the transition rate for the linearly coupled scalar field detector following the same trajectory.

Uniformly accelerated detector

Considering once again a detector following the uniformly accelerated worldline (5.17). We find, as expected, that the correlation function is invariant under translations in τ and the transition rate (5.150) is given by

$$\dot{F}_\tau(\omega) = -\frac{1}{64\pi^4}\lim_{\epsilon\to 0}\int_{-\infty}^{\infty} ds\, \frac{e^{-i\omega s}}{\left(\alpha\sinh\left(\frac{s}{2\alpha}\right) - i\epsilon\cosh\left(\frac{s}{2\alpha}\right)\right)^6}\ . \tag{5.164}$$

The integral may be done by contour integration (appendix D). The result is

$$\dot{F}_\tau(\omega) = \frac{1}{60\pi^3\alpha^4}\frac{\omega}{(e^{2\pi\alpha\omega}-1)}(4+5(\alpha\omega)^2+(\alpha\omega)^4)\ . \tag{5.165}$$

The response is thermal in the sense that it satisfies the KMS condition at the temperature $T=(2\pi\alpha)^{-1}$. It is interesting to note that there is no fermionic factor in the response, instead we have the usual Planckian factor found in the scalar case.

We can see the fermionic factor appearing however if we consider the power spectrum

(5.162) of the Dirac noise. For the uniformly accelerated worldline we have

$$S(\tau) = \cosh\left(\frac{\tau}{2\alpha}\right) - \gamma^0\gamma^1 \sinh\left(\frac{\tau}{2\alpha}\right) \ , \tag{5.166}$$

and the power spectrum (5.162) is given by

$$P(\omega) = \frac{i}{16\pi^2}\int_{-\infty}^{\infty} d\tau \, \frac{e^{-i\omega\tau}}{\left(\alpha\sinh\left(\frac{\tau}{2\alpha}\right) - i\epsilon\cosh\left(\frac{\tau}{2\alpha}\right)\right)^3} \ . \tag{5.167}$$

Again the integral may be done by contour integration to give

$$P(\omega) = \frac{1}{2\pi}\frac{\left(\omega^2 + \frac{1}{4\alpha^2}\right)}{\left(1 + e^{2\pi\alpha\omega}\right)} \ . \tag{5.168}$$

Our expression here agrees with that found by Takagi [31]. The power spectrum therefore does include the expected fermionic factor.

5.6.2 Dirac detector for automorphic fields

Next we wish to consider this fermionic detector on M_0 and M_- and in particular address the issues concerning spin structure on M_-. We consider an automorphic Dirac field on Minkowski space. The main difference from the scalar case is that we must take care of what vierbeins our expressions are written with respect to. In particular our vierbein might not be invariant under the quotient group Γ.

We begin with a massless Dirac field ψ on M, expressed with respect to a vierbein that is invariant under Γ. The automorphic field is then defined by

$$\hat{\psi}(x) = \frac{1}{\left(\sum_{\gamma\in\Gamma} p(\gamma)^2\right)^{1/2}} \sum_{\gamma\in\Gamma} p(\gamma)\psi(\gamma^{-1}x) \ , \tag{5.169}$$

where the normalization is such that, at equal times,

$$\left\{\hat{\psi}_\alpha(x), \hat{\psi}^\dagger_\beta(x')\right\} = \delta^{(d-1)}(x - x')\delta_{\alpha\beta} + \text{image terms} \ . \tag{5.170}$$

The two-point function for the automorphic field is then given by the method of images,

$$S^{+}_{M/\Gamma}(x,x') = \langle 0|\hat{\psi}(x)\bar{\hat{\psi}}(x')|0\rangle = \sum_{\gamma\in\Gamma} p(\gamma)\langle 0|\psi(x)\bar{\psi}(\gamma^{-1}x')|0\rangle \ . \tag{5.171}$$

We consider a detector coupled to the automorphic field via the interaction Hamiltonian

$$H_{\text{int}} = cm(\tau)\bar{\hat{\psi}}(\tau)\hat{\psi}(\tau) \ , \tag{5.172}$$

where

$$\hat{\psi}(\tau) = \int d^3\xi \, W_\epsilon(\xi) S(\tau,\xi)\hat{\psi}(x(\tau,\xi)) \ . \tag{5.173}$$

The transition rate is given by

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds \, Re\left(e^{-i\omega s}\langle 0|\bar{\hat{\psi}}(\tau)\hat{\psi}(\tau)\bar{\hat{\psi}}(\tau - s)\hat{\psi}(\tau - s)|0\rangle\right) \ . \tag{5.174}$$

Further we may show, with a calculation similar to that leading to (5.145) and (5.151), that

$$\langle 0|\bar{\hat{\psi}}(\tau)\hat{\psi}(\tau)\bar{\hat{\psi}}(\tau')\hat{\psi}(\tau')|0\rangle = \text{Tr}\left(\langle 0|\hat{\psi}(\tau)\bar{\hat{\psi}}(\tau')|0\rangle^2\right) \ , \tag{5.175}$$

with

$$\langle 0|\hat{\psi}(\tau)\bar{\hat{\psi}}(\tau')|0\rangle = \sum_{\gamma\in\Gamma} p(\gamma)\langle 0|\psi(\tau)\bar{\psi}(\gamma^{-1}\tau')|0\rangle \ . \tag{5.176}$$

Therefore the method of images may be directly applied to our Minkowski space correlation functions here.

It is important to note that the above mode sum expressions are changed when considering a vierbein not invariant under the action of Γ. Suppose we consider two vierbeins, one invariant under Γ (labelled by an I) and another not invariant (labelled by N). In the vierbein I the automorphic field is given by the mode sum expression (5.169). The transformation from I to N will transform the spinors as $\hat{\psi}_I(x) \to \hat{\psi}_N(x) = S(x)\hat{\psi}_I(x)$. Then from

(5.169)

$$\hat{\psi}_N(x) = \frac{1}{\left(\sum_{\gamma\in\Gamma} p(\gamma)^2\right)^{1/2}} \sum_{\gamma\in\Gamma} p(\gamma) S(x) \psi_I(\gamma^{-1}x) \ , \tag{5.177}$$

and hence the mode sum expression for the automorphic field in terms of the non-invariant vierbein is

$$\hat{\psi}_N(x) = \frac{1}{\left(\sum_{\gamma\in\Gamma} p(\gamma)^2\right)^{1/2}} \sum_{\gamma\in\Gamma} p(\gamma) S(x) S^{-1}(\gamma^{-1}x) \psi_N(\gamma^{-1}x) \ . \tag{5.178}$$

Similarly the two-point function tranforms as

$$S^{+}_{IM/\Gamma}(x,x') \to S^{+}_{NM/\Gamma}(x,x') = S(x) S^{+}_{IM/\Gamma}(x,x') S^{-1}(x') \ . \tag{5.179}$$

From (5.171), the mode sum expression for the two point function in terms of the non-invariant vierbein is hence

$$\begin{aligned} S^{+}_{NM/\Gamma}(x,x') &= \sum_{\gamma\in\Gamma} p(\gamma) S(x) \langle 0|\psi_I(x)\bar{\psi}_I(\gamma^{-1}x')|0\rangle S^{-1}(x') \ , \\ &= \sum_{\gamma\in\Gamma} p(\gamma) \langle 0|\psi_N(x)\bar{\psi}_N(\gamma^{-1}x')|0\rangle S^{-1}(\gamma^{-1}x') S^{-1}(x') \ . \end{aligned} \tag{5.180}$$

We have seen explicit examples of this in sections 4.3.2 and 4.4.2, and on the $\mathbb{RP}^3$ geon in section 3.4.

In sections 5.6.3 and 5.6.4 we shall work throughout in vierbeins invariant under J_0 and J_- respectively.

5.6.3 Dirac detector on M_0

Consider now Dirac field theory on M_0 as an automorphic field theory on M where expressions are written with respect to the standard Minkowski vierbein. From (5.176) the M_0

correlation function is given by

$$S^{+}_{M_0}(\tau,\tau') = \sum_{n\in\mathbb{Z}} \eta^n S^{+}_{M}(\tau, J_0(\tau')) \ , \tag{5.181}$$

where $\eta = 1, (-1)$ labels spinors with periodic (antiperiodic) boundary conditions. We therefore find an explicit expression for $S^{+}_{M_0}(\tau,\tau')$ from (5.181) and (5.157). As we work throughout in the standard Minkowski vierbein here, writing the smeared field operator as in (5.173), we may again argue in an analogous way to in the previous section that the spinor transformation $S(\tau,\xi)$ can be dropped here.

Inertial detector

First consider a Dirac detector following the inertial worldline (5.93) on M_0. The power spectrum (5.162) for the noise is found to be

$$P(\omega) = \left(\frac{\omega^2}{2\pi} + 2\sum_{n=1}^{\infty} \frac{\eta^n \omega}{4\pi n a} \sin(2n\omega a) \right) \Theta(-\omega) \ . \tag{5.182}$$

As on Minkowski space the power spectrum is $-\omega$ times the transition rate of the linearly coupled scalar field detector following the same trajectory (5.83). The summation thus may be performed to give $-\omega$ times (5.84) and (5.85).

For the transition rate we find

$$\dot{F}_\tau(\omega) = -\frac{1}{\pi^4} \sum_{n\in\mathbb{Z}} \sum_{m\in\mathbb{Z}} \int_{-\infty}^{\infty} ds \, \frac{\eta^n \eta^m ((s-2i\epsilon)^2 - 4nma^2)}{[(s-2i\epsilon)^2 - (2na)^2]^2 [(s-2i\epsilon)^2 - (2ma)^2]^2} \ . \tag{5.183}$$

The $n=0, m=0$ term gives the transition rate on Minkowski space (5.160). The integral for other terms may be done by residues, with the result

$$\begin{aligned}\dot{F}_\tau(\omega) \quad = \quad & \Bigg(-\frac{\omega^5}{60\pi^3} + \frac{1}{32\pi} \sum_{\substack{n.m=-\infty \\ n,m\neq 0}}^{\infty} \frac{\eta^n \eta^m}{(m-n)(m+n)^3 a^5} \Bigg[\Bigg(\left(\frac{2m\omega a}{n} + 2\omega a \right) \cos(2\omega n a) \\ & - \frac{(m+3n)}{n^2} \sin(2\omega n a) \Bigg) + \Bigg(\left(\frac{2\omega a n}{m} + 2\omega a \right) \cos(2\omega m a) \\ & - \frac{(n+3m)}{m^2} \sin(2\omega m a) \Bigg) \Bigg] \Bigg) \Theta(-\omega) \ , \end{aligned} \tag{5.184}$$

where the $n = m$ and $n = -m$ terms are understood in the limiting sense and can be verified to be finite. As expected the response does not depend on the velocity.

Uniformly accelerated detector

Next consider the power spectrum for the Rindler noise, that is we consider $g(\tau, \tau')$ on the uniformly accelerated worldline. From (5.162) we find

$$P(\omega) = \frac{i}{16\pi^2} \int_{-\infty}^{\infty} d\tau \sum_{n\in\mathbb{Z}} \frac{\eta^n e^{-i\omega\tau} \left(\alpha \sinh\left(\frac{\tau}{2\alpha}\right) - i\epsilon \cosh\left(\frac{\tau}{2\alpha}\right) \right)}{\left(\left(\alpha \sinh\left(\frac{\tau}{2\alpha}\right) - i\epsilon \cosh\left(\frac{\tau}{2\alpha}\right) \right)^2 - (na)^2 \right)^2} \ . \tag{5.185}$$

The contributions to the integral from each term in the sum may be calculated separately by contour integration. The result is

$$\begin{aligned} P(\omega) \quad = \quad & \frac{1}{2\pi} \frac{(\omega^2 + \frac{1}{4\alpha^2})}{(1 + e^{2\pi\alpha\omega})} \\ & + 2 \sum_{n=1}^{\infty} \frac{\eta^n}{(1 + e^{2\pi\alpha\omega})} \left[\frac{\alpha^2 n^2 a^2 \cos\left(2\omega\alpha \operatorname{arctanh}\left(\frac{(\alpha^2 n^2 a^2 + n^4 a^4)^{1/2}}{\alpha^2 + n^2 a^2} \right) \right)}{4\pi \left(\frac{\alpha^2}{\alpha^2 + n^2 a^2} \right)^{1/2} (2n^6 a^6 + 4n^4 a^4 \alpha^2 + 2n^2 a^2 \alpha^4)} \right. \\ & \left. + \frac{\alpha^3 (\alpha^2 n^2 a^2 + n^4 a^4)^{1/2} \omega \sin\left(2\omega\alpha \operatorname{arctanh}\left(\frac{(\alpha^2 n^2 a^2 + n^4 a^4)^{1/2}}{\alpha^2 + n^2 a^2} \right) \right)}{2\pi \left(\frac{\alpha^2}{\alpha^2 + n^2 a^2} \right)^{1/2} (2n^6 a^6 + 4n^4 a^4 \alpha^2 + 2n^2 a^2 \alpha^4)} \right] \ , \end{aligned} \tag{5.186}$$

where again η labels the spin structure. We see that the power spectrum is dependent on the spin structure. The $n = 0$ term in (5.186) agrees with the Minkowski space power spectrum (5.168) as expected, and both the $n = 0$ and $n > 0$ terms in (5.186) contain the fermionic factor. Note that no simple relation holds between the power spectrum (5.186) and the transition rate of the linearly coupled scalar field detector (5.86), in contrast to the relation we observed on the inertial worldline.

For the transition rate of a fermionic detector on M_0 we have

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds\, Re\left(e^{-i\omega s}\mathrm{Tr}\left(S^+_{M_0}(\tau,\tau-s)^2\right)\right) \ . \tag{5.187}$$

We may evaluate (5.187) on the uniformly accelerated worldline by substituting the worldline into (5.181) and (5.156). It is easy to show that the $n=0$ term leads to the transition rate found on Minkowski space (5.165) as expected. The evaluation of the other terms is not so straightforward as the residues are not so easy to calculate. We shall not present the result here.

5.6.4 Dirac detector on M_-

On M_- we can again build expressions from those on M (or M_0) via the method of images. The transition rate is given by

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds\, Re\left(e^{-i\omega s}\mathrm{Tr}\left(S^+_{M_-}(\tau,\tau-s)^2\right)\right) \ , \tag{5.188}$$

and

$$\begin{aligned} S^+_{M_-}(\tau,\tau') &= S^+_{M_0}(\tau,\tau') + \rho S^+_{M_0}(\tau, J_-(\tau')) \ , & (5.189) \\ &= \sum_{n\in\mathbb{Z}} \rho^n S^+_M(\tau, J^n_-(\tau')) \ , & (5.190) \end{aligned}$$

where $S^+_{M_0}(\tau,\tau')$ and $S^+_M(\tau,\tau')$ are written in terms of a vierbein which rotates by 2π in the (x,y) plane as $z\mapsto z+2a$ (i.e. the one spin structure on M_0 compatible with the two on M_-). $\rho = 1(-1)$ labels spinors with periodic (antiperiodic) boundary conditions on M_- with respect to this vierbein. That is ρ labels the two possible spin structures on M_-.

Now on M_- our main question of interest is whether or not our detector can distinguish the two possible spin structures. As we saw in section 4.3.2 the stress tensor for the massless spinor field in M_- (4.41) has non-zero shear components, $\langle 0_-|T_{xz}|0_-\rangle$ and $\langle 0_-|T_{yz}|0_-\rangle$, which change sign under a change of spin structure. It is therefore conceivable that a detector with a non-zero z-component of angular momentum could detect the spin structure. However as the relation between $\langle 0_-|T_{\mu\nu}|0_-\rangle$ and the detector response is not clear it is not possible to tell in advance whether or not our detector model will be sensitive to the spin structure.

We consider therefore a detector following the trajectory

$$t = t(\tau) \ , \quad x = x(\tau) \ , \quad y = y_0 \ , \quad z = z_0 \ , \tag{5.191}$$

where $y_0 \neq 0$ and z_0 are constants. First we note that there is no direct analogue of the Rindler noise power spectrum here as the power spectrum is defined in [31] only for stationary trajectories. We therefore look directly at the transition rate. From (5.189) and (5.188) the transition rate will contain four terms. The first term, coming from $\mathrm{Tr}\left(S^+_{M_0}(\tau,\tau')\right)^2$, will give us the same response as on M_0 (for $\eta = -1$ in (5.181), as there expressions are written with respect to the standard Minkowski vierbein). This part is independent of spin structure (ρ) on M_-. The fourth term, coming from $\mathrm{Tr}\left(\left(\rho S^+_{M_0}(\tau, J_-(\tau'))\right)^2\right)$, will also be independent of spin structure, as it contains only $\rho^2 = 1$ in both cases. Thus the only way in which the transition rate may be sensitive to the spin structure on M_- is through the cross terms, $\mathrm{Tr}(\rho S^+_{M_0}(\tau, J_-(\tau'))S^+_{M_0}(\tau,\tau'))$ and $\mathrm{Tr}(\rho S^+_{M_0}(\tau,\tau')S^+_{M_0}(\tau, J_-(\tau')))$. However it is a reasonably straightforward matter to show that these traces are both 0 on the trajectory (5.191), due to simple cancellations in the products of the Wightman functions. Thus we see, even without an explicit calculation on a specific trajectory, that the transition rate cannot depend on ρ, and so the detector is not sensitive to the spin structure, for *any* motion at constant y and z.

Unfortunately an explict evaluation of the transition rate on the inertial or uniformly accelerated worldlines, as on M_0, is difficult to obtain and we shall not discuss it further here.

5.7 Static detectors on the $\mathbb{RP}^3$ geon

In the recent literature Deser and Levin [28,116,117] have presented kinematical arguments for the calculation of the Hawking-Unruh effects in a large class of black hole and cosmological spaces by mapping the trajectories of detectors in these spacetimes to Rindler trajectories in higher dimensional embedding spaces (known as GEMS, or global embedding Minkowski spacetimes) in which these spacetimes have global embeddings. In [116] uniformly accelerated observers in de Sitter and Anti de Sitter space are considered. It is seen that in de Sitter space their experience is thermal with temperature $T = a_5/(2\pi)$ where a_5 is their associated acceleration in the 5-dimensional embedding space. In Anti de Sitter space their experience is thermal provided the acceleration is above a certain threshold. In [117] static observers in Schwarzschild space are considered via a 6-dimensional flat embedding space and the expected temperature and entropy are recovered. In [28] this GEMS approach for the derivations of temperature and entropy is extended to Schwarzschild-(anti) de Sitter and Reissner-Nordström spaces in 4 dimensions and rotating BTZ spaces in 3 dimensions, and the methods of [28] can be readily adapted to other cases. We note that indeed any Einstein geometry has a GEMS [118].

[119] considers GEMS calculations on a large class of higher dimensional black holes, generalising the 4-dimensional results of Deser and Levin (and the results for the 4-dimensional AdS hole and others in [120]). In particular, d-dimensional Schwarzschild and Reissner-Nordstrom in asymptotically flat, de Sitter and Anti de Sitter spaces are discussed. The case of 4-dimensional asymptotically locally anti-de Sitter is particularly interesting as solutions with planar, cylindrical, toroidal and hyperbolic horizon topology exist. The higher dimensional versions of these non-spherical AdS black holes are also considered. Their global embeddings in higher dimensional Minkowski spaces are found and the associated temperatures and entropies obtained. Other references on GEMS come from the group of Hong, Park, Kim, Soh and Oh [120–124]. These include the 4-dimensional AdS hole as mentioned above, static rotating and charged BTZ holes, $(2+1)$ de Sitter holes, scalar tensor theories, charged dilatonic black holes in $1+1$ dimensions, charged and uncharged black strings in

(2+1) dimensions, and a few other cases. A recent paper by Chen and Tian [125] argues that the GEMS approach holds for general stationary motions in curved spacetimes. However these authors argue further that the approach in general fails for non-stationary motions. The example they use is that of a freely falling observer in the Schwarzschild geometry. We note here that although their argument does prove that the GEMS argument is not valid for some non-stationary trajectories by use of an example, it does not prove that the GEMS approach is useless for all such trajectories.

Within the kinematical arguments employed in all the work reviewed above the great simplification in working with these GEMS is that we are mapping situations in curved spacetimes to corresponding ones in a flat spacetime, where calculations are always simpler, both conceptually and technically. It seems reasonable following the success of the GEMS programme that the responses of particle detectors in black hole and cosmological backgrounds could also be related in some way to responses of corresponding detectors in their GEMS. We note immediately that such a mapping of detector responses is clearly not trivial as we would expect different responses to occur due to the different dimensions which the spacetimes and their GEMS have, however some relation is still expected. In this section then we present an argument which should be relevant to the response of a static detector in the single exterior of the $\mathbb{RP}^3$ geon black hole (and the Kruskal spacetime) via an embedding of the Kruskal manifold into a 7-dimensional Minkowski space. This embedding space is different to the 6-dimensional embedding of Kruskal so far used in the GEMS literature [126], but we use it as it is more easily adapted to the $\mathbb{RP}^3$ geon.

We begin by first presenting the embedding (see [99]). The complexified Kruskal manifold M_C here is considered to be an algebraic variety in $\mathbb{C}^7$. With coordinates $(z_1, \ldots, z_7)$ and metric

$$ds^2 = -(dz_1)^2 - (dz_2)^2 - \ldots - (dz_6)^2 + (dz_7)^2 \ , \tag{5.192}$$

z_7 being the timelike coordinate, M_C is determined by

$$\begin{aligned}
(z_6)^2 - (z_7)^2 + 4/3(z_5)^2 &= 16M^2 \ , \\
\left((z_1)^2 + (z_2)^2 + (z_3)^2\right)(z_5)^4 &= 576M^6 \ , \\
\sqrt{3} z_4 z_5 + (z_5)^2 &= 24M^2 \ . \qquad (5.193)
\end{aligned}$$

The Lorentzian section of M_C, denoted by $\hat{M}_L$, is the subset stabilised by $J_L : (z_1, \ldots, z_7) \mapsto (z_1^*, \ldots, z_7^*)$. $\hat{M}_L$ consists of two connected components, one with $z_5 > 0$ and one $z_5 < 0$, both of which are isometric to the Kruskal manifold, which we denote by M_L. An explicit embedding of M_L into $\hat{M}_L$ with $z_5 > 0$ is given by

$$\begin{aligned}
z_1 &= r \sin\theta \cos\phi \ , \\
z_2 &= r \sin\theta \sin\phi \ , \\
z_3 &= r \cos\theta \ , \\
z_4 &= 4M \left(\frac{r}{2M}\right)^{1/2} - 2M \left(\frac{2M}{r}\right)^{1/2} \ , \\
z_5 &= 2M \left(\frac{6M}{r}\right)^{1/2} \ , \\
z_6 &= 4M \left(\frac{2M}{r}\right)^{1/2} \exp\left(-\frac{r}{4M}\right) X \ , \\
z_7 &= 4M \left(\frac{2M}{r}\right)^{1/2} \exp\left(-\frac{r}{4M}\right) T \ , \qquad (5.194)
\end{aligned}$$

with $X^2 - T^2 > -1$ and $r = r(T, X)$ defined as the unique solution to

$$\left(\frac{r}{2M} - 1\right) \exp\left(\frac{r}{2M}\right) = X^2 - T^2 \ . \qquad (5.195)$$

Here (T, X, θ, ϕ) are a set of usual Kruskal coordinates, giving the usual Kruskal metric on M_L. In each of the four regions of M_L, $|X| \neq |T|$, one can introduce as usual local

Schwarzschild coordinates (t, r, θ, ϕ). For $X > |T|$, the transformation reads

$$\begin{aligned} T &= \left(\frac{r}{2M} - 1\right)^{1/2} \exp\left(\frac{r}{4M}\right) \sinh\left(\frac{t}{4M}\right) \ , \\ X &= \left(\frac{r}{2M} - 1\right)^{1/2} \exp\left(\frac{r}{4M}\right) \cosh\left(\frac{t}{4M}\right) \ , \end{aligned} \tag{5.196}$$

where $r > 2M$, and the expressions for $z_1, z_2, \ldots, z_5$ are as in (5.194) while those for z_6 and z_7 become

$$\begin{aligned} z_6 &= 4M \left(1 - \frac{2M}{r}\right)^{1/2} \cosh\left(\frac{t}{4M}\right) \ , \\ z_7 &= 4M \left(1 - \frac{2M}{r}\right)^{1/2} \sinh\left(\frac{t}{4M}\right) \ . \end{aligned} \tag{5.197}$$

Recalling that z_7 is the timelike coordinate in the embedding space, we see immediately that an observer static in the exterior region $X > |T|$ at $r = \text{const}, \theta = \text{const}, \phi = \text{const}$ is a Rindler observer in the 7-dimensional embedding space with $(z_1, \ldots, z_5)$ constant and acceleration in the z_6-direction of magnitude

$$a = 1/\alpha = \frac{1}{4M\left(1 - \frac{2M}{r}\right)^{1/2}} \ . \tag{5.198}$$

As we have seen, the response of such a Rindler detector in the embedding space is thermal with the associated temperature

$$T = \frac{a}{2\pi} = \frac{1}{2\pi\alpha} = \frac{1}{8\pi M\left(1 - \frac{2M}{r}\right)^{1/2}} \ . \tag{5.199}$$

This gives the Hawking temperature as seen by the static observer in the black hole spacetime. The associated black hole temperature, i.e. the temperature as seen at infinity, in the Kruskal spacetime is given by the Tolman relation

$$T_0 = g_{00}^{1/2} T = \frac{1}{8\pi M} \ , \tag{5.200}$$

(5.199) and (5.200) are the expected expressions on Kruskal space [1]. Thus the black hole temperature as seen by a static observer has been derived from the Unruh temperature seen by the associated Rindler observer in the global embedding Minkowski spacetime.

Next we consider the $\mathbb{RP}^3$ geon. This is built as a quotient of the Kruskal manifold under the involutive isometry $J_G : (T, X, \theta, \phi) \mapsto (T, -X, \pi - \theta, \phi + \pi)$. We now extend the action of the group generated by J_G to the 7-dimensional embedding space M_C in order to obtain a suitable embedding space for the geon. A suitable extension of J_G is $\bar{J}_G : (z_1, z_2, z_3, z_4, z_5, z_6, z_7) \mapsto (-z_1, -z_2, -z_3, z_4, z_5, -z_6, z_7)$, which is an involution on M_C. Again the worldline of a static detector in the $\mathbb{RP}^3$ geon exterior $X > |T|$ is mapped to the worldline of a Rindler observer with acceleration in the z_6-direction with magnitude (5.198) in this embedding space. We suggest therefore that the calculations of the time dependent responses of an accelerated observer in the d-dimensional quotients of Minkowski space, done in section 5.4.4, should have relevance to the response of a static detector in the exterior of the $\mathbb{RP}^3$ geon (although the exact nature of the relationship is not clear). In particular if we specialise the results of section 5.4.4 to a detector with uniform acceleration (5.198) in the quotient of a 7-dimensional Minkowski space under involution $\bar{J}_G$ we see that the response has two parts. The thermal time-independent part is given by (5.53), which in the present case reads

$$\dot{F}_{M\tau}(\omega) = \frac{a^4}{64\pi^2(e^{\frac{2\pi\omega}{a}} + 1)}(1/4 + \omega^2/a^2)(9/4 + \omega^2/a^2) \ . \tag{5.201}$$

Again this is a thermal response associated with a temperature $T = \frac{a}{2\pi}$. Clearly the response is different to that of the static detector on Kruskal due to the higher number of dimensions though the two should be related. The image part of the response for this Rindler detector is, in the case of a detector switched on in the infinite past,

$$\dot{F}_{I\tau}(\omega) = \frac{3}{8\pi^3}\int_0^\infty ds \, \frac{\cos(\omega s)}{\left(\frac{4}{a^2}\cosh^2\left(\frac{a(2\tau - s)}{2}\right) + 4r^2\right)^{5/2}} \ . \tag{5.202}$$

The total response for certain values of the parameters is shown in figure 5.9. The



Figure 5.9: Transition rate for a detector uniformly accelerated in the z_1 direction on the quotient of 7-dimensional Minkowski space under the involution $\bar{J}_G$. The parameters are $\alpha = 1$, $C = (2z_1)^2 + (2z_2)^2 + (2z_3)^2 = 4r^2 = 64/9$, and $\omega = 1$ (upper curve), $\omega = 1.5$ and $\omega = 2$ (lower curve).

comments of section 5.4 then follow. The image part consists of a term periodic in τ with period π/ω plus a term bounded by a function which dies off exponentially for large τ. The numerical evidence exhibits behaviour qualitatively very similar to that of figure 5.5. The comments made in the section 5.4.3 about finite time detections also follow here. In particular the oscillatory behaviour of the boundary part as $\tau \to \infty$ is a property only of the case of infinite time detection. For a detector switched on at $-\infty < \tau_0 < 0$ the transition rate oscillates for some time period with $\tau > 0$, but eventually it will fall to the thermal response and so at late times the difference between the response on the Minkowski space and that on the quotient space vanishes. This implies for finite time static detectors on the $\mathbb{RP}^3$ geon the difference between the response there and that on Kruskal spacetime falls off also to 0 at late times, which is in agreement with the comments made in [22]. Note however the different behaviour in the case of the infinite time detection.

5.8 Detectors in conformally flat spacetimes

In this section we consider monopole detectors in conformally flat spacetimes coupled to a massless conformally coupled scalar field. Conformally flat spacetimes are of particular interest as they include all two-dimensional spacetimes and all spatially flat Friedmann-

Robertson-Walker models, which include a particular coordinatization of half of the de Sitter manifold as a special case. We will be considering detectors in de Sitter space, and $\mathbb{RP}^3$ de Sitter space, in section 5.9. The main aim of this section is to argue for a regularization of the Wightman function which is similar to that considered for the scalar field in Minkowski space in the earlier sections. Transition rates of monopole detectors can then be calculated for detectors travelling along arbitrary trajectories. The discussion at the start of this section follows that of [1].

The metric on a conformally flat spacetime satisfies

$$g_{\mu\nu} = \Omega^2(x)\eta_{\mu\nu} \ , \tag{5.203}$$

where $\eta_{\mu\nu}$ is the usual Minkowski metric and $\Omega(x)$ is a continuous, non-vanishing, finite, real function. We use associated Minkowski coordinates $(\eta, \mathbf{x})$ in which (5.203) reads

$$ds^2 = \Omega^2(d\eta^2 - d\mathbf{x} \cdot d\mathbf{x}) \ . \tag{5.204}$$

We consider here a conformally coupled massless scalar field. The wave equation in d dimensions is

$$\left[\Box + \frac{(d-2)R}{4(d-1)}\right]\phi = 0 \ . \tag{5.205}$$

From (5.203) it follows that (5.205) takes the form

$$\eta^{\mu\nu}\partial_\mu\partial_\nu\bar{\phi} = 0 \ , \tag{5.206}$$

where $\bar{\phi} = \Omega^{(d-2)/2}\phi$. The exponent $(d-2)/2$ is called the conformal weight of the scalar field. Equation (5.206) has the familiar Minkowski form and admits the usual plane wave mode solutions,

$$\bar{\phi}_{\mathbf{k}}(\eta, \mathbf{x}) = \frac{1}{(2\omega(2\pi)^{d-1})^{1/2}} e^{-i\omega\eta + i\mathbf{k}\cdot\mathbf{x}} \ , \tag{5.207}$$

where $\omega = |\mathbf{k}|$. These modes are positive frequency with respect to the timelike conformal

Killing vector ∂_η. A set of mode solutions to (5.205) are then clearly given by

$$\phi_{\mathbf{k}}(\eta, \mathbf{x}) = \frac{1}{(2\omega(2\pi)^{d-1}\Omega^{d-2}(\eta, \mathbf{x}))^{1/2}} e^{-i\omega\eta + i\mathbf{k}\cdot\mathbf{x}} \; . \tag{5.208}$$

The field is then expanded in these mode solutions

$$\phi(x) = \int d^{d-1}k \left[a_{\mathbf{k}}\phi_{\mathbf{k}}(x) + a^\dagger_{\mathbf{k}}\phi^*_{\mathbf{k}}(x) \right] \; , \tag{5.209}$$

and canonically quantized in the usual way, by imposing the usual commutation relations on annihilation and creation operators. The vacuum state annihilated by all the annihilation operators above $|0\rangle$ is known as the conformal vacuum state.

Now we consider a model monopole particle detector coupled to this conformally invariant scalar field. The model of the unregularized detector is similar to that in the previous sections. The transition rate for a detector turned on at time τ_0, when the field was in the conformal vacuum, and off at $\tau > \tau_0$, is hence

$$\dot{F}_\tau(\omega) = 2 \int_0^{\tau - \tau_0} ds \, Re \left(e^{-i\omega s} \langle 0|\phi(\tau)\phi(\tau - s)|0\rangle \right) \; . \tag{5.210}$$

From (5.209) it follows that the Wightman function in (5.210) is given by

$$\langle 0|\phi(x)\phi(x')|0\rangle = \int d^{d-1}k \, \phi_{\mathbf{k}}(x)\phi^*_{\mathbf{k}}(x') \tag{5.211}$$

$$= \Omega^{(2-d)/2}(x) \int d^{d-1}k \, \bar{\phi}_{\mathbf{k}}(x)\bar{\phi}^*_{\mathbf{k}}(x')\Omega^{(2-d)/2}(x') \; . \tag{5.212}$$

The curved spacetime Wightman function is hence related to the Minkowski Wightman function by

$$\langle 0|\phi(x)\phi(x')|0\rangle = \Omega^{(2-d)/2}(x)\langle 0|\bar{\phi}(x)\bar{\phi}(x')|0\rangle\Omega^{(2-d)/2}(x') \; . \tag{5.213}$$

We now ask: What is a suitable regularization for (5.211) on a given trajectory $x = x(\tau)$?

That is, we wish to regularize the integral

$$\begin{aligned}\langle 0|\phi(x(\tau))\phi(x(\tau'))|0\rangle &= \frac{\Omega^{(2-d)/2}(x(\tau))\Omega^{(2-d)/2}(x(\tau'))}{(2\pi)^{d-1}} \\ &\quad \times \int \frac{d^{d-1}k}{2|\mathbf{k}|}\, e^{-i|\mathbf{k}|(\eta(\tau)-\eta(\tau'))+i\mathbf{k}\cdot(\mathbf{x}(\tau)-\mathbf{x}(\tau'))} \;. \qquad (5.214)\end{aligned}$$

We argue, as with the case discussed in Minkowski space, that a naive $i\epsilon$-prescription where we just introduce an ultraviolet cut-off factor $e^{-\epsilon|\mathbf{k}|}$ (as seems to have been considered by Birrell and Davies in the case of de Sitter space in the spatially flat coordinates [1]) is not suitable here. Indeed we have shown via a numerical analysis that in the case of a comoving observer in de Sitter space in the spatially flat coordinates the transition rate for a detector switched on in the infinite past is time dependent. We interpret this, as Schlicht does for the uniformly accelerated detector in Minkowski space, as an unphysical result.

We introduce instead a regularization based on a frequency cut-off in a frame adapted to the detector's motion. Our proposal is to take the mode solutions in the curved space (5.208) and multiply them by the factor $\Omega^{(d-2)/2}(\eta, \mathbf{x})$ to give the modes (5.207). For a given mode ϕ (5.207) we then define the frequency as seen by the moving detector operationally by

$$i\frac{d\phi}{d\tau} = \omega'\phi \;, \qquad (5.215)$$

where τ is the detector's proper time in the curved space. We thus find from (5.207) that $\omega' = \omega\dot{\eta}(\tau) - \mathbf{k}\cdot\dot{\mathbf{x}}(\tau)$ and we regularize the integral (5.214) by introducing a cut-off of the form $e^{-\epsilon(\omega(\dot{\eta}(\tau)+\dot{\eta}(\tau'))-\mathbf{k}\cdot(\dot{\mathbf{x}}(\tau)+\dot{\mathbf{x}}(\tau')))}$. The integral is then exactly as in (5.33) and the result is just the correlation function with conformal relation (5.213) to (5.34) and (5.36). That is,

$$\begin{gathered}\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \tfrac{\Gamma[d/2-1]}{4\pi^{d/2}}\tfrac{\Omega^{(2-d)/2}(x(\tau))\Omega^{(2-d)/2}(x(\tau'))}{A^{d/2-1}} \;, \\ A = \left[i^2(\eta(\tau)-\eta(\tau') - i\epsilon(\dot{\eta}(\tau)+\dot{\eta}(\tau')))^2 + (\mathbf{x}(\tau)-\mathbf{x}(\tau') - i\epsilon(\dot{\mathbf{x}}(\tau)+\dot{\mathbf{x}}(\tau')))^2\right] \;, \end{gathered} \qquad (5.216)$$

for $d > 2$, while for $d = 2$ the expression is just (5.36) with t replaced by η.

If we were to define the frequency as seen by the detector by

$$i\frac{d\bar{\phi}}{d\tau} = \omega' \bar{\phi} \ , \tag{5.217}$$

that is, using (5.208) we would find $\omega' = \omega\dot{\eta}(\tau) - \mathbf{k} \cdot \dot{\mathbf{x}}(\tau) - i\frac{(d-2)}{2}\frac{\dot{\Omega}(\tau)}{\Omega(\tau)}$. This frequency would have an additional imaginary part. Introducing a cut-off $e^{-\epsilon(\omega'(\tau)+\omega'(\tau'))}$ into (5.214) would give a correlation function without the relation (5.213) to the Minkowski one. Our prescription leading to (5.216) is then equivalent to introducing the cut-off $e^{-\epsilon \mathrm{Re}(\omega'(\tau)+\omega'(\tau'))}$, where Re denotes the real part. It is interesting to note that in the cases considered in section 5.9, that is, the comoving and uniformly accelerated observers on de Sitter space, the imaginary part of ω' is found to be a constant, which then does not contribute to the transition rate integrals, and thus makes no difference to the results in the $\epsilon \to 0$ limit. On a more general trajectory however the imaginary part of ω', defined by (5.217), could depend on τ and so could conceivably affect the transition rate.

We shall examine the consequences of the proposal (5.216) in section 5.9. In particular we shall recover the expected thermal responses for comoving and uniformly accelerated detectors in de Sitter space. Further we will show that for the comoving observer, in the coordinates that admit flat spatial sections, this regularization may also arise from considering a spatially averaged field operator and so from a model of an extended detector.

5.9 De Sitter and $\mathbb{RP}^3$ de Sitter spaces

In this section we begin by considering a model detector in de Sitter space. Throughout we consider a conformally coupled massless scalar field moving through the Euclidean vacuum [127]. We see that here a similar situation is encountered to that found by Schlicht for the uniformly accelerated detector in Minkowski space. It is seen that in the case of a comoving detector switched on in the infinite past and off at finite τ, if the correlation function is regularized by a naive $i\epsilon$ prescription, as for example is done in Birrell and Davies [1], we are led to an unphysical, τ-dependent response. We therefore use the regularization introduced in section 5.8. Further, for a comoving detector, we show that such a regularization can arise also by considering a model detector with spatial extent, that is by considering a smeared field operator in the interaction Hamiltonian. We recover the usual time independent thermal response for comoving and uniformly accelerated detectors.

Subsection 5.9.2 then considers comoving observers in $\mathbb{RP}^3$ de Sitter space, such that the motion is orthogonal to the distinguished foliation [100]. In addition to the thermal part seen in de Sitter space, the transition rate contains an image part, related to that found in section 5.4 for a uniformly accelerated detector on a 4-dimensional Minkowski space with a planar boundary. We also address a comoving detector in de Sitter and $\mathbb{RP}^3$ de Sitter space in a GEMS approach, by considering the response of the associated uniformly accelerated detectors in higher dimensional Minkowski (with boundaries in the case of $\mathbb{RP}^3$ de Sitter) embedding spaces. As we are able to do the calculations both in the original curved spaces and in the global embedding spaces, the results help to clarify the relation and validity of relating detector responses to those in embedding spaces.

5.9.1 Detectors in de Sitter space

We represent d-dimensional de Sitter space as the hyperboloid

$$z_0^2 - z_1^2 - \cdots - z_d^2 = -\alpha^2 \ , \tag{5.218}$$

embedded in the $d+1$-dimensional Minkowski space,

$$ds^2 = dz_0^2 - dz_1^2 - \cdots - dz_d^2 \ , \tag{5.219}$$

with z_i real-valued coordinates. Let us consider the coordinates $(t, \mathbf{x})$ defined by

$$\begin{aligned} z_0 &= \alpha \sinh(t/\alpha) + \frac{e^{t/\alpha}}{2\alpha}|\mathbf{x}|^2 \ , \\ z_d &= \alpha \cosh(t/\alpha) - \frac{e^{t/\alpha}}{2\alpha}|\mathbf{x}|^2 \ , \\ z_i &= e^{t/\alpha} x_i \ . \end{aligned} \tag{5.220}$$

These coordinates cover the half of the de Sitter hyperboloid given by $z_0 + z_d > 0$. The line element is that of a d-dimensional Friedman-Robertson-Walker spacetime with exponentially expanding flat spatial sections,

$$ds^2 = dt^2 - e^{2t/\alpha} \left(dx_1^2 + \cdots + d_{d-1}^2 \right) \ . \tag{5.221}$$

Introducing the conformal time $\eta = -\alpha e^{-t/\alpha}$, the line element becomes conformal to Minkowski,

$$ds^2 = \frac{\alpha^2}{\eta^2} \left[d\eta^2 - \sum_{i=1}^{d-1} (dx_i)^2 \right] \ , \tag{5.222}$$

where $-\infty < \eta < 0$.

Consider a conformally coupled scalar field in the line element (5.222). As we have a conformally coupled field in a conformally flat spacetime the situation is conformally trivial, and we shall apply the discussion in section 5.8, with $\Omega^2(x) = \alpha^2/\eta^2$. The vacuum state associated with a complete set of modes positive frequency with respect to conformal Killing time η, that is the conformal vacuum, coincides with the state known as the Euclidean vacuum [1]. The Euclidean vacuum $|0_E\rangle$ is uniquely characterised as the state whose correlation function $\langle 0_E|\phi(x)\phi(x')|0_E\rangle$ is invariant under the connected component of the de Sitter group, and the only singularities of the correlation function are when x' is on the lightcone of x [127]. Even though we have here defined the Euclidean vacuum in coordinates that only cover half of the de Sitter hyperboloid, it is worth mentioning that the state is well defined on the whole hyperboloid [1].

We shall now specialize to 4-dimensional de Sitter space (although the extension to higher dimensions is straightforward) and consider a uniformly accelerated detector following the worldline

$$\begin{aligned} z_0 &= \alpha \sinh(t/\alpha) + \frac{e^{-t/\alpha}}{2\alpha} r^2 \ , \\ z_4 &= \alpha \cosh(t/\alpha) - \frac{e^{-t/\alpha}}{2\alpha} r^2 \ , \\ z_1 &= z_2 = 0 \ , \\ z_3 &= r, \end{aligned} \tag{5.223}$$

with $r =$ constant. The worldline of such an observer in the embedding space is a hyperbola $(z_4)^2 - (z_0)^2 = \alpha^2 - r^2$, $z_1 = z_2 = 0$, $z_3 = r$. In the de Sitter space the observer has constant proper acceleration a, where $a^2 = -g_{\mu\nu}\dot{u}^\mu \dot{u}^\nu$, $\dot{u}^\mu = u^\nu(\nabla_\nu u^\mu)$ and u^μ is the tangent vector of the trajectory, of magnitude

$$a = \frac{r}{\alpha(\alpha^2 - r^2)^{1/2}} \ . \tag{5.224}$$

The proper time for the accelerated observer is $\tau = (\alpha^2 - r^2)^{1/2} t/\alpha$. The response function for the detector is

$$\dot{F}_\tau(\omega) = 2 \int_0^{\tau - \tau_0} ds \, Re \left(e^{-i\omega s} \langle 0_E | \phi(\tau)\phi(\tau - s) | 0_E \rangle \right) \ , \tag{5.225}$$

and

$$\begin{gathered} \langle 0_E | \phi(\tau)\phi(\tau') | 0_E \rangle = -\tfrac{1}{4\pi^2\alpha^2} \tfrac{\eta(\tau)\eta'(\tau)}{A} \\ A = [(\eta(\tau) - \eta(\tau') - i\epsilon(\dot{\eta}(\tau) + \dot{\eta}(\tau'))^2 - (\mathbf{x}(\tau) - \mathbf{x}(\tau') - i\epsilon(\dot{\mathbf{x}}(\tau) + \dot{\mathbf{x}}(\tau')))^2] \ . \end{gathered} \tag{5.226}$$

Substituting (5.223) into (5.226) we find

$$\langle 0_E | \phi(\tau)\phi(\tau') | 0_E \rangle = -\frac{1}{16\pi^2 \left((\alpha^2 - r^2)^{1/2} \sinh\left(\frac{\tau - \tau'}{2(\alpha^2 - r^2)^{1/2}} \right) - i\epsilon \cosh\left(\frac{\tau - \tau'}{2(\alpha^2 - r^2)^{1/2}} \right) \right)^2} \ , \tag{5.227}$$

and the transition rate of the detector switched on in the infinite past and off at time τ is independent of τ and is given by

$$\dot{F}_\tau(\omega) = \frac{\omega}{2\pi(e^{2\pi\omega(\alpha^2-r^2)^{1/2}} - 1)} \ . \tag{5.228}$$

The accelerated detector thus experiences a thermal response at temperature

$$T = \frac{1}{2\pi}\left(\frac{1}{\alpha^2} + a^2\right)^{1/2} \ . \tag{5.229}$$

The response of a comoving detector is obtained by setting $a = 0$. The transition rate is still thermal at temperature $T = 1/(2\pi\alpha)$. These results agree with the previous literature (e.g [1, 116]). What is new is that we have obtained these results in a causal way for a detector switched on in the infinite past and read at a finite time, as opposed to the case usually considered of a detection over the entire worldline.

We end this section by showing that the regularization in (5.226), in the case of a comoving observer, may be obtained by considering the monopole detector as the limit of an extended detector in de Sitter space. The reason why this is simple in the case of a comoving observer but not for other trajectories is that spatial hypersurfaces of constant t in the coordinates $(t, \mathbf{x})$ are flat Euclidean spaces, which allow us to introduce a detector with infinite spatial extent along these slices. Care must be taken when defining the shape function for the detector however, because the hypersurfaces are expanding with increasing t, with a shape which is rigid in the proper distance the regularization follows. The averaging over spatial hypersurfaces in effect introduces a short distance, high frequency cut-off in the modes.

The detector model is that of section 5.3.1. It is a multi-level quantum mechanical system coupled to a massless conformally coupled scalar field via the interaction Hamiltonian

$$H_{\text{int}} = cm(\tau)\phi(\tau) \ . \tag{5.230}$$

We consider a detector following the trajectory $t = \tau$ that is $\eta = -\alpha e^{-\tau/\alpha}$ and $\mathbf{x} = 0$. In (5.230) we consider the field smeared with a detector profile function over constant τ hypersurfaces, that is

$$\phi(\tau) = \int d^3x\, W_\epsilon(\mathbf{x})\phi(\tau, \mathbf{x}) \ . \tag{5.231}$$

For the profile function we choose

$$W_\epsilon(\mathbf{x}) = \frac{1}{\pi^2}\frac{\epsilon e^{-\tau/\alpha}}{(\mathbf{x}^2 + \epsilon^2 e^{-2\tau/\alpha})^2} \ . \tag{5.232}$$

The detector shape (5.232) is now time dependent! The reason for this is that we want a detector which is rigid in its rest frame. That is, we want a detector which is rigid with respect to proper distance and not comoving distance. The two distances are related by $L_{\text{prop}} = e^{\tau/\alpha}L_{\text{comov}}$. In (5.231) the integration is done over $\mathbf{x}$, which is a comoving coordinate, and using a time independent shape function there would mean that the detector is rigid with respect to comoving distance. It is a simple matter to show that a shape function which selects a distance scale L' may be obtained from one which selects a distance scale L by

$$W_{L'}(\mathbf{x}) = \frac{L^3}{L'^3}W_L\left(\frac{L}{L'}\mathbf{x}\right) \ . \tag{5.233}$$

If we write (5.232) now in terms of proper distance we find

$$W_{\epsilon_{\text{prop}}}(\xi) = \frac{1}{\pi^2}\frac{\epsilon_{\text{prop}}}{\left(\mathbf{x}^2 + \epsilon_{\text{prop}}^2\right)^2} \ , \tag{5.234}$$

and so in terms of proper distance (5.232) is a rigid shape in the sense that it is time independent. Using this shape function we find, using the mode expansion of ϕ,

$$\begin{aligned}\langle 0_E|\phi(\tau)\phi(\tau')|0_E\rangle &= \frac{\eta\eta'}{(2\pi)^3\alpha^2}\int\frac{d^3k}{2\omega}\int d^3x\, W_\epsilon(\mathbf{x})e^{-i(\omega\eta(\tau)-\mathbf{k}\cdot\mathbf{x})} \\ &\quad\times\int d^3x'\, W_\epsilon(\mathbf{x}')e^{i(\omega\eta(\tau')-\mathbf{k}\cdot\mathbf{x})} \ .\end{aligned} \tag{5.235}$$

Further we find

$$\begin{aligned} g_\epsilon(\mathbf{k};\tau) &= \int d^3x\, W_\epsilon(\mathbf{x}) e^{-i(\omega\eta(\tau)-\mathbf{k}\cdot\mathbf{x})} \ , \\ &= e^{-i|\mathbf{k}|\eta(\tau)} e^{-\epsilon|\mathbf{k}|e^{-\tau/\alpha}} \\ &= e^{-i|\mathbf{k}|\eta(\tau)} e^{-\epsilon|\mathbf{k}|\dot{\eta}(\tau)} \ , \end{aligned} \tag{5.236}$$

where the integration is done by transforming to spherical coordinates (appendix B). Hence

$$\langle 0_E|\phi(\tau)\phi(\tau')|0_E\rangle = \frac{\eta\eta'}{(2\pi)^3\alpha^2} \int \frac{d^3k}{2\omega}\, e^{-i\omega(\eta-\eta'-i\epsilon(\dot{\eta}+\dot{\eta}'))} \ . \tag{5.237}$$

The expression (5.237) agrees with that found above from the ultraviolet cut-off regularization.

5.9.2 $\mathbb{RP}^3$ de Sitter space

In this section we consider an inertial detector that is linearly coupled to a conformally coupled massless scalar field in $\mathbb{RP}^3$ de Sitter spacetime [100]. [8]

$\mathbb{RP}^3$ de Sitter space is built as a quotient of de Sitter space under the group generated by the discrete isometry

$$J : (z_0, z_1, z_2, z_3, z_4) \mapsto (z_0, -z_1, -z_2, -z_3, -z_4) \ , \tag{5.238}$$

which induces a map $\tilde{J}$ on the hyperboloid (5.218). Although J has fixed points on M, $\tilde{J}$ acts freely on the hyperboloid. The isometry group of 4-dimensional de Sitter space is $O(1,4)$, being the largest subgroup of the isometry group of the 5-dimensional Minkowski embedding space which preserves (5.218). The isometry group of $\mathbb{RP}^3$ de Sitter space is then the largest subgroup of $O(1,4)$ which commutes with J. That is, $\mathbb{Z}_2 \times O(4)/\mathbb{Z}_2$ where the non-trivial element of the first $\mathbb{Z}_2$ factor sends z_0 to $-z_0$ while the non-trivial element of the $\mathbb{Z}_2$ in the second factor is given by J which clearly acts trivially on $\mathbb{RP}^3$ de Sitter space. The connected component of the isometry group is $SO(4)$. The foliation given by $z_0 =$ constant hypersurfaces is a geometrically distinguished one as it is the only foliation whose spacelike hypersurfaces are orbits of the connected component of the isometry group. This is made clearer by introducing the globally defined coordinates (t, χ, θ, ϕ)

---

[8] See also [101] for a nice discussion on de Sitter space vs $\mathbb{RP}^3$ de Sitter.

$$\begin{aligned} z_0 &= \alpha \sinh(t/\alpha) \ , \\ z_4 &= \alpha \cosh(t/\alpha) \cos\chi \ , \\ z_1 &= \alpha \cosh(t/\alpha) \sin\chi \cos\theta \ , \\ z_2 &= \alpha \cosh(t/\alpha) \sin\chi \sin\theta \cos\phi \ , \\ z_3 &= \alpha \cosh(t/\alpha) \sin\chi \sin\theta \sin\phi \ , \end{aligned} \tag{5.239}$$

in which the metric reads

$$ds^2 = dt^2 - \alpha^2 \cosh^2(t/\alpha)[d\chi^2 + \sin^2\chi(d\theta^2 + \sin^2\theta\, d\phi^2)] \ , \tag{5.240}$$

where (χ, θ, ϕ) on de Sitter ($\mathbb{RP}^3$ de Sitter) space are a set of hyperspherical coordinates on S^3 ($\mathbb{RP}^3$) respectively. (t, χ, θ, ϕ) make manifest the $O(4)$ isometry subgroup.

We denote by $|0_G\rangle$ the vacuum state induced by the Euclidean vacuum $|0_E\rangle$ on de Sitter space (see [100] for more details). We consider a particle detector that is linearly coupled to a massless conformally coupled scalar field. The detector and field are assumed to be in the states $|E_0\rangle$ and $|0_G\rangle$ respectively at time $\tau_0 = -\infty$, and we seek the probability that at time $\tau > \tau_0$ the detector is found in the state $|E_1\rangle$. Through arguments analogous to those in section 5.3.1, we find that the transition rate is

$$\dot{F}_\tau(\omega) = 2\int_0^\infty ds\, Re\left(e^{-i\omega s}\langle 0_G|\phi(\tau)\phi(\tau - s)|0_G\rangle\right) \ . \tag{5.241}$$

By the method of images we have

$$\langle 0_G|\phi(x)\phi(x')|0_G\rangle = \langle 0_E|\phi(x)\phi(x')|0_E\rangle + \langle 0_E|\phi(x)\phi(Jx')|0_E\rangle \ , \tag{5.242}$$

where on the RHS expressions live in de Sitter space, and the correlation function $\langle 0_E|\phi(x)\phi(x')|0_E\rangle$ is given by (5.226).

Consider now a detector that follows the geodesic worldline

$$\begin{aligned} z_0 &= \alpha\sinh(\tau/\alpha)\ , \\ z_4 &= \alpha\cosh(\tau/\alpha)\ , \\ z_1 &= z_2 = z_3 = 0\ . \end{aligned} \tag{5.243}$$

On $\mathbb{RP}^3$ de Sitter space this represents the motion of any geodesic observer whose motion is orthogonal to the distinguished foliation. The transition rate (5.241) is in two parts, a de Sitter part and an image part. We have calculated already the de Sitter part, coming from the first term in (5.242), in section 5.9.1. The result was the usual thermal, Planckian, response at temperature $T = 1/(2\pi\alpha)$. We need the image term. In order to find $\langle 0_E|\phi(x)\phi(Jx')|0_E\rangle$ on this worldline we first write $\langle 0_E|\phi(x)\phi(x')|0_E\rangle$ in terms of the coordinates $(z_0, z_1, z_2, z_3, z_4)$ of the embedding space and then act on x' with J, finding

$$\langle 0_E|\phi(x)\phi(Jx')|0_E\rangle = \frac{1}{8\pi^2}\frac{1}{(+\mathbf{z}\cdot\mathbf{z}' + z_0 z_0' + \alpha^2)}\ , \tag{5.244}$$

where $\mathbf{z} = (z_1, z_2, z_3, z_4)$. The regularization has been omitted as in the image part of the correlation function no regularization is required: The geometric reason for this, as with the uniformly accelerated detector on M with boundary, is that the wordline and its image under J are completely spacelike separated. The image term gives to the transition rate the contribution

$$\dot{F}_{I\tau}(\omega) = 2\int_0^\infty ds\, \frac{\cos(\omega s)}{16\pi^2\alpha^2\cosh^2\left(\frac{2\tau - s}{2\alpha}\right)}\ . \tag{5.245}$$

We see that the image term contribution (5.245) is exactly the same as the image term contribution in the response of a uniformly accelerated detector on a 4-dimensional Minkowski space with boundary at $x = 0$ ((5.101) with $d = 4$). Therefore the total response of our inertial detector in $\mathbb{RP}^3$ de Sitter space (with Ricci scalar $R = 12/\alpha^2$) is identical to the response of a uniformly accelerated detector travelling in 4-dimensional Minkowski space with a boundary at $x = 0$ with the acceleration $1/\alpha$ perpendicular to the boundary. It follows that our numerical results in figure 5.5 also give the response on $\mathbb{RP}^3$ de Sitter space, with the appropriate interpretation for α. In particular, the image term breaks the KMS condition and the response is non-thermal and non-Planckian. When the detector is switched on in the infinite past, the response at large τ is oscillatory in τ with period π/ω. When the detector is switched on at a finite time τ_0, $\tau_0 \ll -1$, numerical evidence indicates that the response is approximately periodic in the region $0 < \tau < -\tau_0$, with period π/ω, but it falls to the thermal response as $\tau \to \infty$, as discussed further in section 5.4.3 and illustrared in figures 5.6, 5.7 and 5.8. This clarifies and adds to the discussion given in [100].

We wish to compare these particle detector results in $\mathbb{RP}^3$ de Sitter space to the asociated GEMS particle detector. We see from (5.243) that the GEMS worldline of interest is a Rindler trajectory with acceleration $a = 1/\alpha$ in the 5-dimensional embedding space. Therefore in the 5-dimensional Minkowski embedding space of de Sitter space we see that the response of a detector following this worldline is a thermal one with associated temperature $T = 1/(2\pi\alpha)$ and so we expect, as indeed we saw in section 5.9.1, the response of the detector in de Sitter space to also be a thermal one with temperature $T = 1/(2\pi\alpha)$. Again the actual responses of detectors in the two situations are not identical (as seen in sections 5.3.1 and 5.9.1) due to the different dimensions of the spaces, the most obvious difference being the presence of the Planckian factor in the de Sitter response and the Fermi factor in Rindler response on the odd dimensional embedding space. As the $\mathbb{RP}^3$ de Sitter spacetime is built as a quotient of de Sitter space under the map $J : (z_0, z_1, z_2, z_3, z_4) \mapsto (z_0, -z_1, -z_2, -z_3, -z_4)$ we have immediately the action of this map on the embedding space. The geodesic worldline of interest maps to a Rindler worldline in this embedding space with acceleration $a = 1/\alpha$, so in the GEMS approach we consider a Rindler particle detector with this acceleration in this 5-dimensional embedding space. The transition rate was found in section 5.4.4. The thermal part of the transition rate is constant in time and is given by

$$\dot{F}_{M\tau}(\omega) = \frac{a^2}{8\pi(e^{\frac{2\pi\omega}{a}}+1)}(1/4+\omega^2/a^2) \ , \qquad (5.246)$$

corresponding to the temperature $T = \frac{a}{2\pi}$. The image part of the transition rate depends on the proper time and is given by

$$\dot{F}_{I\tau}(\omega) = \frac{1}{4\pi^2}\int_0^\infty ds \, \frac{\cos(\omega s)}{\left(4/a^2 \cosh^2\left(\frac{a(2\tau-s)}{2}\right)\right)^{3/2}} \ . \qquad (5.247)$$

The total response is shown for various values of the parameters in figure 5.10. The qualitative similarities to the $\mathbb{RP}^3$ de Sitter transition rate are apparent. They provide evidence that, at least in some cases, the GEMS procedure may be applied to quotient spaces such as $\mathbb{RP}^3$ de Sitter space and the $\mathbb{RP}^3$ geon where the embedding spaces are Minkowski spaces with suitable identifications.



Figure 5.10: Transition rate for a detector uniformly accelerated in the z_1 direction on the quotient space of 5-dimensional Minkowski space under the involution J. The parameters are $\alpha = 1$, $C = (2z_1)^2 + (2z_2)^2 + (2z_3)^2 = 0$ and $\omega = 1$ (upper curve), $\omega = 1.5$ and $\omega = 2$ (lower curve).

Chapter 6

Conclusions

This thesis has been concerned with quantum field theory on topologically non-trivial manifolds and with particle detector models. Chapters 2 and 3 discussed thermal effects for the free Dirac field on the $\mathbb{RP}^3$ geon and on a topologically analogous flat spacetime M_- via a Bogolubov transformation analysis. Compared with the scalar field [22], the main new issue with fermions is that the spacetimes admit two inequivalent spin structures, and there are hence two inequivalent Hartle-Hawking like vacua on the geon and two inequivalent Minkowski-like vacua on M_-. We showed that an observer in the exterior region of the geon can detect both the nonthermality of the Hartle-Hawking like state and the spin structure of this state by suitable interference measurements, and similar results hold for a Rindler observer on M_-. When probed with suitably restricted operators, such as operators at asymptotically late Schwarzschild (respectively Rindler) times, these states nevertheless appear thermal in the usual Hawking (Unruh) temperature, for the same geometric reasons as in the scalar case [22]. As a by-product of the analysis, we presented the Bogolubov transformation for the Unruh effect for the massive Dirac field in $(3+1)$-dimensional Minkowski space, complementing and correcting the previous literature.

As a late time observer in the geon exterior sees a thermal state in the usual Hawking temperature, the classical laws of black hole mechanics lead the observer to assign to the geon the same entropy as to a conventional Schwarzschild hole with the same mass. It was found in [22] that an attempt to evaluate the geon entropy by path-integral methods leads to half of the Bekenstein-Hawking entropy of a Schwarzschild hole of the same mass, and it was suggested that state-counting computations of the geon entropy could shed light on this discrepancy. Our work says little of what the full framework of such a computation would be, but our work would presumably provide part of the fermionic machinery in the computation. In particular, the issue of the spin structure would need to be faced seriously: Does an entropy computation by state-counting need to count the two spin structures as independent degrees of freedom?

In chapter 4 we computed the stress-energy expectation value on M_0 and M_- in Minkowski-like vacuum states, for both massive scalar and spinor fields. For the scalar field our results on M_0 agree with those in [46]. For the spinor field on M_0 they agree with [48]. On M_- the results for the massive fields are new. Further in the massless limit our expectation values agree with the previous literature [22, 36, 47]. We presented also the calculation for 2-component massless spinors. It was shown on both M_0 and M_- that the stress tensor is independent of the handedness of the spinor. In all cases the values fall off exponentially in the large m limit, and the leading order correction for small mass is $O(m^2)$. Further it is noted that for the scalar field in the large a limit on M_- there is an exponential decay in the massive case while for the massless field the behaviour is a^{-4}. For the massive field the difference between $\langle 0|T_{\mu\nu}|0\rangle$ on M_- and the corresponding values on M_0 falls off exponentially in the limit of large $r^2 := x^2 + y^2$, while for the massless field it behaves as $O(r^{-3})$. It is seen that for spinors the two spin structures are distinguished by the sign of a non-vanishing shear component.

Our underlying interest in the expectation values arises from the role of M_0 and M_- in modelling, in the context of accelerated observers on flat spacetimes, the Hawking(-Unruh) effect on respectively the Kruskal manifold and the $\mathbb{RP}^3$ geon. As we have seen in chapters 2, 3 and 5 certain aspects of the thermal and non-thermal effects for scalar and spinor fields on M_0 and M_- are at present understood from the viewpoint of Bogolubov transformations and particle detector analyses (see also [22]), but the connections between (non-)thermality and stress-energy remain less clear. We view our results in chapter 4, in conjunction with those in chapters 2, 3, 5 and in [22], as data points to which we anticipate future work on this question to provide a deeper understanding.

In chapter 5 we considered particle detector models in the context of quantum field theory in curved spacetime. In particular we investigated the model of Schlicht [25, 26]. The model is that of a monopole detector linearly coupled to a massless scalar field which is smeared with a window function in order to regularize the correlation function in the transition rate. We extended the regularization of the correlation function for the massless linearly coupled scalar field to d-dimensional Minkowski space, and we showed that it leads to the expected responses for inertial and uniformly accelerated detectors switched on in the infinite past and off at $\tau < \infty$. Further we extended the regularization of Schlicht to the massive scalar field in Minkowski space and have shown that it reduces to that of [25, 26] in the massless limit. Next we introduced a model of a linearly coupled massless scalar field detector on spacetimes built as quotients of Minkowski space under certain discrete isometries. In a number of cases the model, when switched on at $\tau_0 = -\infty$ and read at $\tau < \infty$, was shown to reproduce the known asymptotic responses. These cases include the uniformly accelerated detector on M_0 [22, 107] as well as the inertial and uniformly accelerated detectors on Minkowski space with boundary when the motion is parallel to the boundary [107]. These results suggest that our model is reasonable. Further we presented a number of new responses, the most interesting of which are the time dependent responses on Minkowski space with boundary and on M_-. An inertial detector approaching the boundary on Minkowski space with boundary was considered and seen to react in a qualitatively similar way to one travelling parallel to the boundary but taking progressively smaller distances (ie comparing figures 5.2 and 5.4). The main difference is that in the detector approaching the boundary a divergence in the transition rate occurs as the boundary is reached. A detector with uniform acceleration perpendicular to the boundary was also considered and an interesting observation made. For a detector which is switched on in the infinite past the transition rate is found to oscillate in τ at late times with period π/ω, never tending to the Minkowski thermal response, no matter how far from the boundary the detector gets in the future. However for a detector switched on at a finite time (that is $\tau_0 > -\infty$), the response will at late times tend to the thermal Minkowski response. For instantaneous, exponential and Gaussian switching functions the conclusion is the same. Responses were also considered on the quotient spaces of Minkowski space under the involution $J_{c_k} : (t, x_1, x_2, \ldots x_{d-1}) \mapsto (t, -x_1, -x_2, - \ldots - x_k, x_{k+1}, \ldots, x_{d-1})$ where $1 < k < d$ and certain relations to the responses on Minkowski space with boundary noted. The responses of uniformly accelerated detectors, where the motion is in the x_1 direction, are relevant for the discussion of static detectors in the $\mathbb{RP}^3$ geon exterior as well as comoving detectors in $\mathbb{RP}^3$ de Sitter via their global embedding Minkowski spacetimes (GEMS). Next we extended the detector model and regularization to the nonlinearly coupled scalar field and to the massless Dirac field. With a few minor technicalities the extension is quite straightforward. In the case of the Dirac field again we obtained the transition rate and the power spectrum of the Dirac noise for a detector switched on in the infinite past on inertial and a uniformly accelerated trajectories. The power spectrum for the accelerated detector agrees with the previous literature (see e.g [31]) and so suggests our model is reasonable. Further we briefly considered the response of the Dirac detector on M_0 and M_-. One aim was to see whether a uniformly accelerated Dirac detector on M_- could distinguish the two spin structures there. Unfortunately we have shown that this is not the case for our particular model.

In section 5.7 we considered the response of a static detector in the exterior region of the $\mathbb{RP}^3$ geon via a global embedding Minkowski space. Although the GEMS program has so far only been applied in a kinematical setting, our aim was to examine the possibility that the response of the detector in the embedding space is related to that in the underlying curved space. We found that the response is related to that of uniformly accelerated detectors given in section 5.4.4. In particular it is shown in the embedding space, and expected on the geon, that the response is not thermal, in the sense that it does not satisfy the KMS condition, for most times. Further it is seen that for a detector switched on in the infinite past the response is approximately thermal at early times but does not return to the thermal response at late times in contrast to expectations (see e.g [22] and sections 2 and 3 here). If the detector however is turned on at some finite time in the distant past then the response is approximately thermal when turned on and returns to being approximately thermal in the distant future.

Lastly we extended our model to conformally flat spacetimes and considered some responses on de Sitter space and $\mathbb{RP}^3$ de Sitter space. The regularization of [25] is not easily adaptable to general motions in conformally flat spacetimes, due to the possibility of spatially closed hypersurfaces in the detector's rest frame. We argued however for a similar regularization by reinterpreting the regularization as an ultraviolet cut-off in the high "frequency" modes. On de Sitter space the transition rate for a detector switched on in the infinite past is found for a uniformly accelerated detector, and it is found to agree with previous literature [116]. This result suggests our regularization is reasonable. In the case of an inertial detector in de Sitter space the regularization we introduced is shown to also come from the consideration of a detector with spatial extension where the detector is rigid in its rest frame. On $\mathbb{RP}^3$ de Sitter space the response of a detector following the comoving worldline was considered in two ways. Firstly a direct calculation showed that the response is exactly that of a uniformly accelerated detector approaching the boundary on Minkowski space with boundary (identifying $1/\alpha$ with the acceleration). Again therefore the response of a detector switched on in the infinite past has an oscillatory behaviour in the distant future and does not tend to the expected thermal result, the image term breaking the KMS condition, in contrast to what was expected [100]. The response of a detector switched on at a finite time does however tend to the expected thermal response at late times. Secondly we considered the same calculation from the GEMS perspective. Although the response is clearly different to that in the original space, due to the different dimensions of $\mathbb{RP}^3$ de Sitter space and the embedding space, it is seen that the response is qualitatively very similar. The calculation therefore provides a good example for investigating detector responses in curved spacetimes and those in their GEMS, and suggests also that the response found in the GEMS of the $\mathbb{RP}^3$ geon in section 5.7 is indeed closely related to the response of the static detector in the geon itself.

Future Work

There are many possible directions for future work on the topics discussed in this thesis. On the $\mathbb{RP}^3$ geon for instance one interesting subject which we have not really discussed here is its entropy. In [22] it is argued from the observed temperature and the classical laws of black hole thermodynamics that an observer at late Schwarzschild time in the exterior will assign an entropy to the hole of 1/4 its late time horizon area, which is the entropy of a Kruskal hole with the same mass. However it was also shown that from a path integral approach the entropy assigned to the hole is 1/8 the late time horizon area. The reason for the difference is not clear. It would be very interesting therefore to consider a state counting argument for the entropy of the geon in the context of a quantum gravity model such as loop quantum gravity, as was done for large non-rotating black holes in [128].

Further work for the particle detector models discussed could include a thorough investigation of the dependency of Schlicht's model on the window function (detector shape). Also it would be interesting to investigate in depth the most general case of an arbitrary trajectory in a general curved spacetime. The GEMS calculations discussed here are also worthy of further investigation. In particular, to what extent do the responses in the GEMS relate to the responses in the original curved spaces? We know that the GEMS procedure may not be applied reliably to the response of an arbitrary non-stationary trajectory in an arbitrary spacetime [125] but we expect for some classes of trajectories and spacetimes the GEMS approach is useful.

Finally, as mentioned when discussing our stress energy results, the relation between the (non)-thermal behaviour seen in various black hole (and flat) spacetimes and the stress energy expectation values is not clear and a deeper understanding is required.

Appendix A

Analytic continuation of Rindler modes across the horizons

In this appendix we provide more details on the analytic continuation of the Rindler modes (2.33) on M_0 into the future and left hand wedges.

(2.33) are Rindler modes in R_0 expressed with respect to the globally defined Minkowski vierbein. We build from them a complete set of positive frequency Minkowski modes (i.e. positive frequency with respect to ∂_t) by continuing them (with $\Omega \in \mathbb{R}$) across the horizons into F and L in the lower half complex t plane. (2.33) are functions of $t-x$ and $t+x$ where in R

$$\begin{aligned} \xi &= (x^2-t^2)^{1/2} = ((x-t)(x+t))^{1/2} \ , && \text{(A.1)} \\ \tau &= \operatorname{arctanh}\left(\frac{t}{x}\right) = \frac{1}{2}\ln\left(\frac{x+t}{x-t}\right) \ , && \text{(A.2)} \end{aligned}$$

Continuing these across the future horizon $t=x$, from $(t-x)<0$ to $(t-x)>0$ we note that $t=x$ is a branch point. In order to analytically continue into the lower half complex t plane we must make the substitutions $\sqrt{x-t} \to i\sqrt{t-x}$ and $\ln(x-t) \to \ln(t-x) + i\pi$.

This gives

$$\begin{aligned}\xi &= (x^2-t^2)^{1/2} \to i((t-x)(x+t))^{1/2} = i\xi \ , &\text{(A.3)}\\ \tau &= \operatorname{arctanh}\left(\frac{t}{x}\right) \to \frac{1}{2}\ln\left(\frac{x+t}{t-x}\right) - \frac{i\pi}{2} = \tau - \frac{i\pi}{2} \ , &\text{(A.4)}\end{aligned}$$

where on the left hand side of these expressions (τ, ξ) are in R and on the right (τ, ξ) are in F as defined by (2.49). Under this substitution it is easy to check that the Rindler modes in R (2.33) become (2.45) and (2.46) in F.

The continuation further to the L wedge is similar. We must cross the horizon $t = -x$ which again is a branch point in the expressions for (τ, ξ) in F. Crossing from $(x+t) > 0$ to $(x+t) < 0$ in the lower half t-plane we make the substitutions $\sqrt{x+t} \to -i\sqrt{-(x+t)}$ and $\ln(x+t) \to \ln(-(x+t)) - i\pi$. Then

$$\begin{aligned}\xi &= (t^2-x^2)^{1/2} \to -i((x-t)(x+t))^{1/2} = -i\xi \ , &\text{(A.5)}\\ \tau &= \operatorname{arctanh}\left(\frac{x}{t}\right) \to \frac{1}{2}\ln\left(\frac{x+t}{x-t}\right) - \frac{i\pi}{2} = \tau - \frac{i\pi}{2} \ , &\text{(A.6)}\end{aligned}$$

where the expressions on the left are in F and those on the right are in L with (τ, ξ) defined by (2.35). Under this continuation the modes (2.45) and (2.46) in F become (2.50).

A complete set of negative frequency Minkowski modes can be obtained in a similar manner by continuing the Rindler modes in R (2.33), with $\Omega \in \mathbb{R}$, across the horizons in the upper half complex t-plane. Alternatively we may just take the charge conjugates of our positive frequency modes.

Appendix B

Evaluating integrals using hyperspherical coordinates

In this appendix we show how to evaluate integrals of the form

$$\int d^{d-1}k\, F(|\mathbf{k}|)e^{i\mathbf{k}\cdot\mathbf{R}}e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} \;, \tag{B.1}$$

where $d > 2$, the integration is over $\mathbf{k} \in \mathbb{R}^{d-1}$, $\mathbf{R}$ and $\tilde{\mathbf{R}}$ are vectors in $\mathbb{R}^{d-1}$ and $F(|\mathbf{k}|)$ is an arbitrary function.

First we transform to hyperspherical coordinates (polar and spherical coordinates for $d = 3$ and $d = 4$ respectively) in $\mathbf{k}$ space $(|\mathbf{k}|, \theta_1, \ldots, \theta_{d-2})$, where

$$\begin{aligned}
k_1 &= |\mathbf{k}| \sin\theta_1 \ldots \sin\theta_{d-3} \sin\theta_{d-2} \;, \\
k_2 &= |\mathbf{k}| \sin\theta_1 \ldots \sin\theta_{d-3} \cos\theta_{d-2} \;, \\
k_3 &= |\mathbf{k}| \sin\theta_1 \ldots \sin\theta_{d-4} \cos\theta_{d-3} \;, \\
&\vdots \\
k_{d-1} &= |\mathbf{k}| \cos\theta_1 \;,
\end{aligned} \tag{B.2}$$

with $0 < |\mathbf{k}| < \infty$, $0 < \theta_i < \pi$ for $i = 1, \ldots, d-3$ and $0 < \theta_{d-2} < 2\pi$. Then

$$\int d^{d-1}k\, F(|\mathbf{k}|) e^{i\mathbf{k}\cdot\mathbf{R}} e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} = \int_0^\infty d|\mathbf{k}|\, |\mathbf{k}|^{d-2} F(|\mathbf{k}|) \int d^{d-2}\Omega\, e^{i\mathbf{k}\cdot\mathbf{R}} e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} \ . \tag{B.3}$$

The angular integral may now be performed by choosing a suitable set of axes in $\mathbf{k}$ space. In particular we choose the k_{d-1} axis in the direction of $\mathbf{R}$ and the k_{d-2} axis such that $\tilde{\mathbf{R}}$ lies in the (k_{d-2}, k_{d-1})-plane. Then $\mathbf{k}\cdot\mathbf{R} = |\mathbf{k}||\mathbf{R}|\cos\theta_1$ and $\mathbf{k}\cdot\tilde{\mathbf{R}} = |\mathbf{k}||\tilde{\mathbf{R}}|(\cos\theta_1\cos\psi + \sin\theta_1\cos\theta_2\sin\psi)$, where ψ is the angle between $\mathbf{R}$ and $\tilde{\mathbf{R}}$, i.e. $|\mathbf{R}||\tilde{\mathbf{R}}|\cos\psi = \mathbf{R}\cdot\tilde{\mathbf{R}}$. Considering first $d > 4$ we have[1]

$$\begin{aligned} \int d^{d-2}\Omega\, e^{i\mathbf{k}\cdot\mathbf{R}} e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} &= \int_0^{2\pi} d\theta_{d-2} \prod_{i=3}^{d-3} \int_0^\pi d\theta_i\, (\sin\theta_i)^{d-2-i} \int_0^\pi d\theta_1\, (\sin\theta_1)^{d-3} e^{i|\mathbf{k}||\mathbf{R}|\cos\theta_1} \\ &\times \int_0^\pi d\theta_2\, (\sin\theta_2)^{d-4} e^{\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|(\cos\theta_1\cos\psi + \sin\theta_1\cos\theta_2\sin\psi)} \ . \end{aligned} \tag{B.4}$$

From [50] we have

$$\int_0^\pi d\theta\, (\sin\theta)^{2\nu} e^{\pm a\cos\theta} = \pi^{1/2}\left(\frac{2}{a}\right)^\nu \Gamma[\nu + 1/2] I_\nu(a) \ , \tag{B.5}$$

for $Re(\nu) > -1/2$, where $I_\nu(a)$ is a Bessel function, and so

$$\begin{aligned} \int_0^\pi d\theta_2\, (\sin\theta_2)^{d-4} e^{\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\sin\theta_1\cos\theta_2\sin\psi} &= \pi^{1/2}\left(\frac{2}{\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\sin\theta_1\sin\psi}\right)^{\frac{d-4}{2}} \Gamma[(d-3)/2] \\ &\times I_{\frac{d-4}{2}}(\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\sin\theta_1\sin\psi) \ . \end{aligned} \tag{B.6}$$

The integral over $\theta_3 \ldots \theta_{d-2}$ in (B.4) may be performed by noting that it is the surface area of a unit $(d-4)$-dimensional sphere (in topologists notation), which is given by

$$S_{d-4} = 2\frac{\pi^{(d-3)/2}}{\Gamma[(d-3)/2]} \ , \tag{B.7}$$

---

[1] For $d = 5$ the expression is (B.4) without the product term.

see e.g [40]. Thus we have

$$\begin{aligned}\int d^{d-2}\Omega\, e^{i\mathbf{k}\cdot\mathbf{R}}e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} &= \frac{(2\pi)^{d/2-1}}{(\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\sin\psi)^{(d-4)/2}}\int_0^\pi d\theta_1\,(\sin\theta_1)^{d/2-1}e^{(i|\mathbf{k}||\mathbf{R}|+\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\cos\psi)\cos\theta_1}\\ &\quad\times I_{\frac{d-4}{2}}(\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\sin\theta_1\sin\psi)\ .\end{aligned}\tag{B.8}$$

The integral over θ_1 may be done by expanding the Bessel function as a series and doing the integral over θ_1 for each term in the sum. We have

$$\begin{aligned}\int_0^\pi d\theta\,(\sin\theta)^{\nu+1}e^{a\cos\theta}I_\nu(b\sin\theta) &= \sum_{k=0}^\infty\left(\frac{b}{2}\right)^{\nu+2k}\frac{1}{k!\Gamma[\nu+k+1]}\int_0^\pi d\theta\,(\sin\theta)^{2\nu+2k+1}e^{a\cos\theta}\ ,\\ &= (2\pi)^{1/2}\sum_{k=0}^\infty\frac{b^{\nu+2k}}{k!2^k}a^{-(\nu+k+1/2)}I_{\nu+k+1/2}(a)\ ,\\ &= (2\pi)^{1/2}b^\nu\sum_{k=0}^\infty\frac{b^{2k}}{k!}\left(\frac{d}{dq}\right)^k\left(q^{-\frac{1}{2}(\nu+\frac{1}{2})}I_{\nu+1/2}(\sqrt{q})\right)\ ,\end{aligned}\tag{B.9}$$

where $q=a^2$, and the second line follows from (B.5). Recognizing the sum in the last expression of (B.9) as a Taylor series we thus have

$$\int_0^\pi d\theta\,(\sin\theta)^{\nu+1}e^{a\cos\theta}I_\nu(b\sin\theta) = (2\pi)^{1/2}\frac{b^\nu}{(\sqrt{a^2+b^2})^{\nu+1/2}}I_{\nu+1/2}(\sqrt{a^2+b^2})\ .\tag{B.10}$$

Further we find

$$-(i|\mathbf{R}|+\epsilon|\tilde{\mathbf{R}}|\cos\psi)^2-(\epsilon|\tilde{\mathbf{R}}|\sin\psi)^2 = (\mathbf{R}-i\epsilon\tilde{\mathbf{R}})\cdot(\mathbf{R}-i\epsilon\tilde{\mathbf{R}}) = (\mathbf{R}-i\epsilon\tilde{\mathbf{R}})^2\ .\tag{B.11}$$

Collecting these results together we have

$$\int d^{d-1}k\,F(|\mathbf{k}|)e^{i\mathbf{k}\cdot\mathbf{R}}e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} = (2\pi)^{(d-1)/2}\int_0^\infty d|\mathbf{k}|\,|\mathbf{k}|^{d-2}F(|\mathbf{k}|)\frac{J_{(d-3)/2}\left(|\mathbf{k}|\sqrt{(\mathbf{R}-i\epsilon\tilde{\mathbf{R}})^2}\right)}{\left(|\mathbf{k}|\sqrt{(\mathbf{R}-i\epsilon\tilde{\mathbf{R}})^2}\right)^{(d-3)/2}}\ ,\tag{B.12}$$

where we have replaced the I Bessel in (B.10) by a J Bessel using $I_\nu(z)=e^{-\frac{\nu\pi i}{2}}J_\nu(ze^{\frac{\pi i}{2}})$.

For $d = 4$ the expression analogous to (B.4) for the angular integral is

$$\begin{aligned}\int d^2\Omega\, e^{i\mathbf{k}\cdot\mathbf{R}}e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} &= \int_0^{\pi} d\theta_1\, \sin\theta_1 e^{i|\mathbf{k}||\mathbf{R}|\cos\theta_1} \\ &\quad\times \int_0^{2\pi} d\theta_2\, e^{\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|(\cos\theta_1\cos\psi+\sin\theta_1\cos\theta_2\sin\psi)} \\ &= 2\pi\int_0^{\pi} d\theta_1\, \sin\theta_1 e^{i|\mathbf{k}||\mathbf{R}|\cos\theta_1+\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\cos\theta_1\cos\psi} I_0(\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|\sin\theta_1\sin\psi)\ ,\end{aligned} \tag{B.13}$$

where we have used (B.5) for the integral over θ_2. Expression (B.13) agrees with (B.8) when $d = 4$ and thus we are led again to the result (B.12).

For $d = 3$ we take the k_2 axis in the direction of $\mathbf{R}$ in (B.3) then $\mathbf{k}\cdot\mathbf{R} = |\mathbf{k}||\mathbf{R}|\cos\theta_1$, $\mathbf{k}\cdot\tilde{\mathbf{R}} = |\mathbf{k}||\tilde{\mathbf{R}}|(\cos\theta_1\cos\psi + \sin\theta_1\sin\psi)$ and

$$\int d\Omega\, e^{i\mathbf{k}\cdot\mathbf{R}}e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} = \int_0^{2\pi} d\theta_1 e^{i|\mathbf{k}||\mathbf{R}|\cos\theta_1} e^{\epsilon|\mathbf{k}||\tilde{\mathbf{R}}|(\cos\theta_1\cos\psi+\sin\theta_1\sin\psi)}\ . \tag{B.14}$$

Hence, [50],

$$\int d^2k\, F(|\mathbf{k}|)e^{i\mathbf{k}\cdot\mathbf{R}}e^{\epsilon\mathbf{k}\cdot\tilde{\mathbf{R}}} = 2\pi\int_0^{\infty} d|\mathbf{k}|\, |\mathbf{k}|F(|\mathbf{k}|)J_0\left(|\mathbf{k}|\sqrt{(\mathbf{R}-i\epsilon\tilde{\mathbf{R}})^2}\right)\ , \tag{B.15}$$

which also agrees with (B.12). Thus (B.12) is the expression we require for all $d > 2$.

In the case of equation (5.28) we want

$$\begin{aligned} g_\epsilon(\mathbf{k};\tau) &= \frac{\epsilon\Gamma(d/2)}{\pi^{d/2}} e^{-i(k\cdot x(\tau))}\int d^{d-1}\xi\, \frac{1}{(\xi^2+\epsilon^2)^{d/2}} e^{-i\bar{\mathbf{k}}\cdot\xi}\ , \\ &= \frac{\epsilon\Gamma(d/2)}{\pi^{d/2}} e^{-i(k\cdot x(\tau))}\frac{(2\pi)^{(d-1)/2}}{|\bar{\mathbf{k}}|^{(d-3)/2}}\int_0^{\infty} d|\xi|\, \frac{|\xi||\xi|^{(d-3)/2}}{(\xi^2+\epsilon^2)^{d/2}} J_{(d-3)/2}(|\xi||\bar{\mathbf{k}}|)\ ,\end{aligned} \tag{B.16}$$

where the second line follows from (B.12). Again the integral may be found in integral tables (see [106] p488 no.11.4.44)

$$\int_0^\infty d|\xi| \, \frac{|\xi||\xi|^{(d-3)/2}}{(\xi^2+\epsilon^2)^{d/2}} J_{(d-3)/2}(|\xi||\bar{\mathbf{k}}|) = \frac{|\bar{\mathbf{k}}|^{(d/2-1)}\epsilon^{-1/2}}{2^{(d/2-1)}\Gamma(d/2)} K_{1/2}(|\bar{\mathbf{k}}|\epsilon) \ , \tag{B.17}$$

K is a modified Bessel function and $K_{1/2}(z) = (\pi/(2|\bar{\mathbf{k}}|\epsilon))^{1/2} e^{-|\bar{\mathbf{k}}|\epsilon}$ so finally we find

$$g_\epsilon(\mathbf{k};\tau) = e^{-i(k\cdot x(\tau))} e^{-\epsilon|\bar{\mathbf{k}}|} \ . \tag{B.18}$$

For equation (5.33) we find

$$\begin{aligned} \langle 0|\phi(\tau)\phi(\tau')|0\rangle &= \frac{1}{(2\pi)^{d-1}} \int \frac{d^{d-1}k}{2|\mathbf{k}|} \, e^{-i|\mathbf{k}|(t-t'-i\epsilon(\dot{t}+\dot{t}'))+i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))} \ , \\ &= \frac{1}{2(2\pi)^{(d-1)/2}} \int_0^\infty d|\mathbf{k}| \, |\mathbf{k}|^{d-3} e^{-i|\mathbf{k}|(t-t'-i\epsilon(\dot{t}+\dot{t}'))} \\ &\quad \times \frac{J_{(d-3)/2}\left(|\mathbf{k}|\sqrt{(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))^2}\right)}{\left(|\mathbf{k}|\sqrt{(\mathbf{x}-\mathbf{x}'-i\epsilon(\dot{\mathbf{x}}+\dot{\mathbf{x}}'))^2}\right)^{(d-3)/2}} \ . \end{aligned} \tag{B.19}$$

The $|\mathbf{k}|$ integral may be found in [50], using

$$\int_0^\infty dx \, e^{-\alpha x} x^\nu J_\nu(\beta x) = \frac{(2\beta)^\nu \Gamma(\nu+1/2)}{\pi^{1/2}(\alpha^2+\beta^2)^{\nu+1/2}} \ , \tag{B.20}$$

for $Re(\alpha) > 0$, the result is (5.34). The same integrations are required in finding the usual correlation function (5.15) where ϵ is taken to 0 in (B.19) and then a cut-off $e^{-\epsilon|\mathbf{k}|}$ is introduced.

For equation (5.69) we want

$$\begin{aligned} g_\epsilon(\mathbf{k};\tau) &= \frac{1}{\epsilon^{d-1}(2\pi)^{(d-1)/2}} e^{-i(k\cdot x(\tau))} \int d^{d-1}\xi \, e^{-\frac{\xi^2}{2\epsilon^2}} e^{-i\bar{\mathbf{k}}\cdot\xi} \ , \\ &= \frac{1}{\epsilon^{d-1}|\bar{\mathbf{k}}|^{(d-3)/2}} e^{-i(k\cdot x(\tau))} \int_0^\infty d|\xi| \, |\xi||\xi|^{(d-3)/2} e^{-\frac{|\xi|^2}{2\epsilon^2}} J_{(d-3)/2}(|\bar{\mathbf{k}}||\xi|) \ . \end{aligned} \tag{B.21}$$

Again from [106] (p486 no.11.4.29) we have

$$\int_0^\infty dt\, e^{-\alpha^2 t^2} t^{\nu+1} J_\nu(bt) = \frac{b^\nu}{(2\alpha^2)^{\nu+1}} e^{-\frac{b^2}{4\alpha^2}} \ , \tag{B.22}$$

and so we find

$$g_\epsilon(\mathbf{k};\tau) = e^{-i(k\cdot x(\tau))} e^{-\frac{\epsilon^2 |\bar{\mathbf{k}}|^2}{2}} \ . \tag{B.23}$$

Appendix C

Expressions in 2 dimensions

In this appendix we present the calculation of some of the expressions appearing in chapter 5 in 2 dimensions.

C.1 The correlation function

In 2-dimensional Minkowski space the solutions to the massless Klein Gordon equation may be expressed as the mode expansion

$$\phi(t,x) = \int_{-\infty}^{\infty} \frac{dk}{(4|k|\pi)^{1/2}} \left(a(k) e^{-i(|k|t-kx)} + a^{\dagger}(k) e^{i(t-kx)} \right) . \tag{C.1}$$

The Wightman function is then

$$\langle 0|\phi(x)\phi(x')|0\rangle = \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{dk}{|k|} e^{-i(|k|(t-t')-k(x-x'))} . \tag{C.2}$$

We see immediately from (C.1) and (C.2) that there is an infrared divergence due to the $k=0$ mode (as well as the usual ultraviolet divergences found in all dimensions). In calculating the Wightman function (C.2) we could introduce an infrared cut-off in order to regularize this divergence, by for example considering

$$\langle 0|\phi(x)\phi(x')|0\rangle = \frac{1}{4\pi}\int_{-\infty}^{\infty}\frac{dk}{|k|}\left(e^{-i(|k|(t-t')-k(x-x'))} - \Theta(C-|k|)\right) \ , \tag{C.3}$$

where $\Theta(x)$ is a step function and C a positive constant (see e.g. [129] for more details). However as we shall see later the divergence does not contribute to the detector response provided we include a temporal window function and consider an asymptotic (infinite time) detection. Without the window function the contribution from these terms may be easily found and so we do not regularize it here but simply drop the divergent term in any calculations done. The ultraviolet divergence in (C.2) still requires regularization. In the usual $i\epsilon$-prescription

$$\begin{aligned}\langle 0|\phi(x)\phi(x')|0\rangle &= \frac{1}{4\pi}\int_{-\infty}^{\infty}\frac{dk}{|k|}\,e^{-i(|k|(t-t'-i\epsilon)-k(x-x'))} \ , \\ &= \frac{1}{4\pi}\left(\int_0^{\infty}\frac{dk}{k}\,e^{-ik((t-t'-i\epsilon)-(x-x'))} + \int_0^{\infty}\frac{dk}{k}\,e^{-ik((t-t'-i\epsilon)+(x-x'))}\right) \ .\end{aligned} \tag{C.4}$$

The integrals may be done by parts

$$\begin{aligned}\int_0^{\infty}\frac{dk}{k}\,e^{-i\mu k} &= \left[\ln(k)e^{-i\mu k}\right]_0^{\infty} + i\mu\int_0^{\infty}dk\,\ln(k)e^{-i\mu k} \ , \\ &= -\ln(0) - (\mathbf{C} + \ln(i\mu)) \ ,\end{aligned} \tag{C.5}$$

when $Im(\mu) < 0$ [50], where $\mathbf{C}$ is Euler's constant. So

$$\begin{aligned}\langle 0|\phi(x)\phi(x')|0\rangle &= \frac{1}{4\pi}(-2\ln(0) - 2\mathbf{C} - \ln(i[(t-t'-i\epsilon)-(x-x')]) \\ &\quad - \ln(i[(t-t'-i\epsilon)+(x-x')])) \\ &= -\frac{1}{4\pi}\ln((t-t'-i\epsilon)^2-(x-x')^2) + C \ ,\end{aligned} \tag{C.6}$$

where C is an ill-defined constant. The usual approach is to say that derivatives of the correlation function are well defined and are given by the derivatives of (C.6). We will show later however that in our particle detector model we may work with (C.6) in cases where the constant term does not contribute to the response.

Next consider the correlation function for the smeared field in two dimensions. The discussion follows as in sections 5.3.1. For a Lorentzian shape function

$$g_\epsilon(\mathbf{k};\tau) = \frac{\epsilon}{\pi} e^{-i(k\cdot x(\tau))} \int_{-\infty}^{\infty} d\xi \, \frac{e^{-i\bar{k}\xi}}{(\xi^2+\epsilon^2)} \ , \tag{C.7}$$

where $\bar{k} := k \cdot e_\xi(\tau)$. Now

$$\int_{-\infty}^{\infty} dx \, \frac{e^{-ipx}}{(x^2+a^2)} = \frac{\pi}{a} e^{-|ap|} \ , \tag{C.8}$$

for $a \neq 0$ and p real [50], so

$$g_\epsilon(\mathbf{k};\tau) = e^{-i(k\cdot x(\tau))} e^{-\epsilon|k\cdot e_\xi(\tau)|} = e^{-i(|k|t(\tau)-kx(\tau))} e^{-\epsilon(|k|\dot{t}-k\dot{x})} \ . \tag{C.9}$$

Then

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{dk}{|k|} \, e^{-i(|k|(t-t'-i\epsilon(\dot{t}+\dot{t}'))-k(x-x'-i\epsilon(\dot{x}+\dot{x}')))} \ . \tag{C.10}$$

An analogous calculation to that leading to (C.6), noting that on a timelike worldline $Im[k(t-t'-i\epsilon(\dot{t}+\dot{t}')) \pm k(x-x'-i\epsilon(\dot{x}+\dot{x}'))] < 0$, gives

$$\langle 0|\phi(\tau)\phi(\tau')|0\rangle = -\frac{1}{4\pi} \ln((t-t'-i\epsilon(\dot{t}+\dot{t}'))^2 - (x-x'-i\epsilon(\dot{x}+\dot{x}'))^2) + C \ , \tag{C.11}$$

where C is an ill-defined constant.

C.2 Inertial detector

Next we consider an inertial detector in 2 dimensions, firstly for a finite time detector switched on instantaneously. From (5.12) and (C.11) the transition rate is

$$\dot{F}_{\Delta\tau}(\omega) = -\frac{1}{4\pi}\int_{-\Delta\tau}^{\Delta\tau} ds\, e^{-i\omega s}\left(\ln((s-2i\epsilon)^2)+C\right) \ , \tag{C.12}$$

where $\Delta\tau = \tau - \tau_0$. The contribution of the constant is

$$\begin{aligned}\dot{F}_{\Delta\tau}(\omega) &= C\int_{-\Delta\tau}^{\Delta\tau} ds\, e^{-i\omega s} \ , \\ &= \frac{2C}{\omega}\sin(\omega(\Delta\tau)) \ .\end{aligned} \tag{C.13}$$

Thus for instant switching constant terms in the Wightman function do contribute to the transition rate, even in the limit as $(\tau-\tau_0)\to\infty$.

Consider then the transition rate for a detector switched smoothly with an exponential switching function.

$$\begin{aligned}\dot{F}_{\Delta\tau}(\omega) &= C\int_{-\infty}^{\infty} ds\, e^{-i\omega s-\frac{|s|}{\Delta\tau}} \ , \\ &= C\left(\int_{0}^{\infty} ds\, e^{-i\omega s-\frac{s}{\Delta\tau}}+\int_{0}^{\infty} ds\, e^{i\omega s-\frac{s}{\Delta\tau}}\right) \ , \\ &= \frac{2C\Delta\tau}{(1+\Delta\tau^2\omega^2)} \ .\end{aligned} \tag{C.14}$$

Again for finite time detections the constant terms do contribute but in the limit as the time of detection $\tau-\tau_0$ goes to infinity these contributions vanish. Further as the constant terms are infinite it seems sensible only to consider the rate of a smoothly switched detector in this asymptotic limit. The remaining transition rate is then

$$\begin{aligned}
\dot{F}_{\Delta\tau}(\omega) &= -\frac{1}{4\pi}\int_{-\infty}^{\infty} ds\, e^{-i\omega s-\frac{|s|}{\Delta\tau}}\ln((s-2i\epsilon)^2)\ , \\
&= -\frac{1}{4\pi}\left(\int_0^{\infty} ds\, e^{-i\omega s-\frac{s}{\Delta\tau}}\ln((s-2i\epsilon)^2)+\int_0^{\infty} ds\, e^{i\omega s-\frac{s}{\Delta\tau}}\ln((s+2i\epsilon)^2)\right)\ , \\
&= -\frac{1}{4\pi}\left[\frac{2\ln(2i\epsilon)}{i\omega+\frac{1}{\Delta\tau}}-\frac{2\ln(2i\epsilon)}{i\omega-\frac{1}{\Delta\tau}}+\frac{2}{i\omega+\frac{1}{\Delta\tau}}\int_0^{\infty} ds\,\frac{e^{-i\omega s-\frac{s}{\Delta\tau}}}{(s-2i\epsilon)}\right. \\
&\quad \left. -\frac{2}{i\omega-\frac{1}{\Delta\tau}}\int_0^{\infty} ds\,\frac{e^{i\omega s-\frac{s}{\Delta\tau}}}{(s+2i\epsilon)}\right]\ , \qquad (C.15)
\end{aligned}$$

where the last line follows by parts. As we are only interested in the asymptotic response we now let $\Delta\tau\to\infty$. Then

$$\dot{F}_{\tau}(\omega) = -\frac{1}{2\pi i\omega}\int_{-\infty}^{\infty} ds\,\frac{e^{-i\omega s}}{(s-2i\epsilon)}\ . \qquad (C.16)$$

The integral may be done simply by residues closing the contour in the upper half complex plane for $\omega<0$ and the lower half plane for $\omega>0$. The result is

$$\dot{F}_{\tau}(\omega) = -\frac{1}{\omega}\Theta(-\omega)\ . \qquad (C.17)$$

Finally we consider the relation with the local density of states (5.41) which in two dimensions is

$$\begin{aligned}
\rho_M(\omega,x) &= \int_{-\infty}^{\infty}\frac{dk}{2\pi}\,\delta(|k|-\omega) = \int_0^{\infty}\frac{dk}{2\pi}\,\delta(k-\omega)+\int_{-\infty}^{0}\frac{dk}{2\pi}\,\delta(-k-\omega)\ , \\
&= \frac{1}{\pi}\Theta(\omega)\ . \qquad (C.18)
\end{aligned}$$

Thus

$$\dot{F}_{\tau}(\omega) = -\frac{\pi}{\omega}\Theta(-\omega)\rho_M(|\omega|,x)\ . \qquad (C.19)$$

C.3 Uniformly accelerated detector

Now let us consider the response of a uniformly accelerated detector in 2 dimensions. Again due to the infinite contributions from the constant terms in the correlation function we consider only the asymptotic limit of a smoothly switched detector. The transition rate, dropping the constant terms which do not contribute in the asymptotic limit, is

$$\begin{aligned}\dot{F}_{\Delta\tau}(\omega) &= -\frac{1}{4\pi}\int_{-\infty}^{\infty} ds\, e^{-i\omega s-\frac{|s|}{\Delta\tau}} \ln\left(4\left(\alpha\sinh\left(\frac{s}{2\alpha}\right)-i\epsilon\cosh\left(\frac{s}{2\alpha}\right)\right)^2\right) \ , \\ &= -\frac{1}{4\pi}\left[\int_{0}^{\infty} ds\, e^{-i\omega s-\frac{s}{\Delta\tau}} \ln\left(4\left(\alpha\sinh\left(\frac{s}{2\alpha}\right)-i\epsilon\cosh\left(\frac{s}{2\alpha}\right)\right)^2\right)\right. \\ &\quad \left.+\int_{0}^{\infty} ds\, e^{i\omega s-\frac{s}{\Delta\tau}} \ln\left(4\left(\alpha\sinh\left(\frac{-s}{2\alpha}\right)-i\epsilon\cosh\left(\frac{s}{2\alpha}\right)\right)^2\right)\right] \ . \end{aligned} \tag{C.20}$$

Integrating by parts twice and then taking the $\Delta\tau \to \infty$ limit we find

$$\dot{F}_{\tau}(\omega) = -\frac{(\alpha^2+\epsilon^2)}{8\pi\omega^2\alpha^2}\int_{-\infty}^{\infty} ds\, \frac{e^{-i\omega s}}{\left(\alpha\sinh\left(\frac{s}{2\alpha}\right)-i\epsilon\cosh\left(\frac{s}{2\alpha}\right)\right)^2} \ . \tag{C.21}$$

Comparing (C.21) with (5.52) when $\Delta\tau \to \infty$ we see that the response is proportional to the response of a uniformly accelerated detector in 4-dimensional Minkowski space. In 4 dimensions we have already recovered the usual thermal response (5.53) thus in 2 dimensions the response is

$$\dot{F}_{\tau}(\omega) = -\frac{1}{\omega(e^{2\pi\alpha\omega}-1)} \ . \tag{C.22}$$

Appendix D

Integrals on uniformly accelerated worldline

In this appendix we evaluate the integral

$$I(\omega) = \lim_{\epsilon\to 0^+} \int_{-\infty}^{\infty} ds \, \frac{e^{-i\omega s}}{(\alpha\sinh(\beta s) - i\epsilon\cosh(\beta s))^k} \ , \tag{D.1}$$

for arbitrary $\alpha > 0$, $\beta > 0$, $\omega \in \mathbb{R}$ and $k \in \mathbb{Z}^+$. Writing $s = z/\beta$, $\epsilon = \alpha\eta$, $q = \omega/\beta$, we may instead consider

$$I(q) = \lim_{\eta\to 0^+} \frac{1}{\beta\alpha^k} \int_{-\infty}^{\infty} dz \, \frac{e^{-iqz}}{(\sinh(z) - i\eta\cosh(z))^k} \ . \tag{D.2}$$

The integrand has a pole at $z = \operatorname{arctanh}(i\eta)$ of order k. We consider the integral as a contour integral and deform the contour to that shown in figure D.1. Now we are free to take $\lim_{\eta\to 0^+}$. Further we may easily show that in the limit as the contour goes off to $\pm\infty$ the contributions from $R1$ and $R2$ vanish and so

$$I(q) = \frac{1}{\beta\alpha^k} \int_C dz \, \frac{e^{-iqz}}{(\sinh(z))^k} \ , \tag{D.3}$$



Figure D.1: Contour used in the integration of $I(q)$.

where C is the line $z = -i\pi/2$ in the complex z-plane. Changing the integration variable by $z = W - i\pi/2$ gives

$$\begin{aligned} I(q) &= \frac{e^{-\pi q/2}}{\beta\alpha^k(-i)^k}\int_{-\infty}^{\infty} dW\,\frac{e^{-iqW}}{(\cosh(W))^k}\ , \\ &= \frac{e^{-\pi q/2}}{\beta\alpha^k(-i)^k}\left(\int_{-\infty}^{\infty} dW\,\frac{\cos(qW)}{(\cosh(W))^k} - i\int_{-\infty}^{\infty} dW\,\frac{\sin(qW)}{(\cosh(W))^k}\right)\ . \end{aligned} \tag{D.4}$$

In the second integral the integrand is odd in W and so the integral is 0. The first integral may be found in [50], with the result

$$I(q) = \frac{2^{k-1}e^{-\pi q/2}}{\Gamma(k)\beta\alpha^k(-i)^k}\Gamma((k+iq)/2)\Gamma((k-iq)/2)\ . \tag{D.5}$$

We may further use the properties of Γ functions [50],

$$\begin{aligned} \Gamma(z+1) &= z\Gamma(z)\ , \\ \Gamma(1-z)\Gamma(z) &= \frac{\pi}{\sin\pi z}\ , \\ \Gamma(1/2+z)\Gamma(1/2-z) &= \frac{\pi}{\cos\pi z}\ , \end{aligned} \tag{D.6}$$

to write the Γ functions that enter the transition rate (5.52) in terms of products of polynomials to obtain the expression (5.53).

Appendix E

Details for automorphic field detector

In this appendix we give some more detail on some statements made in section 5.4.1 regarding automorphic fields.

The automorphic field on Minkowski space, which represents fields on M/Γ is defined by

$$\hat{\phi}(x) := \frac{1}{\left(\sum_{\gamma\in\Gamma} p(\gamma)^2\right)^{1/2}} \sum_{\gamma\in\Gamma} p(\gamma)\phi(\gamma^{-1}x) \ , \tag{E.1}$$

where $p(\gamma)$ is a representation of Γ in $SL(\mathbb{R})$. Then

$$\left[\hat{\phi}(x), \dot{\hat{\phi}}(x')\right] = \frac{1}{\sum_{g\in\Gamma} p(g)^2} \sum_{g\in\Gamma}\sum_{h\in\Gamma} p(g)p(h) \left[\phi(g^{-1}x), \dot{\phi}(h^{-1}x')\right] \ . \tag{E.2}$$

Now writing $h \to hg$ we have, at equal times

$$\begin{aligned}\left[\hat{\phi}(x), \dot{\hat{\phi}}(x')\right] &= \frac{1}{\sum_{g\in\Gamma} p(g)^2} \sum_{g\in\Gamma}\sum_{h\in\Gamma} p(g)p(hg)[\phi(g^{-1}x), \dot{\phi}(g^{-1}h^{-1}x')] \ , \\ &= \frac{1}{\sum_{g\in\Gamma} p(g)^2} \sum_{g\in\Gamma} p(g)^2 \sum_{h\in\Gamma} p(h)[\phi(x), \dot{\phi}(h^{-1}x')] \ , \\ &= \sum_{h\in\Gamma} p(h)[\phi(x), \dot{\phi}(h^{-1}x')] = i\delta^{(d-1)}(x-x') + \text{image terms} \ . \end{aligned} \tag{E.3}$$

Next consider the two-point function $\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle$, where

$$\hat{\phi}(\tau) = \int d^{d-1}\xi \, W_\epsilon(\xi)\hat{\phi}(x(\tau,\xi)) \ . \tag{E.4}$$

With a similar calculation we find

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = \frac{1}{\sum_{g\in\Gamma} p(g)^2} \sum_{g\in\Gamma}\sum_{h\in\Gamma} p(g)p(h)\langle 0|\phi(g^{-1}\tau)\phi(h^{-1}\tau')|0\rangle \ . \tag{E.5}$$

Writing $h \to hg$ and following through the steps which lead to (E.3) we have

$$\langle 0|\hat{\phi}(\tau)\hat{\phi}(\tau')|0\rangle = \sum_{h\in\Gamma} p(h)\langle 0|\phi(\tau)\phi(h^{-1}\tau')|0\rangle \ . \tag{E.6}$$

That is, the two-point function for the smeared automorphic field (E.4) is given by the method of images applied to the two point function for the smeared Minkowski field.

Appendix F

Evaluating (5.95) for inertial detector approaching boundary

In this appendix we evaluate the integral

$$\dot{F}_\tau(\omega) = -\frac{\beta}{2\pi^2}(1-v^2)\lim_{\epsilon\to 0^+}\int_0^\infty ds\, Re\left(\frac{e^{-i\omega s}}{(s-2i\epsilon)^2 - v^2(s-2\tau)^2}\right) \ , \tag{F.1}$$

with $-1 < v < 0$ and $-\infty < \tau < 0$. The integrand has two poles, at $s = (2i\epsilon \pm 2v\tau)/(1 \pm v)$. When the limit is taken only the pole with $(+)$ lies in the integration region. We consider the integration before the limit is taken as a contour integral and we deform the contour as shown in figure F.1. Then we are free to take the limit and obtain

$$\dot{F}_\tau(\omega) = -\frac{\beta}{2\pi^2}\int_C ds\, Re\left(\frac{e^{-i\omega s}}{(s+c)(s-b)}\right) \ , \tag{F.2}$$

where $c = 2v\tau/(1-v)$ and $b = 2v\tau/(1+v)$.

We compute the integral in (F.2) by rotating the contour onto the imaginary s axis. For $\omega > 0$ we rotate to the negative imaginary axis, $s \to -it$. We find after separating the



Figure F.1: Contour used in the integration. The right hand pole is at $s = (2i\epsilon + 2v\tau)/(1+v)$, the left at $s = (2i\epsilon - 2v\tau)/(1-v)$. Recall that $-\infty < \tau < 0$ and $-1 < v < 0$.

integrand by partial fractions

$$\begin{aligned}
\dot{F}_\tau(\omega) &= -\frac{\beta}{2\pi^2(c+b)} \int_0^\infty dt\, Re\left(\frac{e^{-\omega t}}{(t-ib)} - \frac{e^{-\omega t}}{(t+ic)}\right) , \\
&= -\frac{\beta}{2\pi^2(c+b)} \int_0^\infty dt\, \frac{e^{-\omega t} t}{(t^2+b^2)} - \frac{e^{-\omega t} t}{(t^2+c^2)} . \qquad \text{(F.3)}
\end{aligned}$$

The integrals may be computed in terms of sine and cosine integrals [50], giving the result (5.96). For $\omega < 0$ we rotate to the positive imaginary axis $s \rightarrow it$. The method is similar to that above except now a contribution is picked up from a pole on the positive real axis. This contribution is given by $2\pi i \mathrm{Res}(e^{-i\omega s}/((s+c)(s-b)), s = b)$. The residue is easily found as $s = b$ is a simple pole. The result is as shown in (5.96).

Appendix G

Asymptotic behaviour of (5.101) and (5.115)

In this appendix we show that (5.101) of section 5.4.3 consists of a function periodic in τ with period π/ω plus a function $B_\tau(\omega)$ bounded in absolute value by $\frac{\beta\Gamma(d/2-1)}{(d-2)\pi^{d/2}\alpha^{d-3}}e^{-\frac{(d-2)\tau}{\alpha}}$. In particular we show when $d = 4$ (5.101) is given by

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\omega\cos(2\tau\omega)}{2\pi\sinh(\omega\pi\alpha)} + B_\tau(\omega)\ . \tag{G.1}$$

Further we present an analytic proof that when the detector is switched on at a finite τ_0 (5.101) and (5.115) (with the upper limit replaced by $\tau - \tau_0$) vanish as $\tau \to \infty$.

In this appendix we also discuss the behaviour of (5.115) as $\tau \to \infty$. We show that (5.115) also consists of a function periodic in τ with period π/ω plus a function which decays exponentially as $\tau \to \infty$.

Consider (5.101),

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}} \int_0^\infty ds\, \frac{\cos(\omega s)}{(2\alpha\cosh(\frac{2\tau-s}{2\alpha}))^{d-2}}\ . \tag{G.2}$$

Changing the integration variables by $s = 2\alpha x + 2\tau$ gives

$$\begin{aligned}
\dot{F}_{B\tau}(\omega) &= \frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}(2\alpha)^{d-3}} \int_{-\tau/\alpha}^{\infty} dx\, \frac{\cos(\omega(2\alpha x + 2\tau))}{\cosh^{d-2} x} \ , \\
&= \frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}(2\alpha)^{d-3}} \left(\int_{-\infty}^{\infty} dx\, \frac{\cos(\omega(2\alpha x + 2\tau))}{\cosh^{d-2} x} - \int_{-\infty}^{-\tau/\alpha} dx\, \frac{\cos(\omega(2\alpha x + 2\tau))}{\cosh^{d-2} x} \right) \ , \\
&= \frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}(2\alpha)^{d-3}} \left(\cos(2\tau\omega) \int_{-\infty}^{\infty} dx\, \frac{\cos(2\alpha\omega x)}{\cosh^{d-2} x} - \sin(2\tau\omega) \int_{-\infty}^{\infty} dx\, \frac{\sin(2\alpha\omega x)}{\cosh^{d-2} x} \right. \\
&\quad \left. - \int_{-\infty}^{-\tau/\alpha} dx\, \frac{\cos(\omega(2\alpha x + 2\tau))}{\cosh^{d-2} x} \right) \ . \qquad \text{(G.3)}
\end{aligned}$$

In the last expression the integrand in the second term is odd in x and hence integrates to 0. The first term is periodic in τ with period π/ω, further the integral may be performed in terms of Gamma functions [50]. When $d = 4$ the result is

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\omega\cos(2\tau\omega)}{2\pi\sinh(\omega\pi\alpha)} - \frac{\beta}{4\pi^2\alpha} \int_{-\infty}^{-\tau/\alpha} dx\, \frac{\cos(\omega(2\alpha x + 2\tau))}{\cosh^2 x} \ . \qquad \text{(G.4)}$$

Writing $y = -x$, we find that the third term in (G.3) is bounded in absolute value by

$$\begin{aligned}
\frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}(2\alpha)^{d-3}} \int_{\tau/\alpha}^{\infty} dy\, \frac{1}{\cosh^{d-2} y} &\leq \frac{\beta\Gamma(d/2-1)}{\pi^{d/2}\alpha^{d-3}} \int_{\tau/\alpha}^{\infty} dy\, \frac{1}{e^{(d-2)y}} \ , \\
&= \frac{\beta\Gamma(d/2-1)}{(d-2)\pi^{d/2}\alpha^{d-3}} e^{-\frac{(d-2)\tau}{\alpha}} \ . \qquad \text{(G.5)}
\end{aligned}$$

Next consider (5.101) for the detector switched on instantaneously at finite τ_0,

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}} \int_{0}^{\tau-\tau_0} ds\, \frac{\cos(\omega s)}{(2\alpha\cosh(\frac{2\tau-s}{2\alpha}))^{d-2}} \ . \qquad \text{(G.6)}$$

We consider the behaviour as $\tau \to \infty$ for fixed τ_0.

$$\dot{F}_{B\tau}(\omega) = \frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}} \left(\int_{-\infty}^{\tau-\tau_0} ds\, \frac{\cos(\omega s)}{(2\alpha\cosh(\frac{2\tau-s}{2\alpha}))^{d-2}} - \int_{-\infty}^{0} ds\, \frac{\cos(\omega s)}{(2\alpha\cosh(\frac{2\tau-s}{2\alpha}))^{d-2}} \right) \ . \qquad \text{(G.7)}$$

Changing variables by $s=-2\alpha y+2\tau$ we have seen in (G.5) that the second term is bounded in absolute value by $\frac{\beta\Gamma(d/2-1)}{(d-2)\pi^{d/2}\alpha^{d-3}}e^{-\frac{(d-2)\tau}{\alpha}}$. In the first term in (G.7) we change variables by $s=\tau-\tau_0-v$. The result is

$$\begin{aligned}\frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}}\int_0^\infty dv\,\frac{\cos(\omega(\tau-\tau_0-v))}{(2\alpha\cosh(\frac{\tau+\tau_0+v}{2\alpha}))^{d-2}} &\leq \frac{\beta\Gamma(d/2-1)}{2\pi^{d/2}}\int_0^\infty dv\,\frac{1}{(\alpha e^{\frac{\tau+\tau_0+v}{2\alpha}})^{d-2}}\ , \\ &= \frac{\beta\Gamma(d/2-1)}{(d-2)\pi^{d/2}\alpha^{d-3}}e^{-\frac{(d-2)(\tau+\tau_0)}{2\alpha}}\ . \end{aligned} \quad \text{(G.8)}$$

This term is therefore also bounded by a function which exponentially decays as $\tau\to\infty$. Hence, for the detector switched on instantaneously at finite τ_0, $\dot{F}_{B\tau}(\omega)\to 0$ as $\tau\to\infty$.

Consider now (5.115), that is

$$\begin{aligned}\dot{F}_{I\tau}(\omega) &= \frac{1}{2\pi^2}\sum_{n=-\infty}^{\infty}\int_0^\infty ds\,\frac{\cos(\omega s)}{4\alpha^2\cosh^2\left(\frac{2\tau-s}{2\alpha}\right)+4y_0^2+a^2(2n-1)^2}\ , \\ &= \frac{1}{2\pi^2}\sum_{n=-\infty}^{\infty}\left(\int_{-\infty}^\infty ds\ -\int_{-\infty}^0 ds\right)\frac{\cos(\omega s)}{4\alpha^2\cosh^2\left(\frac{2\tau-s}{2\alpha}\right)+4y_0^2+a^2(2n-1)^2}\ .\end{aligned} \quad \text{(G.9)}$$

Changing integration variables by $s=2\alpha x+2\tau$ we find the first term in (G.9) becomes

$$\frac{2\alpha}{2\pi^2}\cos(2\tau\omega)\sum_{n=-\infty}^{\infty}\int_{-\infty}^\infty dx\,\frac{\cos(2\alpha\omega x)}{4\alpha^2\cosh^2 x+y_0^2+a^2(2n-1)^2}\ , \quad \text{(G.10)}$$

which is periodic in τ with period π/ω. Changing coordinates by $s=-2\alpha z$ the second term in (G.9) becomes

$$I(\omega)=-\frac{1}{2\alpha\pi^2}\sum_{n=1}^{\infty}\int_0^\infty dz\,\frac{\cos(2\alpha\omega z)}{\cosh^2\left(z+\frac{\tau}{\alpha}\right)+\left(\frac{y_0}{\alpha}\right)^2+\left(\frac{a}{2\alpha}\right)^2(2n-1)^2}\ . \quad \text{(G.11)}$$

Thus

$$
\begin{aligned}
|I(\omega)| \quad \leq \quad & \frac{1}{2\alpha\pi^2}\left(\int_0^\infty dz\,\frac{1}{\cosh^2\left(z+\frac{\tau}{\alpha}\right)+\left(\frac{y_0}{\alpha}\right)^2+\left(\frac{a}{2\alpha}\right)^2}\right. \\
& \left. +\sum_{n=2}^{\infty}\int_0^\infty dz\,\frac{1}{\cosh^2\left(z+\frac{\tau}{\alpha}\right)+\left(\frac{y_0}{\alpha}\right)^2+\left(\frac{a}{2\alpha}\right)^2(2n-1)^2}\right) \ . \qquad \text{(G.12)}
\end{aligned}
$$

It is simple to show that the first term is (G.12) is bounded in absolute value by $\frac{1}{\alpha\pi^2}e^{-2\tau/\alpha}$. The integrand in the second term of (G.12), call it $I_2(\omega)$, is a monotonically decreasing function of n. We can therefore bound the sum by an integral

$$
\begin{aligned}
I_2(\omega) \quad & \leq \quad \frac{1}{2\alpha\pi^2}\int_0^\infty dz\int_1^\infty dn\,\frac{1}{\cosh^2\left(z+\frac{\tau}{\alpha}\right)+\left(\frac{y_0}{\alpha}\right)^2+\left(\frac{a}{2\alpha}\right)^2(2n-1)^2} \ , \\
& \leq \quad \frac{1}{2\alpha\pi^2}\int_0^\infty dz\int_{1/2}^\infty dn\,\frac{1}{\cosh^2\left(z+\frac{\tau}{\alpha}\right)+\left(\frac{y_0}{\alpha}\right)^2+\left(\frac{a}{2\alpha}\right)^2(2n-1)^2} \ . \qquad \text{(G.13)}
\end{aligned}
$$

Changing variables by $n=\frac{1}{2}+\frac{\alpha}{a}v$

$$
\begin{aligned}
I_2(\omega) \quad & \leq \quad \frac{1}{2a\pi^2}\int_0^\infty dz\int_0^\infty dv\,\frac{1}{\cosh^2\left(z+\frac{\tau}{\alpha}\right)+\left(\frac{y_0}{\alpha}\right)^2+v^2} \ , \\
& = \quad \frac{1}{4a\pi}\int_0^\infty dz\,\frac{1}{\left(\cosh^2\left(z+\frac{\tau}{\alpha}\right)+\left(\frac{y_0}{\alpha}\right)^2\right)^{1/2}} \ , \\
& \leq \quad \frac{1}{2a\pi}\int_0^\infty dz\,e^{-(z+\tau/\alpha)}=\frac{1}{2a\pi}e^{-\tau/\alpha} \ . \qquad \text{(G.14)}
\end{aligned}
$$

Thus we see that $I(\omega)\leq\frac{1}{\alpha\pi^2}e^{-2\tau/\alpha}+\frac{1}{2\pi a}e^{-\tau/\alpha}$ and so $I(\omega)$ vanishes as $\tau\to\infty$ and (5.115) becomes periodic in τ with period π/ω at late times.

Finally consider (5.115) when the detector is switched on at $\tau_0>-\infty$, that is

$$
\begin{aligned}
\dot{F}_{I\tau}(\omega) \quad & = \quad \frac{1}{2\pi^2}\sum_{n=-\infty}^{\infty}\int_0^{\tau-\tau_0}ds\,\frac{\cos(\omega s)}{4\alpha^2\cosh^2\left(\frac{2\tau-s}{2\alpha}\right)+4y_0^2+a^2(2n-1)^2} \ , \\
& = \quad \frac{1}{2\pi^2}\sum_{n=-\infty}^{\infty}\left(\int_{-\infty}^{\tau-\tau_0}ds-\int_{-\infty}^{0}ds\right)\frac{\cos(\omega s)}{4\alpha^2\cosh^2\left(\frac{2\tau-s}{2\alpha}\right)+4y_0^2+a^2(2n-1)^2} \ . \qquad \text{(G.15)}
\end{aligned}
$$

We consider the limit as $\tau \to \infty$ for fixed τ_0. We saw in the calculation (G.11) to (G.14) that the second term in (G.15) is bounded in absolute value by $\frac{1}{\alpha\pi^2}e^{-2\tau/\alpha} + \frac{1}{2\pi a}e^{-\tau/\alpha}$. Consider then the first term in (G.15). With a change of variables by $s = \tau - \tau_0 - v$ this term becomes

$$\begin{aligned} I(\omega) &= \frac{1}{2\pi^2}\sum_{n=-\infty}^{\infty}\int_0^\infty dv\, \frac{\cos(\omega(\tau-\tau_0-v))}{4\alpha^2\cosh^2\left(\frac{\tau+\tau_0+v}{2\alpha}\right) + 4y_0^2 + a^2(2n-1)^2}\ , \\ &= \frac{1}{4\pi^2\alpha^2}\sum_{n=1}^{\infty}\int_0^\infty dv\, \frac{\cos(\omega(\tau-\tau_0-v))}{\cosh^2\left(\frac{\tau+\tau_0+v}{2\alpha}\right) + \left(\frac{y_0}{\alpha}\right)^2 + \left(\frac{a}{2\alpha}\right)^2(2n-1)^2}\ , \\ &= \frac{1}{4\pi^2\alpha^2}\left[\int_0^\infty dv\, \frac{\cos(\omega(\tau-\tau_0-v))}{\cosh^2\left(\frac{\tau+\tau_0+v}{2\alpha}\right) + \left(\frac{y_0}{\alpha}\right)^2 + \left(\frac{a}{2\alpha}\right)^2}\right. \\ &\quad \left. + \sum_{n=2}^{\infty}\int_0^\infty dv\, \frac{\cos(\omega(\tau-\tau_0-v))}{\cosh^2\left(\frac{\tau+\tau_0+v}{2\alpha}\right) + \left(\frac{y_0}{\alpha}\right)^2 + \left(\frac{a}{2\alpha}\right)^2(2n-1)^2}\right]\ . \end{aligned} \tag{G.16}$$

It is a simple matter to show the first term in (G.16) is bounded in absolute value by $\frac{e^{-\frac{\tau+\tau_0}{\alpha}}}{\pi^2\alpha}$. In the second term in (G.16), call it $I_2(\omega)$, the integrand is a monotonically decreasing function of n. The sum can be bounded by an integral and we find

$$|I_2(\omega)| \le \frac{1}{4\pi^2\alpha^2}\int_0^\infty dv \int_{1/2}^\infty dn\, \frac{\cos(\omega(\tau-\tau_0-v))}{\cosh^2\left(\frac{\tau+\tau_0+v}{2\alpha}\right) + \left(\frac{y_0}{\alpha}\right)^2 + \left(\frac{a}{2\alpha}\right)^2(2n-1)^2}\ . \tag{G.17}$$

Changing variables by $n = \frac{1}{2} + \frac{\alpha}{a}z$

$$\begin{aligned} |I_2(\omega)| &\le \frac{1}{4\pi^2\alpha a}\int_0^\infty dv \int_0^\infty dz\, \frac{1}{\cosh^2\left(\frac{\tau+\tau_0+v}{2\alpha}\right) + \left(\frac{y_0}{\alpha}\right)^2 + z^2}\ , \\ &= \frac{1}{8\pi\alpha a}\int_0^\infty dv\, \frac{1}{\left(\cosh^2\left(\frac{\tau+\tau_0+v}{2\alpha}\right) + \left(\frac{y_0}{\alpha}\right)^2\right)^{1/2}}\ , \\ &\le \frac{1}{4\pi\alpha a}\int_0^\infty dv\, e^{-\frac{\tau+\tau_0+v}{2\alpha}}\ , \\ &= \frac{1}{2\pi a}e^{-\frac{\tau+\tau_0}{2\alpha}}\ . \end{aligned} \tag{G.18}$$

Thus finally we see that $|I_2(\omega)| \to 0$ as $\tau \to \infty$, and so (5.115) also vanishes in this limit.

Bibliography

[1] N. D. Birrell and P. C. W. Davies, *"Quantum Fields in Curved Space"* Cambridge University Press (Cambridge, 1982)

[2] S. A. Fulling, *"Aspects of Quantum Field Theory in Curved Spacetime"* Cambridge University Press (Cambridge, 1989)

[3] R. M. Wald, *"Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics"* The University of Chicago Press (Chicago, 1994)

[4] L. Parker, *"The Creation of Particles in an Expanding Universe"* PhD Thesis, (Harvard University, 1966)

[5] S. W. Hawking, Commun.Math.Phys **43**, 199 (1975)

[6] J. D. Bekenstein, Phys.Rev.D **7**, 2333 (1973)

[7] R. Kaul and P. Majumdar, Phys.Rev.Lett **84**, 5255 (2000)

[8] S. Das, P. Majumdar and R. K. Bhaduri, Class.Quant.Grav **19**, 2355 (2002)

[9] S. Das, Pramana **63**, 797 (2004)

[10] M. D. Kruskal, Phys.Rev **119**, 1743 (1960)

[11] G. Szekeres, Publ.Mat.Debrecen **7**, 285 (1960)

[12] J. B. Hartle and S. W. Hawking, Phys.Rev.D **13**, 2188 (1976)

[13] W. Israel, Phys.Lett **A57**, 107 (1976)

[14] W. G. Unruh, Phys.Rev.D **14**, 870 (1976)

[15] G. W. Gibbons and S. W. Hawking, Phys.Rev.D **15**, 2752 (1977)

[16] B. S. Kay and R. M. Wald, Phys.Rep **207**, 49 (1991)

[17] S. A. Fulling, Phys.Rev.D **7**, 2850 (1973)

[18] P. Chen and T. Tajima, Phys.Rev.Lett **83**, 256 (1999)

[19] J. L. Friedman, K. Schleich and D. M. Witt, Phys.Rev.Lett **71**, 1486 (1993): Erratum, Phys.Rev.Lett. **75**, 1872 (1995)

[20] C. W. Misner and J. A. Wheeler, Ann.Phys **2**, 525 (1957)

[21] D. Giulini, Ph.D. thesis (University of Cambridge, 1989)

[22] J. Louko and D. Marolf, Phys.Rev.D **58**, 024007 (1998)

[23] P. Langlois, Phys.Rev.D **70**, 104008 (2004)

[24] P. Langlois, Class.Quant.Grav **22**, 4141 (2005)

[25] S. Schlicht, Class.Quant.Grav **21**, 4647 (2004)

[26] S. Schlicht, PhD Thesis (University of Freiburg, 2002)

[27] B. S. DeWitt, *"Quantum gravity: the new synthesis" in "General Relativity an Einstein centenary survey"* ed S. W. Hawking, W. Israel, Cambridge University Press (Cambridge, 1979)

[28] S. Deser and O. Levin, Phys.Rev.D **59**, 064004 (1999)

[29] P. Langlois, gr-qc/0510049

[30] P. Candelas and D. Deutsch, Proc.R.Soc.Lond.A **362**, 251 (1978)

[31] S. Takagi, Prog.Theor.Phys.Supp **88**, 1 (1986)

[32] M. Soffel, B. Muller and W. Greiner, Phys.Rev.D **22**, 1935 (1979)

[33] D. Oriti, Nuovo.Cim **B115**, 1005 (2000)

[34] D. Oriti Masters thesis (unpublished)

[35] R. Banach and J. S. Dowker, J.Phys.A:Math.Gen **12**, 2527 (1979)

[36] R. Banach and J. S. Dowker, J.Phys.A:Math.Gen **12**, 2545 (1979)

[37] S. J. Avis and C. J. Isham in *"Recent Developments in Gravitation-Cargese 1978"* ed M Levy and Deser, Plenum Press (New York, 1979)

[38] J. D. Bjorken and S. D. Drell, *"Relativistic Quantum Mechanics"* McGraw-Hill Book Company (1964)

[39] C. Itzykson and J. B. Zuber, *"Quantum Field Theory"* McGraw-Hill Book Company (New York, 1975)

[40] D. Zwillinger, *"CRC Standard Mathematical Tables and Formulae"* Chapman and Hall/CRC Press LLC (2003)

[41] A. Hatcher, *"Algebraic Topology"* Cambridge University Press (Cambridge, 2002), Proposition 1.40 and Exercise 1.3.24.

[42] D. G. Boulware, Phys.Rev.D **12**, 350 (1975)

[43] D. R. Brill and J. A. Wheeler, Rev.Mod.Phys **29**, 465 (1957)

[44] S. Chandrasekhar in *"The Mathematical Theory of Black Holes"* Oxford University Press (New York, 1985)

[45] A. Messiah, *"Quantum Mechanics"* North-Holland (Amsterdam, 1991), Vol.I.

[46] T. Tanaka and W. A. Hiscock, Phys.Rev.D **52**, 4503 (1995)

[47] B. S. Dewitt, C. F. Hart and C. J. Isham, Physica **96A**, 197 (1979)

[48] N. Ahmadi and M. Nouri-Zonoz, Phys.Rev.D **71**, 104012 (2005)

[49] S. M. Christensen, Phys.Rev.D **14**, 2490 (1976)

[50] I. S. Gradshteyn and I. M. Ryzhik, *"Table of Integrals Series and Products"* Academic Press (2000)

[51] J. D. Bjorken and S. D. Drell, *"Relativistic Quantum Fields"* McGraw-Hill Book Company (1965)

[52] J. R. Letaw, Phys.Rev.D **23**, 1709 (1981)

[53] J. S. Dowker, Phys.Rev.D **36**, 3095 (1987)

[54] P. Candelas and D. W. Sciama, Phys.Rev.Lett **38**, 1372 (1977)

[55] D. W. Sciama, P. Candelas and D. Deutsch, Adv.Phys **30**, 327 (1981)

[56] S. Takagi, Prog.Theor.Phys **72**, 505 (1984)

[57] S. Takagi, Prog.Theor.Phys **74**, 142 (1985)

[58] W. G. Unruh, Phys.Rev.D **34**, 1222 (1986)

[59] S. Takagi, Prog.Theor.Phys **74**, 501 (1985)

[60] R. Kubo, J.Phys.Soc.Jpn **12**, 570 (1957)

[61] P. C. Martin and J. Schwinger, Phys.Rev **115**, 1342 (1959)

[62] J. R. Letaw and J. D. Pfautsch, Phys.Rev.D **22**, 1345 (1980)

[63] P. C. W. Davies, T. Dray and C. A. Manogue, Phys.Rev.D **53**, 4382 (1996)

[64] O. Levin, Y. Peleg and A. Peres, J.Phys.A **26**, 3001 (1993)

[65] N. Nicolaevici, Class.Quant.Grav **18**, 5407 (2001)

[66] A. C. Ottewill and E. Winstanley, Phys.Rev.D **62**, 084018 (2000)

[67] G. Duffy and A. C. Ottewill gr-qc/0507116v1

[68] L. Sriramkumar and T. Padmanabhan, Int.J.Mod.Phys **D11**, 1 (2002)

[69] V. A. De Lorenci, R. D. M. De Paola and N. F. Svaiter, Class.Quant.Grav **17**, 4241 (2000) Erratum, Class.Quant.Grav **18**, 205 (2001)

[70] V. A. De Lorenci and N. F. Svaiter, Found.Phys. **29**,1233 (1999)

[71] R. D. M. De Paola and N. F. Svaiter, Class.Quant.Grav **18**, 1799 (2001)

[72] K. J. Hinton, J.Phys.A **16**, 1937 (1983)

[73] K. J. Hinton, Class.Quant.Grav **1**, 27 (1984)

[74] L. Sriramkumar, Mod.Phys.Lett **A17**, 1059 (2002)

[75] L. Sriramkumar, Gen.rel.Grav **35**, 1699 (2003)

[76] T. Padmanabhan and T. P. Singh, Class.Quant.Grav **4**, 1397 (1987)

[77] L. H. Ford and T. A. Roman, Phys.Rev.D **48**, 776 (1993)

[78] P. G. Grove and A. C. Ottewill, J.Phys.A **16**, 3905 (1983)

[79] B. F Svaiter and N. F. Svaiter, Phys.Rev.D **46**, 5267 (1992) Erratum, Phys.Rev.D **47**, 4802 (1993)

[80] P. G. Grove, Class.Quant.Grav **5**, 1381 (1988)

[81] A. Higuchi, G. E. A. Matsas and C. B. Peres, Phys.Rev.D **48**, 3731 (1993)

[82] L. Sriramkumar and T. Padmanabhan, Class.Quant.Grav **13**, 2061 (1996)

[83] P. Martinetti and C. Rovelli, Class.Quant.Grav **20**, 4919 (2003)

[84] P. Martinetti, gr-qc/0401116

[85] Y. Tian, J.High Energy.Phys JHEP06(2005)045, (2005)

[86] T. Jacobson, gr-qc/0308048

[87] D. H. Coule, Class.Quant.Grav **22**, R125 (2005)

[88] L. H. Ford, Phys.Rev.D **43**, 3972 (1991)

[89] L. H. Ford and T. A. Roman, Phys.Rev.D **55**, 2082 (1997)

[90] M. J. Pfenning and L. H. Ford, Phys.Rev.D **55**, 4813 (1997)

[91] M. J. Pfenning and L. H. Ford, Phys.Rev.D **57**, 3489 (1998)

[92] L. H. Ford, M. J. Pfenning and T. A. Roman, Phys.Rev.D **57**, 4839 (1998)

[93] C. J. Fewster and S. P. Eveson, Phys.Rev.D **58**, 084010 (1998)

[94] C. J. Fewster and E. Teo, Phys.Rev.D **59**, 104016 (1999)

[95] C. J. Fewster, Class.Quant.Grav **17**, 1897 (2000)

[96] L. H. Ford, N. F. Svaiter and M. L. Lyra, Phys.Rev.A **49**, 1378 (1994)

[97] P. C. W. Davies and A. C. Ottewill, Phys.Rev.D **65**, 104014 (2002)

[98] A. Vilenkin, Phys.Rev.D **23**, 852 (1981)

[99] A. Chamblin and G. W. Gibbons, Phys.Rev.D **55**, 2177 (1997)

[100] J. Louko and K. Schleich, Class.Quant.Grav **16**, 2005 (1999)

[101] B. McInnes, J.High Energy.Phys JHEP 09(2003)009, (2003)

[102] W. Greiner and J. Reinhardt, *"Field Quantization"* Springer (Berlin, 1996)

[103] A. S. Wightman, *in "1964 Cargese Lectures in Theoretical Physics"* ed M. Levy, Gordon and Breach (1967)

[104] C. W. Misner, K. S. Thorne and J. A. Wheeler, *"Gravitation"* Freeman (New York 1973)

[105] A. Ottewill and S. Takagi, Prog.Theor.Phys **77**, 310 (1987)

[106] M. Abramowitz and I. A. Stegun, *"Handbook of Mathematical Functions"* Dover Publications, Inc (New York, 1970)

[107] P. C. W. Davies, Z. X. Liu and A. C. Ottewill, Class.Quant.Grav **6**, 1041 (1989)

[108] N. Suzuki, Class.Quant.Grav **14**, 3149 (1997)

[109] K. Schleich and D. M. Witt Nucl.Phys.B **402**, 411 (1993)

[110] K. Schleich and D. M. Witt Nucl.Phys.B **402**, 469 (1993)

[111] R. P. Geroch and J. Traschen, Phys.Rev.D **36**, 1017 (1987)

[112] D. Garfinkle, Class.Quant.Grav **16**, 4101 (1999)

[113] M. R. Anderson, *"The mathematical theory of cosmic strings"* IOP Publishing (2003)

[114] B. F. Svaiter and N. F. Svaiter, Class.Quant.Grav **11**, 347 (1994)

[115] P. C. W. Davies and V. Sahni, Class.Quant.Grav **5**, 1 (1988)

[116] S. Deser and O. Levin, Class.Quant.Grav **14**, L163 (1997)

[117] S. Deser and O. Levin, Class.Quant.Grav **15**, L85 (1998)

[118] H. F Goenner in *"General Relativity and Gravitation"* Plenum Press (New York 1980)

[119] N. L. Santos, O. J. C. Dias and J. P. S. Lemos, Phys.Rev.D **70**, 124033 (2004)

[120] Y. Kim, Y. Park and K. Soh, Phys.Rev.D **62**, 104020 (2000)

[121] S. Hong, Y. Kim, and Y. Park, Phys.Rev.D **62**, 024024 (2000)

[122] S. Hong, W. Kim, Y. Kim, and Y. Park, Phys.Rev.D **62**, 064021 (2000)

[123] S. Hong, W. Kim, J. Oh and Y. Park, Phys.Rev.D **63**, 127502 (2001)

[124] S. Hong, W. Kim, J. Oh and Y. Park, J.Korean.Phys.Soc **42**, 23 (2003)

[125] H. Chen and Y. Tian, Phys.Rev.D **71**, 104008 (2005)

[126] C. Fronsdal, Phys.Rev **116**, 778 (1959)

[127] B. Allen, Phys.Rev.D **32**, 3136 (1985)

[128] A. Ashtekar, J. Baez, A. Corichi and K. Krasnov, Phys.Rev.Lett **80**, 904 (1998)

[129] N. Nakanishi, Prog.Theor.Phys **57**, 269 (1977)